

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
**WO 01/72956 A2**


- (51) International Patent Classification<sup>7</sup>: **C12N**
- (21) International Application Number: **PCT/IB01/01400**
- (22) International Filing Date: **27 March 2001 (27.03.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
**09/535,852** **27 March 2000 (27.03.2000)** **US**
- (71) Applicant (for all designated States except US): **ADHEREX TECHNOLOGIES INC. [CA/CA]**; Suite 340, 600 Peter Morand Crescent, Ottawa, Ontario K1G 5Z3 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BLASCHUK, Orest, W. [CA/CA]**; Suite 1520, 4998 De Maisonneuve

West, Westmount, Quebec H3Z 1N2 (CA). **SYMONDS, James, Matthew [CA/CA]**; Apartment 1, 560 McLeod Street, Ottawa, Ontario K1R 5R3 (CA). **GOUR, Barbara, J. [CA/CA]**; RR#4, 2890 Donnelly Drive, Kemptville, Ontario K0G 1J0 (CA).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **COMPOUNDS AND METHODS FOR MODULATING DESMOSOMAL CADHERIN-MEDIATED FUNCTIONS**



(57) Abstract: Modulating agents for inhibiting or enhancing desmosomal cadherin mediated cell adhesion are provided. The modulating agents comprise one or more of: a) a peptide sequence that is at least 50% identical to a desmosomal cadherin CAR sequence; b) a non-peptide mimetic of a desmosomal cadherin CAR sequence; c) a substance, such as an antibody or antigen-binding fragment thereof, that specifically binds a desmosomal cadherin CAR sequence; and/or d) a polynucleotide encoding a polypeptide that comprises a desmosomal cadherin CAR sequence or analogue thereof. Methods for using such modulating agents for modulating desmosomal cadherin-mediated cell adhesion in a variety of contexts are also provided.

WO 01/72956 A2

BEST AVAILABLE COPY



**Published:**

- without international search report and to be republished upon receipt of that report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## COMPOUNDS AND METHODS FOR MODULATING DESMOSOMAL CADHERIN-MEDIATED FUNCTIONS

### TECHNICAL FIELD

5                   The present invention relates generally to methods for modulating desmosomal cadherin-mediated functions, and more particularly to the use of modulating agents derived from desmocollin and desmoglein cell adhesion recognition sequences for inhibiting or enhancing functions mediated by such desmosomal cadherins.

### 10 BACKGROUND OF THE INVENTION

Cadherins are a rapidly expanding superfamily of calcium-dependent cell adhesion molecules (CAMs) (*for review, see* Munro et al., *In: Cell Adhesion and Invasion in Cancer Metastasis*, P. Brodt, ed., pp. 17-34, RG Landes Co., Austin TX, 1996). All cadherins appear to be membrane glycoproteins that generally promote cell  
15 adhesion through homophilic interactions (a cadherin on the surface of one cell binds to an identical cadherin on the surface of another cell), although cadherins also appear to be capable of forming heterotypic complexes with one another under certain circumstances and with lower affinity.

There are many different types of cadherins. The most extensively  
20 studied group of cadherins is known as the classical, or type I, cadherins. Classical cadherins have been shown to regulate epithelial, endothelial, neural and cancer cell adhesion, with different cadherins expressed on different cell types. All classical cadherins have a similar structure. As illustrated in Figure 1A, classical cadherins are composed of five extracellular domains (EC1-EC5), a single hydrophobic domain (TM)  
25 that transverse the plasma membrane (PM), and two cytoplasmic domains (CP1 and CP2). The calcium binding motifs DXNDN (SEQ ID NO:1), DXD and LDRE (SEQ ID NO:2) are interspersed throughout the extracellular domains, and each 110 amino acid region that contains such motifs is considered a cadherin repeat. The first extracellular domain (EC1) contains the cell adhesion recognition (CAR) sequence, HAV (His-Ala-

---

Val), along with flanking sequences on either side of the CAR sequence that play a role in conferring specificity. Synthetic peptides containing the HAV sequence and antibodies directed against such peptides have been shown to inhibit classical cadherin-dependent processes (Munro et al., *supra*; Blaschuk et al., *J. Mol. Biol.* 211:679-82, 5 1990; Blaschuk et al., *Develop. Biol.* 139:227-29, 1990; Alexander et al., *J. Cell. Physiol.* 156:610-18, 1993).

Cadherins that contain calcium binding motifs within extracellular domain cadherin repeats, but do not contain an HAV CAR sequence, are considered to be nonclassical cadherins (illustrated in Figures 1B to 1AA). To date, nine groups of 10 nonclassical cadherins have been identified (types II - X). These cadherins are also membrane glycoproteins. Type II, or atypical, cadherins include OB-cadherin (cadherin-11; see Getsios et al., *Developmental Dynamics* 211:238-247, 1998; Simonneau et al., *Cell Adhesion and Communication* 3:115-130, 1995; Okazaki et al., *J. Biological Chemistry* 269:12092-12098, 1994), cadherin-5 (VE-cadherin; see Navarro 15 et al., *J. Cell Biology* 140:1475-1484, 1998), cadherin-6 (K-cadherin; see Shimoyama et al., *Cancer Research* 55:2206-2211, 1995; Shimazui et al., *Cancer Research* 56:3234-3237, 1996; Inoue et al., *Developmental Dynamics* 211:338-351, 1998; Getsios et al., *Developmental Dynamics* 211:238-247, 1998), cadherin-7 (see Nakagawa et al., *Development* 121:1321-1332, 1995), cadherin-8 (see Suzuki et al., *Cell Regulation* 20 2:261-270, 1991), cadherin-12 (Br-cadherin; see Tanihara et al., *Cell Adhesion and Communication* 2:15-26, 1994), cadherin-14 (see Shibata et al., *J. Biological Chemistry* 272:5236-5240, 1997), cadherin-15 (M-cadherin; see Shimoyama et al., *J. Biological Chemistry* 273:10011-10018, 1998), and PB-cadherin (see Sugimoto et al., *J. Biological Chemistry* 271:11548-11556, 1996). For a general review of atypical cadherins, see 25 Redies and Takeichi, *Developmental Biology* 180:413-423, 1996 and Suzuki et al., *Cell Regulation* 2:261-270, 1991.

Types III-X include LI-cadherin (type III; see Berndorff et al., *J. Cell Biology* 125:1353-1369, 1994), T-cadherin (type IV; see Ranscht, U.S. Patent No. 5,585,351; Tkachuk et al., *FEBS Lett.* 421:208-212, 1998; Ranscht et al., *Neuron* 7:391- 30 402, 1991; Sacristan et al., *J. Neuroscience Research* 34:664-680, 1993; Vestal and



Ranscht, *J. Cell Biology* 119:451-461, 1992; Fredette and Ranscht, *J. Neuroscience* 14:7331-7346, 1994; Ranscht and Bronner-Fraser, *Development* 111:15-22, 1991), protocadherins (type V; e.g., protocadherins 42, 43 and 68; see Sano et al., *EMBO J.* 12:2249-2256, 1993; GenBank Accession Number AF029343), desmocollins (type VI; 5 e.g., desmocollins 1, 2, 3 and 4; see King et al., *Genomics* 18:185-194, 1993; Parker et al., *J. Biol. Chem.* 266:10438-10445, 1991; King et al., *J. Invest. Dermatol.* 105:314-321, 1995; Kawamura et al., *J. Biol. Chem.* 269:26295-26302, 1994), desmogleins (type VII; e.g., desmogleins 1 and 2; see Wheeler et al., *Proc. Natl. Acad. Sci. USA* 88:4796-4800; Koch et al., *Eur. J. Cell. Biol.* 55:200-208, 1991), and cadherin-related neuronal 10 receptors (type X; see Kohmura et al., *Neuron* 20:1137-1151, 1998).


Most studies of nonclassical cadherins have focused on atypical or type II cadherins. The structure of such cadherins is similar to that of the type I cadherins, but they do not contain the CAR sequence, HAV (Figure 1B). Atypical cadherins appear to mediate a wide variety of functions. Additional variation is seen in the 15 structures of types III-X cadherins. Although less studied, such cadherins also appear to play a role in diverse functions. Desmosomal cadherins, for example, are present within desmosomes, the intercellular junctions that provide adhesion and membrane anchors for the intermediate filament cytoskeleton. These cadherins (e.g., desmogleins and desmocollins) play a role in desmosomal adhesion, which is critically important for 20 normal tissue construction and epidermis structure. Abnormal desmosomes appear in certain types of carcinomas and other skin disorders (see Chidgey, *Histol. Histopathol.* 12:1159-1168, 1997).

Notwithstanding these recent advances, nonclassical cadherin function remains poorly understood at the biological and molecular levels. Accordingly, there is 25 a need in the art for identifying sequences involved in modulating nonclassical cadherin-dependent functions, such as cell adhesion mediated by desmosomal cadherins, and for the development of methods employing such sequences to inhibit processes such as cell adhesion. The present invention fulfills these needs and further provides other related advantages.

## SUMMARY OF THE INVENTION

Briefly stated, this invention provides compositions and methods for modulating desmosomal cadherin-mediated functions, such as cell adhesion. Within certain aspects, modulating agents are provided. Such agents are capable of modulating (i.e., inhibiting or enhancing) one or more functions mediated by the desmosomal cadherins desmoglein (dsg) or desmocollin (dsc). A modulating agent may comprise at least one of: (a) a native dsc or dsg CAR sequence; (b) an analogue of such a CAR sequence that is capable of modulating dsc- or dsg-mediated cell adhesion; (c) a non-peptide peptidomimetic of a dsg or dsc CAR sequence that is capable of modulating dsc- or dsg-mediated cell adhesion; (d) an antibody, or antigen-binding fragment thereof, that specifically binds a dsc or dsg CAR sequence; and/or (e) a polynucleotide encoding a native dsc or dsg CAR sequence or analogue thereof that is capable of modulating dsc- or dsg-mediated cell adhesion. Certain preferred modulating agents comprise a desmoglein-1, desmoglein-2, desmoglein-3, desmocollin-1, desmocollin-2, desmocollin-3 or desmocollin-4 CAR sequence, or an analogue of any of the foregoing CAR sequences. An analogue is generally a peptide sequence that is at least 50% identical to a dsg or dsc CAR sequence, and (i) detectably inhibits a function that is modulated by the dsg or dsc; or (ii) detectably enhances adhesion of cells that express the dsg or dsc. Modulating agents preferably contain no more than 85, and preferably no more than 50, consecutive amino acid residues present within the dsg or dsc. Certain modulating agents comprise a dsg or dsc CAR sequence and are 3-16 amino acid residues in length. Within certain specific embodiments, a modulating agent as described above is a peptide ranging in size from 3 to 50, preferably from 4 to 16, amino acid residues.

For certain modulating agents, the dsg or dsc CAR sequence has the formula:



wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group

consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine or asparagine. For other modulating agents as described above, the dsg or dsc CAR sequence consists of at least three consecutive amino acid residues, and preferably at least five consecutive amino acid residues, of a region of the dsg or dsc having the formula recited above.

Within certain specific embodiments, a modulating agent as provided herein comprises: (a) one or more desmoglein CAR sequences selected from the group consisting of NQK, NQKT (SEQ ID NO:63), NQKTG (SEQ ID NO:64), INQK (SEQ ID NO:65), INQKT (SEQ ID NO:66), INQKTG (SEQ ID NO:67), VINQK (SEQ ID NO:68), VINQKT (SEQ ID NO:69), VINQKTG (SEQ ID NO:70), FVINQK (SEQ ID NO:71), FVINQKT (SEQ ID NO:72), FVINQKTG (SEQ ID NO:73), IFVINQK (SEQ ID NO:74), IFVINQKT (SEQ ID NO:75), IFVINQKTG (SEQ ID NO:76), NRN, NRNT (SEQ ID NO:77), NRNTG (SEQ ID NO:78), INRN (SEQ ID NO:79), INRNT (SEQ ID NO:80), INRNTG (SEQ ID NO:81), IINRN (SEQ ID NO:82), IINRNT (SEQ ID NO:83), IINRNTG (SEQ ID NO:84), FIINRN (SEQ ID NO:85), FIINRNT (SEQ ID NO:86), FIINRNTG (SEQ ID NO:87), MFIINRN (SEQ ID NO:88), MFIINRNT (SEQ ID NO:89), MFIINRNTG (SEQ ID NO:90), NKD, NKDT (SEQ ID NO:91), NKDTG (SEQ ID NO:92), LNKD (SEQ ID NO:93), LNKDT (SEQ ID NO:94), LNKDTG (SEQ ID NO:95), YLNKD (SEQ ID NO:96), YLNKDT (SEQ ID NO:97), YLNKDTG (SEQ ID NO:98), FYLNKD (SEQ ID NO:99), FYLNKDT (SEQ ID NO:100), FYLNKDTG (SEQ ID NO:101), VFYLNKD (SEQ ID NO:102), VFYLNKDT (SEQ ID NO:103) and VFYLNKDTG (SEQ ID NO:104); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmoglein-mediated function is not substantially diminished. For example, the agent may comprise a linear peptide having the sequence N-Ac-IFVINQKTG-NH<sub>2</sub> (SEQ ID NO:105), N-Ac-MFIINRNTG-NH<sub>2</sub> (SEQ ID NO:106) or N-Ac-VFYLNKDTG-NH<sub>2</sub> (SEQ ID NO:107). The desmoglein CAR sequence may, but need not, be present within a cyclic peptide.

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more desmocollin CAR sequences selected from the

group consisting of EKD, EKDT (SEQ ID NO:108), EKDTG (SEQ ID NO:109), IEKD (SEQ ID NO:110), IEKDT (SEQ ID NO:111), IEKDTG (SEQ ID NO:112), YIEKD (SEQ ID NO:113), YIEKDT (SEQ ID NO:114), YIEKDTG (SEQ ID NO:115), FYIEKD (SEQ ID NO:116), FYIEKDT (SEQ ID NO:117), FYIEKDTG (SEQ ID NO:118), LFYIEKD (SEQ ID NO:119), LFYIEKDT (SEQ ID NO:120), LFYIEKDTG (SEQ ID NO:121), ERD, ERDT (SEQ ID NO:122), ERDTG (SEQ ID NO:123), VERD (SEQ ID NO:124), VERDT (SEQ ID NO:125), VERDTG (SEQ ID NO:126), YVERD (SEQ ID NO:127), YVERDT (SEQ ID NO:128), YVERDTG (SEQ ID NO:129), FYVERD (SEQ ID NO:130), FYVERDT (SEQ ID NO:131), FYVERDTG (SEQ ID NO:132), LFYVERD (SEQ ID NO:133), LFYVERDT (SEQ ID NO:134), LFYVERDTG (SEQ ID NO:135), IERD (SEQ ID NO:136), IERDT (SEQ ID NO:137), IERDTG (SEQ ID NO:138), YIERD (SEQ ID NO:139), YIERDT (SEQ ID NO:140), YIERDTG (SEQ ID NO:141), FYIERD (SEQ ID NO:142), FYIERDT (SEQ ID NO:143), FYIERDTG (SEQ ID NO:144), LFYIERD (SEQ ID NO:145), LFYIERDT (SEQ ID NO:146) and LFYIERDTG (SEQ ID NO:147); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmocollin-mediated function is not substantially diminished. For example, the agent may comprise a linear peptide having the sequence N-Ac-LFYIEKDTG-NH<sub>2</sub> (SEQ ID NO:148), N-Ac-LFYVERDTG-NH<sub>2</sub> (SEQ ID NO:149) or N-Ac-LFYIERDTG-NH<sub>2</sub> (SEQ ID NO:150). The desmocollin CAR sequence may, but need not, be present within a cyclic peptide.

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more Desmoglein-1 CAR sequences selected from the group consisting of: RAL, RALN (SEQ ID NO:151), RALNS (SEQ ID NO:152, RALNSM (SEQ ID NO:153), RALNSL (SEQ ID NO:154), RALNSMG (SEQ ID NO:155), RALNSLG (SEQ ID NO:156), CRAL (SEQ ID NO:157), CRALN (SEQ ID NO:158), CRALNS (SEQ ID NO:159), CRALNSM (SEQ ID NO:160), CRALNSL (SEQ ID NO:161), CRALNSMG (SEQ ID NO:162), CRALNSLG (SEQ ID NO:163), YCRAL (SEQ ID NO:164), YCRALN (SEQ ID NO:165), YCRALNS (SEQ ID

NO:166), YCRALNSM (SEQ ID NO:167), YCRALNSL (SEQ ID NO:168), YCRALNSMG (SEQ ID NO:169), YCRALNSLG (SEQ ID NO:170), IYRAL (SEQ ID NO:171), IYRALN (SEQ ID NO:172), IYRALNS (SEQ ID NO:173), IYRALNSM (SEQ ID NO:174), IYRALNSL (SEQ ID NO:175), IYRALNSMG (SEQ ID NO:176) or IYRALNSLG (SEQ ID NO:177); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmoglein-mediated function is not substantially diminished. Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-IYRALNSMG-NH<sub>2</sub> (SEQ ID NO:178) and N-Ac-IYRALNSLG-NH<sub>2</sub> (SEQ ID NO:179).

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more Desmoglein-2 CAR sequences selected from the group consisting of: YAL, YALD (SEQ ID NO:180), YALDA (SEQ ID NO:181), YALDAR (SEQ ID NO:182), YALDARG (SEQ ID NO:183), GYAL (SEQ ID NO:184), GYALD (SEQ ID NO:185), GYALDA (SEQ ID NO:186), GYALDAR (SEQ ID NO:187), GYALDARG (SEQ ID NO:188), TGYAL (SEQ ID NO:189), TGYALD (SEQ ID NO:190), TGYALDA (SEQ ID NO:191), TGYALDAR (SEQ ID NO:192), TGYALDARG (SEQ ID NO:193), LTGYAL (SEQ ID NO:194), LTGYALD (SEQ ID NO:195), LTGYALDA (SEQ ID NO:196), LTGYALDAR (SEQ ID NO:197) or LTGYALDARG (SEQ ID NO:198); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmoglein-mediated function is not substantially diminished. Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac- LTGYALDARG -NH<sub>2</sub> (SEQ ID NO:199).

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more Desmoglein-3 CAR sequences selected from the group consisting of: RAL, RALN (SEQ ID NO:151), RALNA (SEQ ID NO:200), RALNAQ (SEQ ID NO:201), RALNAL (SEQ ID NO:202), RALNAQG (SEQ ID NO:203), RALNALG (SEQ ID NO:204), CRAL (SEQ ID NO:157), CRALN (SEQ ID

NO:158), CRALNA (SEQ ID NO:205), CRALNAQ (SEQ ID NO:206), CRALNAL (SEQ ID NO:207), CRALNAQG (SEQ ID NO:208), CRALNALG (SEQ ID NO:209), TCRAL (SEQ ID NO:210), TCRALN (SEQ ID NO:211), TCRALNA (SEQ ID NO:212), TCRALNAQ (SEQ ID NO:213), TCRALNAL (SEQ ID NO:214),  
 5 TCRALNAQG (SEQ ID NO:215), TCRALNALG (SEQ ID NO:216), ITCRAL (SEQ ID NO:217), ITCRALN (SEQ ID NO:218), ITCRALNA (SEQ ID NO:219), ITCRALNAQ (SEQ ID NO:220), ITCRALNAL (SEQ ID NO:221), ITCRALNAQG (SEQ ID NO:222), ITCRALNALG (SEQ ID NO:223); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or  
 10 insertions such that that ability of the analogue to modulate a desmoglein-mediated function is not substantially diminished. Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-ITCRALNAQG-NH<sub>2</sub> (SEQ ID NO:224) and N-Ac-ITCRALNALG-NH<sub>2</sub> (SEQ ID NO:225).

Within other specific embodiments, a modulating agent as provided  
 15 herein may comprise: (a) one or more Desmocollin-1 CAR sequences selected from the group consisting of: YAT, YATT (SEQ ID NO:226), YATTA (SEQ ID NO:227), YATTAD (SEQ ID NO:228), YATTADG (SEQ ID NO:229), GYAT (SEQ ID NO:230), GYATT (SEQ ID NO:231), GYATTA (SEQ ID NO:232), GYATTAD (SEQ ID NO:233), GYATTADG (SEQ ID NO:234), AYAT (SEQ ID NO:235), AYATT (SEQ ID NO:236), AYATTA (SEQ ID NO:237), AYATTAD (SEQ ID NO:238),  
 20 AYATTADG (SEQ ID NO:239), YGYAT (SEQ ID NO:240), YGYATT (SEQ ID NO:241), YGYATTA (SEQ ID NO:242), YGYATTAD (SEQ ID NO:243), YGYATTADG (SEQ ID NO:244), YAYAT (SEQ ID NO:245), YAYATT (SEQ ID NO:246), YAYATTA (SEQ ID NO:247), YAYATTAD (SEQ ID NO:248),  
 25 YAYATTADG (SEQ ID NO:249), LYGYAT (SEQ ID NO:250), LYGYATT (SEQ ID NO:251), LYGYATTA (SEQ ID NO:252), LYGYATTAD (SEQ ID NO:253), LYGYATTADG (SEQ ID NO:254), LYAYAT (SEQ ID NO:255), LYAYATT (SEQ ID NO:256), LYAYATTA (SEQ ID NO:257), LYAYATTAD (SEQ ID NO:258), LYAYATTADG (SEQ ID NO:259), VYGYAT (SEQ ID NO:260), VYGYATT (SEQ  
 30 ID NO:261), VYGYATTA (SEQ ID NO:262), VYGYATTAD (SEQ ID NO:263),


VYGYATTADG (SEQ ID NO:264), VYAYAT (SEQ ID NO:265), VYAYATT (SEQ ID NO:266), VYAYATTA (SEQ ID NO:267), VYAYATTAD (SEQ ID NO:268), VYAYATTADG (SEQ ID NO:269), IYGYAT (SEQ ID NO:270), IYGYATT (SEQ ID NO:271), IYGYATTA (SEQ ID NO:272), IYGYATTAD (SEQ ID NO:273),  
 5 IYGYATTADG (SEQ ID NO:274), IYAYAT (SEQ ID NO:275), IYAYATT (SEQ ID NO:276), IYAYATTA (SEQ ID NO:277), IYAYATTAD (SEQ ID NO:278) or IYAYATTADG (SEQ ID NO:279); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmocollin-mediated function is  
 10 not substantially diminished. Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-LYGYATTADG-NH<sub>2</sub> (SEQ ID NO:280) N-Ac-LYAYATTADG-NH<sub>2</sub> (SEQ ID NO:281) N-Ac-VYGYATTADG-NH<sub>2</sub> (SEQ ID NO:282) N-Ac- VYAYATTADG-NH<sub>2</sub> (SEQ ID NO:283) N-Ac-IYGYATTADG-NH<sub>2</sub> (SEQ ID NO:284) and N-Ac- IYAYATTADG-NH<sub>2</sub> (SEQ ID  
 15 NO:285).

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more Desmocollin-2 CAR sequences selected from the group consisting of: FAT, FATT (SEQ ID NO:286), FATTP (SEQ ID NO:287), FATTPD (SEQ ID NO:288), FATTPDG (SEQ ID NO:289), AFAT (SEQ ID NO:290),  
 20 AFATT (SEQ ID NO:291), AFATTP (SEQ ID NO:292), AFATTPD (SEQ ID NO:293), AFATTPDG (SEQ ID NO:294), IAFAT (SEQ ID NO:295), IAFATT (SEQ ID NO:296), IAFATTP (SEQ ID NO:297), IAFATTPD (SEQ ID NO:298), IAFATTPDG (SEQ ID NO:299), IIAFAT (SEQ ID NO:300), IIAFATT (SEQ ID NO:301), IIAFATTP (SEQ ID NO:302), IIAFATTPD (SEQ ID NO:303),  
 25 IIAFATTPDG (SEQ ID NO:304), LIAFAT (SEQ ID NO:305), LIAFATT (SEQ ID NO:306), LIAFATTP (SEQ ID NO:307), LIAFATTPD (SEQ ID NO:308), LIAFATTPDG (SEQ ID NO:309); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmocollin-mediated function is not  
 30 substantially diminished. Linear peptides having such sequences may be modified at

the N- and/or C-termini, as in the peptides N-Ac-IIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:310) and N-Ac-LIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:311).

Within other specific embodiments, a modulating agent as provided herein may comprise: (a) one or more desmocollin-3 or desmocollin-4 CAR sequences  
 5 selected from the group consisting of: YAS, YAST (SEQ ID NO:312), YASTA (SEQ ID NO:313), YASTAD (SEQ ID NO:314), YASTADG (SEQ ID NO:315), AYAS (SEQ ID NO:316), AYAST (SEQ ID NO:317), AYASTA (SEQ ID NO:318), AYASTAD (SEQ ID NO:319), AYASTADG (SEQ ID NO:320), IAYAS (SEQ ID NO:321), IAYAST (SEQ ID NO:322), IAYASTA (SEQ ID NO:323), IAYASTAD  
 10 (SEQ ID NO:324), IAYASTADG (SEQ ID NO:325), LIAYAS (SEQ ID NO:326), LIAYAST (SEQ ID NO:327), LIAYASTA (SEQ ID NO:328), LIAYASTAD (SEQ ID NO:329), LIAYASTADG (SEQ ID NO:330); or (b) an analogue of any of the foregoing sequences that differs in one or more substitutions, deletions, additions and/or insertions such that that ability of the analogue to modulate a desmocollin-mediated function is  
 15 not substantially diminished. Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-LIAYASTADG-NH<sub>2</sub> (SEQ ID NO:331).

Within other embodiments, a modulating agent comprises a dsc or dsg CAR sequence that is present within a cyclic peptide. Such cyclic peptides may have  
 20 the formula:



wherein W is a tripeptide selected from the group consisting of NQK, NRN, NKD, EKD, ERD, RAL, YAL, YAT, FAT, and YAS; wherein X<sub>1</sub> and X<sub>2</sub> are optional, and if  
 25 present, are independently selected from the group consisting of amino acid residues and combinations thereof in which the residues are linked by peptide bonds, and wherein X<sub>1</sub> and X<sub>2</sub> independently range in size from 0 to 10 residues, such that the sum of residues contained within X<sub>1</sub> and X<sub>2</sub> ranges from 1 to 12; wherein Y<sub>1</sub> and Y<sub>2</sub> are independently selected from the group consisting of amino acid residues, and wherein a  
 30 covalent bond is formed between residues Y<sub>1</sub> and Y<sub>2</sub>; and wherein Z<sub>1</sub> and Z<sub>2</sub> are optional, and if present, are independently selected from the group consisting of amino



acid residues and combinations thereof in which the residues are linked by peptide bonds.

Within other aspects of the present invention, polynucleotides encoding a modulating agent as described above are provided, along with expression vectors  
5 comprising such a polynucleotide and host cells transformed or transfected with such an expression vector.

The present invention further provides modulating agents that comprise an antibody or antigen-binding fragment thereof that specifically binds to a dsg or dsc CAR sequence as described above and modulates a desmosomal cadherin-mediated  
10 function.

Within further aspects, the present invention provides modulating agents comprising a non-peptide mimetic of any one of the desmosomal cadherin CAR sequences provided above.

In certain embodiments, a modulating agent may be linked to a drug, a  
15 detectable marker, a targeting agent and/or a support material. Modulating agents may also, or alternatively, comprise one or more of: (a) a cell adhesion recognition sequence other than a dsc or dsg CAR sequence; and/or (b) an antibody or antigen-binding fragment thereof that specifically binds to a CAR sequence other than a dsc or dsg CAR sequence. For example, such an adhesion molecule may be a classical cadherin, nonclassical cadherin, integrin, occludin, claudin, N-CAM, fibronectin, laminin or other  
20 extracellular matrix protein.

Within further aspects, pharmaceutical compositions are provided, comprising a modulating agent as described above, in combination with a pharmaceutically acceptable carrier. Such compositions may additionally comprise a  
25 drug and/or one or more of: (a) a peptide comprising a cell adhesion recognition sequence other than a dsc or dsg CAR sequence; and/or (b) an antibody or antigen-binding fragment thereof that specifically binds to a CAR sequence other than a dsc or dsg CAR sequence.

The present invention further provides, within other aspects, methods for  
30 modulating one or more desmosomal cadherin-mediated functions. Such methods

generally comprise contacting a dsg- or dsc-expressing cell with a modulating agent as described above. Suitable cells include, but are not limited to, epithelial cells, endothelial cells and tumor cells. Within such methods, the modulating agent may, but need not, be present within a pharmaceutical composition as recited above.

5           The present invention further provides, within other aspects, methods for modulating cell adhesion, comprising contacting a dsc- and/or dsg-expressing cell with a modulating agent or a pharmaceutical composition as described above. Such modulating agents and compositions may inhibit or enhance cell adhesion.

          Within other aspects, the present invention provides methods for  
10   inhibiting adhesion of dsc- and/or dsg-expressing cells in a mammal, comprising administering to a mammal a modulating agent as described above, wherein the modulating agent inhibits dsc- and/or dsg-mediated cell adhesion.

          Within further aspects, the present invention provides methods for enhancing the delivery of a drug through the skin of a mammal, comprising contacting  
15   epithelial cells of a mammal with a drug and a modulating agent as described above, wherein the step of contacting is performed under conditions and for a time sufficient to allow passage of the drug across the epithelial cells, and wherein the modulating agent inhibits dsg- or dsc-mediated cell adhesion. Such modulating agents may pass into the blood stream of the mammal. Within certain embodiments, the modulating agent is  
20   linked to the drug. The step of contacting may, but need not, be performed via a skin patch comprising the modulating agent and the drug, and such skin patches are further provided herein.

          Methods are further provided for facilitating blood sampling in a mammal, comprising contacting epithelial cells of a mammal with a modulating agent  
25   as described above, wherein the modulating agent inhibits dsg- or dsc-mediated cell adhesion, and wherein the step of contacting is performed under conditions and for a time sufficient to allow passage of one or more blood components across the epithelial cells. The step of contacting may be performed via a skin patch comprising the modulating agent, and (optionally) a reagent for detecting a blood component of

interest, and such kits are specifically provided herein. Within certain embodiments, the epithelial cells are skin cells or are gum cells.

Within further aspects, methods are provided for enhancing the delivery of a drug to a tumor in a mammal, comprising administering to a mammal a modulating agent as described above, wherein the modulating agent inhibits dsg- or dsc-mediated cell adhesion. Suitable tumors include, but are not limited to, bladder tumors, ovarian tumors, breast tumors, stomach tumors and kidney tumors, and the modulating agent may be administered locally to the tumor or may be administered systemically.

Within other aspects, the present invention provides methods for treating cancer in a mammal, comprising administering to a mammal a modulating agent as described above, wherein the modulating agent inhibits dsg- or dsc-mediated cell adhesion. The mammal may be afflicted with a cancer such as a carcinoma, leukemia or melanoma, and the modulating agent may be administered to a tumor or systemically.

The present invention further provides, within other aspects, methods for inducing apoptosis in a desmosomal cadherin-expressing cell, comprising contacting a desmosomal cadherin-expressing cell with a modulating agent as described above, wherein the modulating agent inhibits desmosomal cadherin-mediated cell adhesion.

Within other aspects, the present invention provides methods for enhancing adhesion of desmosomal cadherin-expressing cells, comprising contacting dsg- or dsc-expressing cells with a modulating agent as described above, wherein the modulating agent enhances dsg- or dsc-mediated cell adhesion, wherein the step of contacting is performed under conditions and for a time sufficient to detectably enhance adhesion of the cells. Within certain embodiments, modulating agents for use within such methods are linked to a support molecule or a solid support.

Within related aspects, the present invention provides methods for facilitating wound healing and/or reducing scar tissue in a mammal, comprising contacting a wound in a mammal with a modulating agent as described above, wherein the modulating agent enhances desmosomal cadherin-mediated cell adhesion. Within

certain embodiments, modulating agents for use within such methods are linked to a support molecules or a solid support.

Methods are also provided, within other aspects, for enhancing adhesion of foreign tissue implanted within a mammal, comprising contacting a site of  
5 implantation of foreign tissue in a mammal with a modulating agent as described above, wherein the modulating agent enhances dsg- or dsc-mediated cell adhesion. Such foreign tissue may be a skin graft or organ implant. Within certain embodiments, modulating agents for use within such methods are linked to a support molecules or a solid support.

10 Within further aspects, methods are provided for treating an autoimmune blistering disorder in a mammal, comprising administering to a mammal a modulating agent as described above, wherein the modulating agent enhances desmosomal cadherin-mediated cell adhesion. Within certain embodiments, such an agent may be administered topically to a blister. Modulating agents for use within such methods may  
15 be linked to a support molecule or a solid support.

The present invention further provides methods for detecting the presence of desmosomal cadherin-expressing cells in a sample, comprising: (a) contacting a sample with an antibody or antigen-binding fragment thereof that binds to a dsg or dsc CAR sequence as described above under conditions and for a time  
20 sufficient to allow formation of an antibody-cadherin complex; and (b) detecting the level of antibody-cadherin complex, and therefrom detecting the presence of desmosomal cadherin-expressing cells in a sample. The antibody may be linked to a support material or a detectable marker such as a fluorescent marker. In certain embodiments, the step of detecting is performed using fluorescence activated cell  
25 sorting.

Kits for detecting the presence of dsg- or dsc-expressing cells in a sample are also provided. Such kits may comprise: (a) an antibody or antigen-binding fragment thereof that specifically binds to a desmosomal cadherin CAR sequence; and (b) a detection reagent.

Within other aspects, the present invention provides methods for identifying a compound capable of modulating a desmosomal cadherin-mediated function, comprising: (a) contacting an antibody or antigen-binding fragment thereof that specifically binds to a dsg or dsc CAR sequence as described above with a test  
5 compound; and (b) detecting the level of antibody or fragment that binds to the test compound, and therefrom identifying a compound capable of modulating desmosomal cadherin-mediated cell adhesion.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All  
10 references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1AA are diagrams depicting the structure of classical (Figure 1A) and nonclassical cadherins (Figures 1B to 1AA). The extracellular domains are  
15 designated EC1-EC5 for most cadherins; EC1-EC7 for LI-cadherin and EC1-EC6 for protocadherins and cnr. The hydrophobic domain that transverses the plasma membrane (PM) is represented by TM, and the varying number of cytoplasmic domains are represented by CP. The calcium binding motifs for classical cadherins are shown in Figure 1A by DXNDN (SEQ ID NO:1), DXD, LDRE (SEQ ID NO:2), DXE, and  
20 DVNE (SEQ ID NO:341), and the calcium binding motifs for other cadherins are also indicated above the extracellular domains. Below the extracellular domains, the nine amino acid CAR sequences are shown.

Figure 2 provides the amino acid sequences of representative mammalian nonclassical cadherin extracellular domains, as indicated (SEQ ID NOs:4-  
25 43). Calcium binding motifs are shown in bold, and representative CAR sequences are shown in bold and underlined.

Figure 3 provides the amino acid sequences of representative mammalian desmosomal cadherin EC1 domains, as indicated. Amino acids are represented by their IUPAC codes and – represents a gap. The desmosomal cadherin  
30 CAR sequence is indicated in bold.

Figure 4 depicts the structures of representative cyclic peptide modulating agents.

## 5 DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention provides methods for modulating desmosomal cadherin-mediated functions, such as cell adhesion. The present invention is based, in part, upon the identification of previously unknown cell adhesion recognition (CAR) sequences present in desmosomal cadherins. A modulating agent  
10 may comprise one or more such desmosomal cadherin CAR sequences (or analogues or mimetics thereof), with or without one or more additional cell adhesion molecule CAR sequences, as described below. Peptide CAR sequences may be present within a linear or cyclic peptide. Alternatively, or in addition, a modulating agent may comprise a polynucleotide encoding a polypeptide comprising one or more desmosomal cadherin  
15 CAR sequences and/or a modulating agent may comprise a substance (such as an antibody or antigen-binding fragment thereof) that specifically binds to a desmosomal cadherin CAR sequence.

In general, to modulate a desmosomal cadherin-mediated function, a cell that expresses a desmosomal cadherin is contacted with a modulating agent either *in vivo* or *in vitro*. Within certain aspects, the methods provided herein inhibit a  
20 desmosomal cadherin-mediated function. Such methods include, for example, methods for treating diseases or other conditions characterized by undesirable cell adhesion or for facilitating drug delivery to a specific tissue or tumor. Certain methods may inhibit cell adhesion (*e.g.*, cancer cell adhesion). Alternatively, a modulating agent may, such  
25 as when linked to a matrix or to another modulating agent via a linker, be used to enhance a desmosomal cadherin-mediated function, such as cell adhesion. Such conjugates may be used, for example, to facilitate wound healing or the adhesion of implants.

## MODULATING AGENTS

The term "modulating agent," as used herein, refers to a molecule comprising at least one of the following components:

- (a) a linear or cyclic peptide sequence that is at least 50% identical to  
5 a desmosomal cadherin CAR sequence (*i.e.*, a desmosomal cadherin CAR sequence or an analogue thereof that retains at least 50% sequence identity);
- (b) a mimetic (*e.g.*, peptidomimetic or small molecule mimic) of a desmosomal cadherin CAR sequence;
- (c) a substance, such as an antibody or antigen-binding fragment  
10 thereof, that specifically binds a desmosomal cadherin CAR sequence; and/or
- (d) a polynucleotide encoding a polypeptide that comprises a desmosomal cadherin CAR sequence or analogue thereof.

A modulating agent may consist entirely of one or more of the above elements, or may additionally comprise further peptide and/or non-peptide regions.  
15 Additional peptide regions may be derived from a desmosomal cadherin (preferably an extracellular domain that comprises a CAR sequence) and/or may be heterologous. Within certain preferred embodiments, a modulating agent contains no more than 85 consecutive amino acid residues, and preferably no more than 50 consecutive amino acid residues, present within a desmosomal cadherin.

20 A modulating agent is further capable of modulating a function mediated by a desmosomal cadherin. Such activity may generally be assessed using, for example, representative assays provided herein. Certain modulating agents inhibit an interaction between desmosomal cadherin molecules and/or between a desmosomal cadherin and a different adhesion molecule. For functions (*e.g.*, cell adhesion) that are inhibited by a  
25 full length desmosomal cadherin, such a modulating agent may inhibit the function with an activity that is not substantially diminished relative to the full length desmosomal cadherin (*i.e.*, the modulating agent inhibits the function at least as well as soluble cadherin, when contacted with cells that express the cadherin). For example, a modulating agent may be as effective as soluble desmosomal cadherin in preventing  
30 and/or disrupting adhesion of desmosomal cadherin-expressing cells. Alternatively, to

enhance adhesion of desmosomal cadherin-expressing cells, a modulating agent may comprise an antibody or antigen-binding fragment thereof and/or multiple peptides or mimetics linked to a support material. Such modulating agents may function as a biological glue to bind desmosomal cadherin-expressing cells, and should result in a detectable enhancement of cell adhesion (preferably an enhancement that is at least as great as that observed for immobilized desmosomal cadherin or antibody directed against the cadherin).

The term "nonclassical cadherin," as used herein, refers to a polypeptide that contains characteristic cadherin repeats, but does not contain an HAV CAR sequence. As used herein, a "cadherin repeat" refers to an amino acid sequence that is approximately 110 amino acid residues in length (generally 100 to 120 residues, preferably 105 to 115 residues), comprises an extracellular domain, and contains three calcium binding motifs (DXD, XDXE and DXXDX; SEQ ID NOS:332 and 333 respectively) in the same order and in approximately the same position (*see, e.g.*, Figure 2). The presence of an extracellular domain may generally be determined using well known techniques, such as the presence of one or more of: a hydrophilic sequence, a region that is recognized by an antibody, a region that is cleaved by trypsin and/or a potential glycosylation site with the glycosylation motif Asn-X-Ser/Thr. The second calcium binding motif commonly has the sequence LDRE (SEQ ID NO:2), although variants of this sequence with conservative substitutions are also observed, including MDRE (SEQ ID NO:334), LDFE (SEQ ID NO:335), LDYE (SEQ ID NO:336), IDRE (SEQ ID NO:337), VDRE (SEQ ID NO:338) and IDFE (SEQ ID NO:339). Within most cadherin repeats, the third calcium binding motif has the sequence [L,I,V]-X-[L,I,V]-X-D-X-N-D-[N,H]-X-P (SEQ ID NO:340), wherein residues indicated in brackets may be any one of the recited residues. A preferred third calcium binding motif has the sequence DXNDN (SEQ ID NO:1), although one or both of the D residues may be replaced by an E. Homology among cadherin repeats is generally at least 20%, preferably at least 30%, as determined by the ALIGN algorithm (Myers and Miller, *CABIOS* 4:11-17, 1988). Most cadherins comprise at least five cadherin repeats, along with a hydrophobic domain that transverses the plasma membrane and,



optionally, one or more cytoplasmic domains, as shown in Figures 1B-1AA. Occasionally, however, a cadherin may substitute an extracellular domain that contains fewer than three calcium binding motifs for one or more of the cadherin repeats. For example, as shown in Figure 2, the second extracellular domain of LI-cadherin  
5 comprises only the first calcium binding motif (DXD). As noted above, atypical, or type II, cadherins include cadherin-5 (VE-cadherin), cadherin-6 (K-cadherin), cadherin-7, cadherin-8, cadherin-11 (OB-cadherin), cadherin-12, cadherin-14, cadherin-15 and PB-cadherin. Types III-X include LI-cadherin, T-cadherin, protocadherins (*e.g.*, protocadherins 42, 43 and 68), desmocollins (*e.g.*, desmocollins 1, 2, 3 and 4),  
10 desmogleins (*e.g.*, desmogleins 1 and 2), and cadherin-related neuronal receptors.

The term "desmosomal cadherin" refers to a nonclassical cadherin that is present within the intercellular junction known as the desmosome. Desmosomal cadherins include desmogleins and desmocollins (*see e.g.*, King et al., *Genomics* 18:185-194, 1993; Parker et al., *J. Biol. Chem.* 266:10438-10445, 1991; King et al., *J.*  
15 *Invest. Dermatol.* 105:314-321, 1995; Kawamura et al., *J. Biol. Chem.* 269:26295-26302, 1994; Wheeler et al., *Proc. Natl. Acad. Sci. USA* 88:4796-4800; and Koch et al., *Eur. J. Cell. Biol.* 55:200-208, 1991). Desmogleins and desmocollins are expressed by cells that possess desmosomes, such as epithelial cells, cardiac muscle cells and meningeal cells. These cadherins are involved in intercellular adhesion of such cells,  
20 and may function in a heterotypic manner, whereby a desmocollin isoform and a desmoglein isoform are both required for adhesion. Desmogleins and desmocollins are involved in a number of autoimmune blistering disorders, such as pemphigus vulgaris, pemphigus foliaceus and intercellular IgA dermatosis, and have been shown to have reduced expression in some human carcinomas. The sequences of various extracellular  
25 domains of known desmosomal cadherins are shown in Figures 2 and 3 (SEQ ID NOs:4-43 and SEQ ID NOs:44-57).

A desmosomal cadherin CAR sequence, as used herein, is an amino acid sequence that is present within in a naturally occurring desmosomal cadherin and that is capable of detectably modulating a desmosomal cadherin-mediated function, such as  
30 cell adhesion, as described herein. In other words, contacting a desmosomal cadherin-

expressing cell with a peptide comprising a CAR sequence results in a detectable change in a desmosomal cadherin-mediated function using at least one of the representative assays provided herein. CAR sequences are generally recognized *in vivo* by a desmosomal cadherin or other adhesion molecule (*i.e.*, a molecule that mediates cell adhesion via a receptor on the cell surface), and are necessary for maximal heterophilic and/or homophilic interaction. CAR sequences may be of any length, but generally comprise at least three amino acid residues, preferably 4-16 amino acid residues, and more preferably 5-9 amino acid residues. A peptide modulating agent may comprise any number of amino acid residues, but preferred agents comprise 3-50 residues, preferably 4-16 residues, and more preferably 6-15 residues.

It has been found, within the context of the present invention, that certain nonclassical cadherin CAR sequences share the consensus sequence:

Aaa-Phe-Baa-Ile/Leu/Val-Asp/Asn/Glu-Caa-Daa-Ser/Thr/Asn-Gly  
(SEQ ID NO:3)

Within the consensus sequence, Aaa, Baa, Caa and Daa indicate independently selected amino acid residues; "Ile/Leu/Val" indicates an amino acid that is isoleucine, leucine or valine; "Asp/Asn/Glu" indicates an amino acid that is aspartic acid, asparagine or glutamic acid; and "Ser/Thr/Asn" indicates an amino acid that is serine, threonine or asparagine. Representative desmosomal cadherin CAR sequences, as well as other nonclassical cadherin CAR sequences, are provided within Table I. CAR sequences specifically provided herein further include portions of such representative CAR sequences, as well as longer polypeptides that comprise at least a portion of such sequences. Additional nonclassical cadherin CAR sequences may be identified based on sequence homology to the nonclassical cadherin CAR sequences provided herein, and based on the ability of a peptide comprising such a sequence to modulate a nonclassical cadherin-mediated function within a representative assay described herein. Within certain embodiments, a modulating agent comprises at least three consecutive residues, preferably at least five consecutive residues and more

preferably at least seven consecutive residues, of a desmosomal cadherin CAR sequence that satisfies the above consensus sequence. Similarly, if a CAR sequence of a different nonclassical cadherin is used in combination with a desmosomal cadherin CAR sequence, such CAR sequence(s) should comprise at least three consecutive residues, preferably at least five consecutive residues and more preferably at least seven consecutive residues, of a nonclassical cadherin CAR sequence that satisfies the above consensus sequence.

Table I - Representative Nonclassical Cadherin CAR Sequences

Cadherin	CAR Sequence
Human OB-cadherin EC1	FFVIEEYTG (SEQ ID NO:367)
Human OB-cadherin EC1	IFVIDDKSG (SEQ ID NO:368)
Human OB-cadherin EC2	YFSVEAQTG (SEQ ID NO:369)
Human cadherin-5 EC1	VFRVDAETG (SEQ ID NO:370)
Human cadherin-6 EC1	FFLLEEYTG (SEQ ID NO:371)
Human cadherin-6 EC1	LFIINENTG (SEQ ID NO:372)
Human cadherin-6 EC2	YFSVESETG (SEQ ID NO:373)
Human cadherin-6 EC4	IFNIDSGNG (SEQ ID NO:374)
Chicken cadherin-7 EC1	IFIIDENTG (SEQ ID NO:375)
Chicken cadherin-7 EC2	YFSVEPKTG (SEQ ID NO:376)
Chicken cadherin-7 EC4	YFNIDANSNG (SEQ ID NO:377)
Human cadherin-8 EC1	MFVLEEFSG (SEQ ID NO:378)
Human cadherin-8 EC1	IFQINDVTG (SEQ ID NO:379)
Human cadherin-12 EC1	VFTIDETTG (SEQ ID NO:380)
Human cadherin-12 EC2	YFSIDPKTG (SEQ ID NO:381)
Human cadherin-14 EC1	IFIIDDTTG (SEQ ID NO:382)
Human cadherin-14 EC2	YFSVDPKTG (SEQ ID NO:383)
Human cadherin-14 EC4	FFNIDANTG (SEQ ID NO:384)
Human cadherin-15 EC1	VFSIDKFTG (SEQ ID NO:385)
Human cadherin-15 EC2	LFSIDELTG (SEQ ID NO:386)
Human T-cadherin EC1	IFRINENTG (SEQ ID NO:387)
Rat PB-cadherin EC1	FFVVEEYTG (SEQ ID NO:388)
Rat PB-cadherin EC1	IFLIDELTG (SEQ ID NO:389)
Rat PB-cadherin EC2	HFTVDPKTG (SEQ ID NO:390)
Rat PB-cadherin EC4	IFDIDADTG (SEQ ID NO:391)
Human LI-cadherin EC2	YFQINNKTG (SEQ ID NO:392)
Human protocadherin 43 EC3	LFALDLVTG (SEQ ID NO:393)
Human protocadherin 43 EC5	YFTINRDNG (SEQ ID NO:394)
Human protocadherin 68 EC3	LFSIDPKTG (SEQ ID NO:395)

Human protocadherin 68 EC6	LFEIDPSSG	(SEQ ID NO:396)
Human desmoglein1 EC1	IFVINQKTG	(SEQ ID NO:397)
Human desmoglein1 EC2	MFIINRNTG	(SEQ ID NO:398)
Human desmoglein2 EC2	VFYLNKDTG	(SEQ ID NO:399)
Human desmocollin 1 EC1	LFYIEKDTG	(SEQ ID NO:400)
Human desmocollin 2 EC1	LFYVERDTG	(SEQ ID NO:401)
Human desmocollin 3/4 EC1	LFYIERDTG	(SEQ ID NO:402)
Mouse Cnr1 EC3	KFHIDPVSG	(SEQ ID NO:403)
Mouse Cnr2 EC3	QFSIDADTG	(SEQ ID NO:404)
Mouse Cnr3 EC3	TFHIDSVSG	(SEQ ID NO:405)
Mouse Cnr5 EC3	AFNIDSNSG	(SEQ ID NO:406)
Mouse Cnr6 EC3	KFTIDSSSG	(SEQ ID NO:407)
Mouse Cnr7 EC3	LFTLDEKNG	(SEQ ID NO:408)
Mouse Cnr8 EC3	KFLINEKTG	(SEQ ID NO:409)
CONSENSUS	xFxidxxtG	(SEQ ID NO:3)
	v n s	
	l e n	

Nonclassical cadherin CAR sequences are generally physically located within the cadherin molecule in or near the binding site of an adhesion molecule (*i.e.*, within 10 amino acids, and preferably within 5 amino acids). The location of a binding site may generally be determined using well known techniques, such as evaluating the ability of a portion of the nonclassical cadherin to bind to the same nonclassical cadherin or to another adhesion molecule. Any standard binding assay may be employed for such an evaluation. Recognition of a CAR sequence by the nonclassical cadherin or other adhesion molecule results in a measurable effect on an adhesion molecule function, such as cell adhesion. Peptides comprising a CAR sequence generally inhibit such a function unless linked, as described herein, to form an enhancer of adhesion molecule function.

Certain preferred desmosomal cadherin CAR sequences comprise 3-9 amino acid residues of a desmoglein or desmocollin sequence provided in Table I. For example, a CAR sequence may comprise 3, 4 or 5 residues of a 9 amino acid sequence in Table I. Within certain embodiments, a CAR sequence may include at least residues 5-7 of a sequence in Table I. A desmoglein CAR sequence may comprise, for example, one or more of the sequences: NQK, NQKT (SEQ ID NO:63), NQKTG (SEQ ID NO:64), INQK (SEQ ID NO:65), INQKT (SEQ ID NO:66), INQKTG (SEQ ID

NO:67), VINQK (SEQ ID NO:68), VINQKT (SEQ ID NO:69), VINQKTG (SEQ ID NO:70), FVINQK (SEQ ID NO:71), FVINQKT (SEQ ID NO:72), FVINQKTG (SEQ ID NO:73), IFVINQK (SEQ ID NO:74), IFVINQKT (SEQ ID NO:75), IFVINQKTG (SEQ ID NO:76), NRN, NRNT (SEQ ID NO:77), NRNTG (SEQ ID NO:78), INRN (SEQ ID NO:79), INRNT (SEQ ID NO:80), INRNTG (SEQ ID NO:81), IINRN (SEQ ID NO:82), IINRNT (SEQ ID NO:83), IINRNTG (SEQ ID NO:84), FIINRN (SEQ ID NO:85), FIINRNT (SEQ ID NO:86), FIINRNTG (SEQ ID NO:87), MFIINRN (SEQ ID NO:88), MFIINRNT (SEQ ID NO:89), MFIINRNTG (SEQ ID NO:90), NKD, NKDT (SEQ ID NO:91), NKDTG (SEQ ID NO:92), LNKD (SEQ ID NO:93), LNKDT (SEQ ID NO:94), LNKDTG (SEQ ID NO:95), YLNKD (SEQ ID NO:96), YLNKDT (SEQ ID NO:97), YLNKDTG (SEQ ID NO:98), FYLNKD (SEQ ID NO:99), FYLNKDT (SEQ ID NO:100), FYLNKDTG (SEQ ID NO:101), VFYLNKD (SEQ ID NO:102), VFYLNKDT (SEQ ID NO:103) and VFYLNKDTG (SEQ ID NO:104). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-IFVINQKTG-NH<sub>2</sub> (SEQ ID NO:105), N-Ac-MFIINRNTG-NH<sub>2</sub> (SEQ ID NO:106) and N-Ac-VFYLNKDTG-NH<sub>2</sub> (SEQ ID NO:107).

A desmocollin CAR sequence may comprise, for example, one or more of the sequences EKD, EKDT (SEQ ID NO:108), EKDTG (SEQ ID NO:109), IEKD (SEQ ID NO:110), IEKDT (SEQ ID NO:111), IEKDTG (SEQ ID NO:112), YIEKD (SEQ ID NO:113), YIEKDT (SEQ ID NO:114), YIEKDTG (SEQ ID NO:115), FYIEKD (SEQ ID NO:116), FYIEKDT (SEQ ID NO:117), FYIEKDTG (SEQ ID NO:118), LFYIEKD (SEQ ID NO:119), LFYIEKDT (SEQ ID NO:120), LFYIEKDTG (SEQ ID NO:121), ERD, ERDT (SEQ ID NO:122), ERDTG (SEQ ID NO:123), VERD (SEQ ID NO:124), VERDT (SEQ ID NO:125), VERDTG (SEQ ID NO:126), YVERD (SEQ ID NO:127), YVERDT (SEQ ID NO:128), YVERDTG (SEQ ID NO:129), FYVERD (SEQ ID NO:130), FYVERDT (SEQ ID NO:131), FYVERDTG (SEQ ID NO:132), LFYVERD (SEQ ID NO:133), LFYVERDT (SEQ ID NO:134), LFYVERDTG (SEQ ID NO:135), IERD (SEQ ID NO:136), IERDT (SEQ ID NO:137), IERDTG (SEQ ID NO:138), YIERD (SEQ ID NO:139), YIERDT (SEQ ID NO:140), YIERDTG (SEQ ID NO:141), FYIERD (SEQ ID NO:142), FYIERDT (SEQ ID NO:143), FYIERDTG (SEQ ID NO:144), LFYIERD (SEQ ID NO:145), LFYIERDT (SEQ ID NO:146), LFYIERDTG (SEQ ID NO:147), and combinations thereof.

NO:143), FYIERDTG (SEQ ID NO:144), LFYIERD (SEQ ID NO:145), LFYIERDT (SEQ ID NO:146) and LFYIERDTG (SEQ ID NO:147). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-LFYIEKDTG-NH<sub>2</sub> (SEQ ID NO:148), N-Ac-LFYVERDTG-NH<sub>2</sub> (SEQ ID NO:149) and N-Ac-LFYIERDTG-NH<sub>2</sub> (SEQ ID NO:150).

Further desmosomal cadherin CAR sequences are derived from the EC1 domains recited in Figure 3. A desmoglein-1 CAR sequence may comprise, for example, one or more of the sequences RAL, RALN (SEQ ID NO: 151), RALNS (SEQ ID NO:152), RALNSM (SEQ ID NO:153), RALNSL (SEQ ID NO:154), RALNSMG (SEQ ID NO:155), RALNSLG (SEQ ID NO:156), CRAL (SEQ ID NO:157), CRALN (SEQ ID NO:158), CRALNS (SEQ ID NO:159), CRALNSM (SEQ ID NO:160), CRALNSL (SEQ ID NO:161), CRALNSMG (SEQ ID NO:162), CRALNSLG (SEQ ID NO:163), YCRAL (SEQ ID NO:164), YCRALN (SEQ ID NO:165), YCRALNS (SEQ ID NO:166), YCRALNSM (SEQ ID NO:167), YCRALNSL (SEQ ID NO:168), YCRALNSMG (SEQ ID NO:169), YCRALNSLG (SEQ ID NO:170), IYRAL (SEQ ID NO:171), IYRALN (SEQ ID NO:172), IYRALNS (SEQ ID NO:173), IYRALNSM (SEQ ID NO:174), IYRALNSL (SEQ ID NO:175), IYRALNSMG (SEQ ID NO:176) and IYRALNSLG (SEQ ID NO:177). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-IYRALNSMG-NH<sub>2</sub> (SEQ ID NO:178) and N-Ac-IYRALNSLG-NH<sub>2</sub> (SEQ ID NO:179).

A desmoglein-2 CAR sequence may comprise, for example, one or more of the sequences YAL, YALD (SEQ ID NO:180), YALDA (SEQ ID NO:181), YALDAR (SEQ ID NO:182), YALDARG (SEQ ID NO:183), GYAL (SEQ ID NO:184), GYALD (SEQ ID NO:185), GYALDA (SEQ ID NO:186), GYALDAR (SEQ ID NO:187), GYALDARG (SEQ ID NO:188), TGYAL (SEQ ID NO:189), TGYALD (SEQ ID NO:190), TGYALDA (SEQ ID NO:191), TGYALDAR (SEQ ID NO:192), TGYALDARG (SEQ ID NO:193), LTGYAL (SEQ ID NO:194), LTGYALD (SEQ ID NO:195), LTGYALDA (SEQ ID NO:196), LTGYALDAR (SEQ ID NO:197), LTGYALDARG (SEQ ID NO:198). Linear peptides having such sequences may be

modified at the N- and/or C-termini, as in the peptides N-Ac-LTGYALDARG-NH<sub>2</sub> (SEQ ID NO:199).

A desmoglein-3 CAR sequence may comprise, for example, one or more of the sequences RAL, RALN (SEQ ID NO:151), RALNA (SEQ ID NO:200),  
5 RALNAQ (SEQ ID NO:201), RALNAL (SEQ ID NO:202), RALNAQG (SEQ ID NO:203), RALNALG (SEQ ID NO:204), CRAL (SEQ ID NO:157), CRALN (SEQ ID NO:158), CRALNA (SEQ ID NO:205), CRALNAQ (SEQ ID NO:206), CRALNAL (SEQ ID NO:207), CRALNAQG (SEQ ID NO:208), CRALNALG (SEQ ID NO:209),  
TCRAL (SEQ ID NO:210), TCRALN (SEQ ID NO:211), TCRALNA (SEQ ID NO:212), TCRALNAQ (SEQ ID NO:213), TCRALNAL (SEQ ID NO:214),  
10 TCRALNAQG (SEQ ID NO:215), TCRALNALG (SEQ ID NO:216), ITCRAL (SEQ ID NO:217), ITCRALN (SEQ ID NO:218), ITCRALNA (SEQ ID NO:219), ITCRALNAQ (SEQ ID NO:220), ITCRALNAL (SEQ ID NO:221), ITCRALNAQG (SEQ ID NO:222), ITCRALNALG (SEQ ID NO:223). Linear peptides having such  
15 sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-ITCRALNAQG-NH<sub>2</sub> (SEQ ID NO:224) and N-Ac-ITCRALNALG-NH<sub>2</sub> (SEQ ID NO:225).

A desmocollin-1 CAR sequence may comprise, for example, one or more of the sequences YAT, YATT (SEQ ID NO:226), YATTA (SEQ ID NO:227),  
20 YATTAD (SEQ ID NO:228), YATTADG (SEQ ID NO:229), GYAT (SEQ ID NO:230), GYATT (SEQ ID NO:231), GYATTA (SEQ ID NO:232), GYATTAD (SEQ ID NO:233), GYATTADG (SEQ ID NO:234), AYAT (SEQ ID NO:235), AYATT (SEQ ID NO:236), AYATTA (SEQ ID NO:237), AYATTAD (SEQ ID NO:238), AYATTADG (SEQ ID NO:239), YGYAT (SEQ ID NO:240), YGYATT (SEQ ID NO:241), YGYATTA (SEQ ID NO:242), YGYATTAD (SEQ ID NO:243), YGYATTADG (SEQ ID NO:244), YAYAT (SEQ ID NO:245), YAYATT (SEQ ID NO:246), YAYATTA (SEQ ID NO:247), YAYATTAD (SEQ ID NO:248), YAYATTADG (SEQ ID NO:249), LYGYAT (SEQ ID NO:250), LYGYATT (SEQ ID NO:251), LYGYATTA (SEQ ID NO:252), LYGYATTAD (SEQ ID NO:253),  
30 LYGYATTADG (SEQ ID NO:254), LYAYAT (SEQ ID NO:255), LYAYATT (SEQ

ID NO:256), LYAYATTA (SEQ ID NO:257), LYAYATTAD (SEQ ID NO:258), LYAYATTADG (SEQ ID NO:259), VYGYAT (SEQ ID NO:260), VYGYATT (SEQ ID NO:261), VYGYATTA (SEQ ID NO:262), VYGYATTAD (SEQ ID NO:263), VYGYATTADG (SEQ ID NO:264), VYAYAT (SEQ ID NO:265), VYAYATT (SEQ ID NO:266), VYAYATTA (SEQ ID NO:267), VYAYATTAD (SEQ ID NO:268), VYAYATTADG (SEQ ID NO:269), IYGYAT (SEQ ID NO:270), IYGYATT (SEQ ID NO:271), IYGYATTA (SEQ ID NO:272), IYGYATTAD (SEQ ID NO:273), IYGYATTADG (SEQ ID NO:274), IYAYAT (SEQ ID NO:275), IYAYATT (SEQ ID NO:276), IYAYATTA (SEQ ID NO:277), IYAYATTAD (SEQ ID NO:278) and IYAYATTADG (SEQ ID NO:279). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-LYGYATTADG-NH<sub>2</sub> (SEQ ID NO:280) N-Ac-LYAYATTADG-NH<sub>2</sub> (SEQ ID NO:281) N-Ac-VYGYATTADG-NH<sub>2</sub> (SEQ ID NO:282) N-Ac-VYAYATTADG-NH<sub>2</sub> (SEQ ID NO:283) N-Ac-IYGYATTADG-NH<sub>2</sub> (SEQ ID NO:284) and N-Ac-IYAYATTADG-NH<sub>2</sub> (SEQ ID NO:285).

A desmocollin-2 CAR sequence may comprise, for example, one or more of the sequences FAT, FATT (SEQ ID NO:286), FATTP (SEQ ID NO:287), FATTPD (SEQ ID NO:288), FATTPDG (SEQ ID NO:289), AFAT (SEQ ID NO:290), AFATT (SEQ ID NO:291), AFATTP (SEQ ID NO:292), AFATTPD (SEQ ID NO:293), AFATTPDG (SEQ ID NO:294), IAFAT (SEQ ID NO:295), IAFATT (SEQ ID NO:296), IAFATTP (SEQ ID NO:297), IAFATTPD (SEQ ID NO:298), IAFATTPDG (SEQ ID NO:299), IIAFAT (SEQ ID NO:300), IIAFATT (SEQ ID NO:301), IIAFATTP (SEQ ID NO:302), IIAFATTPD (SEQ ID NO:303), IIAFATTPDG (SEQ ID NO:304), LIAFAT (SEQ ID NO:305), LIAFATT (SEQ ID NO:306), LIAFATTP (SEQ ID NO:307), LIAFATTPD (SEQ ID NO:308), LIAFATTPDG (SEQ ID NO:309). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac- IIAFATTPDG -NH<sub>2</sub> (SEQ ID NO:310) and N-Ac-LIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:311).

A desmocollin-3 or desmocollin-4 CAR sequence may comprise, for example, one or more of the sequences YAS, YAST (SEQ ID NO:312), YASTA (SEQ



ID NO:313), YASTAD (SEQ ID NO:314), YASTADG (SEQ ID NO:315), AYAS (SEQ ID NO:316), AYAST (SEQ ID NO:317), AYASTA (SEQ ID NO:318), AYASTAD (SEQ ID NO:319), AYASTADG (SEQ ID NO:320), IAYAS (SEQ ID NO:321), IAYAST (SEQ ID NO:322), IAYASTA (SEQ ID NO:323), IAYASTAD (SEQ ID NO:324), IAYASTADG (SEQ ID NO:325), LIAYAS (SEQ ID NO:326), LIAYAST (SEQ ID NO:327), LIAYASTA (SEQ ID NO:328), LIAYASTAD (SEQ ID NO:329), LIAYASTADG (SEQ ID NO:330). Linear peptides having such sequences may be modified at the N- and/or C-termini, as in the peptides N-Ac-LIAYASTADG-NH<sub>2</sub> (SEQ ID NO:331).

10           Those of ordinary skill in the art will recognize that similar peptide sequences may be designed to modulate a function mediated by other cadherins, following identification of a CAR sequence as described herein.

          It will be apparent that certain of the peptide sequences provided above may modulate a function mediated by multiple nonclassical cadherins. In general, peptides comprising a greater number of consecutive residues derived from a desmosomal cadherin have a greater specificity for that cadherin. In addition, further flanking sequences may be included to enhance specificity. Such flanking sequences may be identified based on the sequences provided in Figures 2 and 3, or based on published sequences. To achieve specificity (*i.e.*, modulation of a desmosomal cadherin function that is enhanced relative to the modulation of a function mediated by a different cadherin), the addition of 2 to 5 flanking residues (preferably at least one residue on either side of the CAR sequence) is generally sufficient.

          As noted above, modulating agents provided herein may comprise an analogue or mimetic of a desmosomal cadherin CAR sequence. An analogue generally retains at least 50% identity to a native desmosomal cadherin CAR sequence, and modulates a desmosomal cadherin-mediated function as described herein. Such analogues preferably contain at least three consecutive residues of, and more preferably at least five consecutive residues of, a desmosomal cadherin CAR sequence. An analogue may contain any of a variety of amino acid substitutions, additions, deletions and/or modifications (*e.g.*, side chain modifications). Preferred amino acid

substitutions are conservative. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. The critical determining feature of a desmosomal cadherin CAR sequence analogue is the ability to modulate a desmosomal cadherin-mediated function, which may be evaluated using the representative assays provided herein.

A mimetic is a non-peptidyl compound that is conformationally similar to a desmosomal cadherin CAR sequence, such that it modulates a desmosomal cadherin-mediated function as described below. Such mimetics may be designed based on techniques that evaluate the three dimensional structure of the peptide. For example, Nuclear Magnetic Resonance spectroscopy (NMR) and computational techniques may be used to determine the conformation of a desmosomal cadherin CAR sequence. NMR is widely used for structural analyses of both peptidyl and non-peptidyl compounds. Nuclear Overhauser Enhancements (NOE's), coupling constants and chemical shifts depend on the conformation of a compound. NOE data provides the interproton distance between protons through space and can be used to calculate the lowest energy conformation for the desmosomal cadherin CAR sequence. This information can then be used to design mimetics of the preferred conformation. Linear peptides in solution exist in many conformations. By using conformational restriction techniques it is possible to fix the peptide in the active conformation. Conformational restriction can be


achieved by i) introduction of an alkyl group such as a methyl which sterically restricts free bond rotation; ii) introduction of unsaturation which fixes the relative positions of the terminal and geminal substituents; and/or iii) cyclization, which fixes the relative positions of the sidechains. Mimetics may be synthesized where one or more of the amide linkages has been replaced by isosteres, substituents or groups which have the same size or volume such as -CH<sub>2</sub>NH-, -CSNH-, -CH<sub>2</sub>S-, -CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, -CONMe- and others. These backbone amide linkages can also be part of a ring structure (*e.g.*, lactam). Mimetics may be designed where one or more of the side chain functionalities of the desmosomal cadherin CAR sequence are replaced by groups that do not necessarily have the same size or volume, but have similar chemical and/or physical properties which produce similar biological responses. Other mimetics may be small molecule mimics, which may be readily identified from small molecule libraries, based on the three-dimensional structure of the CAR sequence. It should be understood that, within embodiments described below, an analogue or mimetic may be substituted for a desmosomal cadherin CAR sequence.

Modulating agents, or peptide portions thereof, may be linear or cyclic peptides. The term "cyclic peptide," as used herein, refers to a peptide or salt thereof that comprises (1) an intramolecular covalent bond between two non-adjacent residues and (2) at least one desmosomal cadherin CAR sequence or an analogue thereof. The intramolecular bond may be a backbone to backbone, side-chain to backbone or side-chain to side-chain bond (*i.e.*, terminal functional groups of a linear peptide and/or side chain functional groups of a terminal or interior residue may be linked to achieve cyclization). Preferred intramolecular bonds include, but are not limited to, disulfide, amide and thioether bonds. One or more of any of the above desmosomal cadherin CAR sequences, or an analogue or mimetic thereof, may be incorporated into a cyclic peptide, with or without one or more other adhesion molecule binding sites. Additional adhesion molecule binding sites are described in greater detail below.

The size of a cyclic peptide ring generally ranges from 5 to about 15 residues, preferably from 5 to 10 residues. Additional residue(s) may be present on the N-terminal and/or C-terminal side of a desmosomal cadherin CAR sequence, and may

be derived from sequences that flank a desmosomal cadherin CAR sequence, with or without amino acid substitutions and/or other modifications. Alternatively, additional residues present on one or both sides of the CAR sequence(s) may be unrelated to an endogenous sequence (*e.g.*, residues that facilitate cyclization, purification or other manipulation and/or residues having a targeting or other function).

Within certain embodiments, a modulating agent may comprise a cyclic peptide that contains a desmosomal cadherin CAR sequence as provided in Table I (or a portion of such a CAR sequence). Certain cyclic peptides have the formula:



Within this formula, W is a tripeptide selected from the group consisting of NQK, NRN, NKD, EKD, ERD, RAL, YAL, YAT, FAT, and YAS;  $X_1$ , and  $X_2$  are optional, and if present, are independently selected from the group consisting of amino acid residues and combinations thereof in which the residues are linked by peptide bonds, and wherein  $X_1$  and  $X_2$  independently range in size from 0 to 10 residues, such that the sum of residues contained within  $X_1$  and  $X_2$  ranges from 1 to 12;  $Y_1$  and  $Y_2$  are independently selected from the group consisting of amino acid residues, and wherein a covalent bond is formed between residues  $Y_1$  and  $Y_2$ ; and  $Z_1$  and  $Z_2$  are optional, and if present, are independently selected from the group consisting of amino acid residues and combinations thereof in which the residues are linked by peptide bonds.

Cyclic peptides may comprise any of the above CAR sequence(s). Such cyclic peptides may be used as modulating agents without modification, or may be incorporated into a modulating agent.

Representative cyclic peptides comprising a desmoglein CAR sequence include: CNQKC (SEQ ID NO:410), CNQKTC (SEQ ID NO:411), CNQKTGC (SEQ ID NO:412), CINQKC (SEQ ID NO:413), CINQKTC (SEQ ID NO:414), CINQKTGC (SEQ ID NO:415), CVINQKC (SEQ ID NO:416), CVINQKTC (SEQ ID NO:417), CVINQKTGC (SEQ ID NO:418), CFVINQKC (SEQ ID NO:419), CFVINQKTC (SEQ ID NO:420), CFVINQKTGC (SEQ ID NO:421), CIFVINQKC (SEQ ID NO:422),

CIFVINQKTC (SEQ ID NO:423), CIFVINQKTGC (SEQ ID NO:424), CNRNC (SEQ ID NO:425), CNRNTC (SEQ ID NO:426), CNRNTGC (SEQ ID NO:427), CINRNC (SEQ ID NO:428), CINRNTC (SEQ ID NO:429), CINRNTGC (SEQ ID NO:430), CIINRNC (SEQ ID NO:431), CIINRNTC (SEQ ID NO:432), CIINRNTGC (SEQ ID NO:433), CFIINRNC (SEQ ID NO:434), CFIINRNTC (SEQ ID NO:435), CFIINRNTGC (SEQ ID NO:436), CMFIINRNC (SEQ ID NO:437), CMFIINRNTC (SEQ ID NO:438), CMFIINRNTGC (SEQ ID NO:439), CNKDC (SEQ ID NO:440), CNKDTC (SEQ ID NO:441), CNKDTGC (SEQ ID NO:442), CLNKDC (SEQ ID NO:443), CLNKDTC (SEQ ID NO:444), CLNKDTGC (SEQ ID NO:445), CYLNKDC (SEQ ID NO:446), CYLNKDTC (SEQ ID NO:447), CYLNKDTGC (SEQ ID NO:448), CFYLNKDC (SEQ ID NO:449), CFYLNKDTGC (SEQ ID NO:450), CFYLNKDTGC (SEQ ID NO:451), CVFYLNKDC (SEQ ID NO:452), CVFYLNKDTGC (SEQ ID NO:453), CVFYLNKDTGC (SEQ ID NO:454), ENQKK (SEQ ID NO:455), ENQKTK (SEQ ID NO:456), ENQKTGK (SEQ ID NO:457), EINQKK (SEQ ID NO:458), EINQKTK (SEQ ID NO:459), EINQKTGK (SEQ ID NO:460), EVINQKK (SEQ ID NO:461), EVINQKTK (SEQ ID NO:462), EVINQKTGK (SEQ ID NO:463), EFVINQKK (SEQ ID NO:464), EFVINQKTK (SEQ ID NO:465), EFVINQKTGK (SEQ ID NO:466), EIFVINQKK (SEQ ID NO:467), EIFVINQKTK (SEQ ID NO:468), EIFVINQKTGK (SEQ ID NO:469), ENRNC (SEQ ID NO:470), ENRNTK (SEQ ID NO:471), ENRNTGK (SEQ ID NO:472), EINRNC (SEQ ID NO:473), EINRNTK (SEQ ID NO:474), EINRNTGK (SEQ ID NO:475), EIINRNC (SEQ ID NO:476), EIINRNTK (SEQ ID NO:477), EIINRNTGK (SEQ ID NO:478), EFIINRNC (SEQ ID NO:479), EFIINRNTK (SEQ ID NO:480), EFIINRNTGK (SEQ ID NO:481), EMFIINRNC (SEQ ID NO:482), EMFIINRNTK (SEQ ID NO:483), EMFIINRNTGK (SEQ ID NO:484), ENKDK (SEQ ID NO:485), ENKDTK (SEQ ID NO:486), ENKDTGK (SEQ ID NO:487), ELNKDK (SEQ ID NO:488), ELNKDTK (SEQ ID NO:489), ELNKDTGK (SEQ ID NO:490), EYLNKDK (SEQ ID NO:491), EYLNKDTK (SEQ ID NO:492), EYLNKDTGK (SEQ ID NO:493), EFYLNKDK (SEQ ID NO:494), EFYLNKDTK (SEQ ID NO:495), EFYLNKDTGK (SEQ ID NO:496), EVFYLNKDK (SEQ ID NO:497), EVFYLNKDTK (SEQ ID NO:498),

EVFYLNKDTGK (SEQ ID NO:499), KNQKD (SEQ ID NO:500), KNQKTD (SEQ ID NO:501), KNQKTGD (SEQ ID NO:502), KINQKD (SEQ ID NO:503), KINQKTD (SEQ ID NO:504), KINQKTGD (SEQ ID NO:505), KVINQKD (SEQ ID NO:506), KVINQKTD (SEQ ID NO:507), KVINQKTGD (SEQ ID NO:508), KFVINQKD (SEQ ID NO:509), KFVINQKTD (SEQ ID NO:510), KFVINQKTGD (SEQ ID NO:511),  
 5 KIFVINQKD (SEQ ID NO:512), KIFVINQKTD (SEQ ID NO:513), KIFVINQKTGD (SEQ ID NO:514), KNRND (SEQ ID NO:515), KNRNTD (SEQ ID NO:516), KNRNTGD (SEQ ID NO:517), KINRND (SEQ ID NO:518), KINRNTD (SEQ ID NO:519), KINRNTGD (SEQ ID NO:520), KIINRND (SEQ ID NO:521), KIINRNTD (SEQ ID NO:522), KIINRNTGD (SEQ ID NO:523), KFIINRND (SEQ ID NO:524), KFIINRNTD (SEQ ID NO:525), KFIINRNTGD (SEQ ID NO:526), KMFIINRND (SEQ ID NO:527), KMFIINRNTD (SEQ ID NO:528), KMFIINRNTGD (SEQ ID NO:529), KNKDD (SEQ ID NO:530), KNKDTD (SEQ ID NO:531), KNKDTGD (SEQ ID NO:532), KLNKDD (SEQ ID NO:533), KLNKDTD (SEQ ID NO:534),  
 15 KLNKDTGD (SEQ ID NO:535), KYLNKDD (SEQ ID NO:536), KYLNKDTD (SEQ ID NO:537), KYLNKDTGD (SEQ ID NO:538), KFYLNKDD (SEQ ID NO:539), KFYLNKDTD (SEQ ID NO:540), KFYLNKDTGD (SEQ ID NO:541), KVFYLNKDD (SEQ ID NO:542), KVFYLNKDTD (SEQ ID NO:543), KVFYLNKDTGD (SEQ ID NO:544), DNQKK (SEQ ID NO:545), DNQKTK (SEQ ID NO:546), DNQKTGK (SEQ ID NO:547), DINQKK (SEQ ID NO:548), DINQKTK (SEQ ID NO:549), DINQKTGK (SEQ ID NO:550), DVINQKK (SEQ ID NO:551), DVINQKTK (SEQ ID NO:552), DVINQKTGK (SEQ ID NO:553), DFVINQKK (SEQ ID NO:554), DFVINQKTK (SEQ ID NO:555), DFVINQKTGK (SEQ ID NO:556), DIFVINQKK (SEQ ID NO:557), DIFVINQKTK (SEQ ID NO:558), DIFVINQKTGK (SEQ ID NO:559), DNRNK (SEQ ID NO:560), DNRNTK (SEQ ID NO:561), DNRNTGK (SEQ ID NO:562), DINRNK (SEQ ID NO:563), DINRNTK (SEQ ID NO:564), DINRNTGK (SEQ ID NO:565), DIINRNK (SEQ ID NO:566), DIINRNTK (SEQ ID NO:567), DIINRNTGK (SEQ ID NO:568), DFIINRNK (SEQ ID NO:569), DFIINRNTK (SEQ ID NO:570), DFIINRNTGK (SEQ ID NO:571), DMFIINRNK (SEQ ID NO:572),  
 25 DMFIINRNTK (SEQ ID NO:573), DMFIINRNTGK (SEQ ID NO:574), DNKDK

(SEQ ID NO:575), DNKDTK (SEQ ID NO:576), DNKDTGK (SEQ ID NO:577),  
DLNKDK (SEQ ID NO:578), DLNKDTK (SEQ ID NO:579), DLNKDTGK (SEQ ID  
NO:580), DYLNKDK (SEQ ID NO:581), DYLNKDTK (SEQ ID NO:582),  
DYLNKDTGK (SEQ ID NO:583), DFYLNKDK (SEQ ID NO:584), DFYLNKDTK  
5 (SEQ ID NO:585), DFYLNKDTGK (SEQ ID NO:586), DVFYLNKDK (SEQ ID  
NO:587), DVFYLNKDTK (SEQ ID NO:588), DVFYLNKDTGK (SEQ ID NO:589),  
KKNOKE (SEQ ID NO:590), KNQKTE (SEQ ID NO:591), KNQKTGE (SEQ ID  
NO:592), KINOKE (SEQ ID NO:593), KINQKTE (SEQ ID NO:594), KINQKTGE  
(SEQ ID NO:595), KVINQKE (SEQ ID NO:596), KVINQKTE (SEQ ID NO:597),  
10 KVINQKTGE (SEQ ID NO:598), KFVINQKE (SEQ ID NO:599), KFVINQKTE (SEQ  
ID NO:600), KFVINQKTGE (SEQ ID NO:601), KIFVINQKE (SEQ ID NO:602),  
KIFVINQKTE (SEQ ID NO:603), KIFVINQKTGE (SEQ ID NO:604), KNRNE (SEQ  
ID NO:605), KNRNTE (SEQ ID NO:606), KNRNTGE (SEQ ID NO:607), KINRNE  
(SEQ ID NO:608), KINRNTE (SEQ ID NO:609), KINRNTGE (SEQ ID NO:610),  
15 KIINRNE (SEQ ID NO:611), KIINRNTE (SEQ ID NO:612), KIINRNTGE (SEQ ID  
NO:613), KFIINRNE (SEQ ID NO:614), KFIINRNTE (SEQ ID NO:615),  
KFIINRNTGE (SEQ ID NO:616), KMFIINRNE (SEQ ID NO:617), KMFIINRNTE  
(SEQ ID NO:618), KMFIINRNTGE (SEQ ID NO:619), KNKDE (SEQ ID NO:620),  
KNKDTE (SEQ ID NO:621), KNKDTGE (SEQ ID NO:622), KLNKDE (SEQ ID  
20 NO:623), KLNKDTE (SEQ ID NO:624), KLNKDTGE (SEQ ID NO:625), KYLNKDE  
(SEQ ID NO:626), KYLNKDTE (SEQ ID NO:627), KYLNKDTGE (SEQ ID NO:628),  
KFYLNKDE (SEQ ID NO:629), KFYLNKDTE (SEQ ID NO:630), KFYLNKDTGE  
(SEQ ID NO:631), KVFYLNKDE (SEQ ID NO:632), KVFYLNKDTE (SEQ ID  
NO:633), KVFYLNKDTGE (SEQ ID NO:634), NQKTG (SEQ ID NO:635), INQKT  
25 (SEQ ID NO:636), INQKTG (SEQ ID NO:637), VINQK (SEQ ID NO:638), VINQKT  
(SEQ ID NO:639), VINQKTG (SEQ ID NO:640), FVINQK (SEQ ID NO:641),  
FVINQKT (SEQ ID NO:642), FVINQKTG (SEQ ID NO:643), IFVINQK (SEQ ID  
NO:644), IFVINQKT (SEQ ID NO:645), IFVINQKTG (SEQ ID NO:646), NRNTG  
(SEQ ID NO:647), INRNT (SEQ ID NO:648), INRNTG (SEQ ID NO:649), IINRN  
30 (SEQ ID NO:650), IINRNT (SEQ ID NO:651), IINRNTG (SEQ ID NO:652), FIINRN

(SEQ ID NO:653), FIINRNT (SEQ ID NO:654), FIINRNTG (SEQ ID NO:655),  
MFIINRN (SEQ ID NO:656), MFIINRNT (SEQ ID NO:657), MFIINRNTG (SEQ ID  
NO:658), NKDTG (SEQ ID NO:659), LNKDT (SEQ ID NO:660), LNKDTG (SEQ ID  
NO:661), YLNKD (SEQ ID NO:662), YLNKDT (SEQ ID NO:663), YLNKDTG (SEQ  
5 ID NO:664), FYLNKD (SEQ ID NO:665), FYLNKDT (SEQ ID NO:666),  
FYLNKDTG (SEQ ID NO:667), VFYLNKD (SEQ ID NO:668), VFYLNKDT (SEQ  
ID NO:669) and VFYLNKDTG (SEQ ID NO:670).

Representative cyclic peptides comprising a desmocollin CAR sequence  
include: CEKDC (SEQ ID NO:671), CEKDTG (SEQ ID NO:672), CEKDTGC (SEQ  
10 ID NO:673), CIEKDC (SEQ ID NO:674), CIEKDTG (SEQ ID NO:675), CIEKDTGC  
(SEQ ID NO:676), CYIEKDC (SEQ ID NO:677), CYIEKDTG (SEQ ID NO:678),  
CYIEKDTGC (SEQ ID NO:679), CFYIEKDC (SEQ ID NO:680), CFYIEKDTG (SEQ  
ID NO:681), CFYIEKDTGC (SEQ ID NO:682), CLFYIEKDC (SEQ ID NO:683),  
CLFYIEKDTG (SEQ ID NO:684), CLFYIEKDTGC (SEQ ID NO:685), CERDC (SEQ  
15 ID NO:686), CERDTG (SEQ ID NO:687), CERDTGC (SEQ ID NO:688), CVERDC  
(SEQ ID NO:689), CVERDTG (SEQ ID NO:690), CVERDTGC (SEQ ID NO:691),  
CYVERDC (SEQ ID NO:692), CYVERDTG (SEQ ID NO:693), CYVERDTGC (SEQ  
ID NO:694), CFYVERDC (SEQ ID NO:695), CFYVERDTG (SEQ ID NO:696),  
CFYVERDTGC (SEQ ID NO:697), CLFYVERDC (SEQ ID NO:698), CLFYVERDTG  
20 (SEQ ID NO:699), CLFYVERDTGC (SEQ ID NO:700), CIERDC (SEQ ID NO:701),  
CIERDTG (SEQ ID NO:702), CIERDTGC (SEQ ID NO:703), CYIERDC (SEQ ID  
NO:704), CYIERDTG (SEQ ID NO:705), CYIERDTGC (SEQ ID NO:706),  
CFYIERDC (SEQ ID NO:707), CFYIERDTG (SEQ ID NO:708), CFYIERDTGC  
(SEQ ID NO:709), CLFYIERDC (SEQ ID NO:710), CLFYIERDTG (SEQ ID  
25 NO:711), CLFYIERDTGC (SEQ ID NO:712), EEKDK (SEQ ID NO:713), EEKDTK  
(SEQ ID NO:714), EEKDTGK (SEQ ID NO:715), EIEKDK (SEQ ID NO:716),  
EIEKDTK (SEQ ID NO:717), EIEKDTGK (SEQ ID NO:718), EYIEKDK (SEQ ID  
NO:719), EYIEKDTK (SEQ ID NO:720), EYIEKDTGK (SEQ ID NO:721),  
EFYIEKDK (SEQ ID NO:722), EFYIEKDTK (SEQ ID NO:723), EFYIEKDTGK  
30 (SEQ ID NO:724), ELFYIEKDK (SEQ ID NO:725), ELFYIEKDTK (SEQ ID



NO:726), ELFYIEKDTGK (SEQ ID NO:727), EERDK (SEQ ID NO:728), EERDTK  
 (SEQ ID NO:729), EERDTGK (SEQ ID NO:730), EVERDK (SEQ ID NO:731),  
EVERDTK (SEQ ID NO:732), EVERDTGK (SEQ ID NO:733), EYVERDK (SEQ ID  
 NO:734), YVERDTK (SEQ ID NO:735), EYVERDTGK (SEQ ID NO:736),  
 5 EFYVERDK (SEQ ID NO:737), EFYVERDTK (SEQ ID NO:738), EFYVERDTGK  
 (SEQ ID NO:739), ELFYVERDK (SEQ ID NO:740), ELFYVERDTK (SEQ ID  
 NO:741), ELFYVERDTGK (SEQ ID NO:742), EIERDK (SEQ ID NO:743),  
EIERDTK (SEQ ID NO:744), EIERDTGK (SEQ ID NO:745), EYIERDK (SEQ ID  
 NO:746), EYIERDTK (SEQ ID NO:747), EYIERDTGK (SEQ ID NO:748),  
 10 EFYIERDK (SEQ ID NO:749), EFYIERDTK (SEQ ID NO:750), EFYIERDTGK  
 (SEQ ID NO:751), ELFYIERDK (SEQ ID NO:752), ELFYIERDTK (SEQ ID  
 NO:753), ELFYIERDTGK (SEQ ID NO:754), KEKDD (SEQ ID NO:755), KEKDTD  
 (SEQ ID NO:756), KEKDTGD (SEQ ID NO:757), KIEKDD (SEQ ID NO:758),  
KIEKDTD (SEQ ID NO:759), KIEKDTGD (SEQ ID NO:760), KYIEKDD (SEQ ID  
 15 NO:761), KYIEKDTD (SEQ ID NO:762), KYIEKDTGD (SEQ ID NO:763),  
KFYIEKDD (SEQ ID NO:764), KFYIEKDTD (SEQ ID NO:765), KFYIEKDTGD  
 (SEQ ID NO:766), KLFYIEKDD (SEQ ID NO:767), KLFYIEKDTD (SEQ ID  
 NO:768), KLFYIEKDTGD (SEQ ID NO:769), KERDD (SEQ ID NO:770), KERDTD  
 (SEQ ID NO:771), KERDTGD (SEQ ID NO:772), KVERDD (SEQ ID NO:773),  
 20 KVERDTD (SEQ ID NO:774), KVERDTGD (SEQ ID NO:775), KYVERDD (SEQ ID  
 NO:776), KYVERDTD (SEQ ID NO:777), KYVERDTGD (SEQ ID NO:778),  
KFYVERDD (SEQ ID NO:779), KFYVERDTD (SEQ ID NO:780), KFYVERDTGD  
 (SEQ ID NO:781), KLFYVERDD (SEQ ID NO:782), KLFYVERDTD (SEQ ID  
 NO:783), KLFYVERDTGD (SEQ ID NO:784), KIERDD (SEQ ID NO:785),  
 25 KIERDTD (SEQ ID NO:786), KIERDTGD (SEQ ID NO:787), KYIERD (SEQ ID  
 NO:788), KYIERDTD (SEQ ID NO:789), KYIERDTGD (SEQ ID NO:790),  
KFYIERDD (SEQ ID NO:791), KFYIERDTD (SEQ ID NO:792), KFYIERDTGD  
 (SEQ ID NO:793), KLFYIERDD (SEQ ID NO:794), KLFYIERDTD (SEQ ID  
 NO:795), KLFYIERDTGD (SEQ ID NO:796), DEKDK (SEQ ID NO:797), DEKDTK  
 30 (SEQ ID NO:798), DEKDTGK (SEQ ID NO:799), DIEKDK (SEQ ID NO:800),

DIEKDTK (SEQ ID NO:801), DIEKDTGK (SEQ ID NO:802), DYIEKDK (SEQ ID NO:803), DYIEKDTK (SEQ ID NO:804), DYIEKDTGK (SEQ ID NO:805), DFYIEKDK (SEQ ID NO:806), DFYIEKDTK (SEQ ID NO:807), DFYIEKDTGK (SEQ ID NO:808), DLFYIEKDK (SEQ ID NO:809), DLFYIEKDTK (SEQ ID NO:810), DLFYIEKDTGK (SEQ ID NO:811), DERDK (SEQ ID NO:812), DERDTK (SEQ ID NO:813), DERDTGK (SEQ ID NO:814), DVERDK (SEQ ID NO:815), DVERDTK (SEQ ID NO:816), DVERDTGK (SEQ ID NO:817), DYVERDK (SEQ ID NO:818), DYVERDTK (SEQ ID NO:819), DYVERDTGK (SEQ ID NO:820), DFYVERDK (SEQ ID NO:821), DFYVERDTK (SEQ ID NO:822), DFYVERDTGK (SEQ ID NO:823), DLFYVERDK (SEQ ID NO:824), DLFYVERDTK (SEQ ID NO:825), DLFYVERDTGK (SEQ ID NO:826), DIERDK (SEQ ID NO:827), DIERDTK (SEQ ID NO:828), DIERDTGK (SEQ ID NO:829), DYIERDK (SEQ ID NO:830), DYIERDTK (SEQ ID NO:831), DYIERDTGK (SEQ ID NO:832), DFYIERDK (SEQ ID NO:833), DFYIERDTK (SEQ ID NO:834), DFYIERDTGK (SEQ ID NO:835), DLFYIERDK (SEQ ID NO:836), DLFYIERDTK (SEQ ID NO:837), DLFYIERDTGK (SEQ ID NO:838), KEKDE (SEQ ID NO:839), KEKDTK (SEQ ID NO:840), KEKDTGE (SEQ ID NO:841), KIEKDE (SEQ ID NO:842), KIEKDTK (SEQ ID NO:843), KIEKDTGE (SEQ ID NO:844), KYIEKDE (SEQ ID NO:845), KYIEKDTK (SEQ ID NO:846), KYIEKDTGE (SEQ ID NO:847), KFYIEKDE (SEQ ID NO:848), KFYIEKDTK (SEQ ID NO:849), KFYIEKDTGE (SEQ ID NO:850), KLFYIEKDE (SEQ ID NO:851), KLFYIEKDTK (SEQ ID NO:852), KLFYIEKDTGE (SEQ ID NO:853), KERDE (SEQ ID NO:854), KERDTK (SEQ ID NO:855), KERDTGE (SEQ ID NO:856), KVERDE (SEQ ID NO:857), KVERDTK (SEQ ID NO:858), KVERDTGE (SEQ ID NO:859), KYVERDE (SEQ ID NO:860), KYVERDTK (SEQ ID NO:861), KYVERDTGE (SEQ ID NO:862), KFYVERDE (SEQ ID NO:863), KFYVERDTK (SEQ ID NO:864), KFYVERDTGE (SEQ ID NO:865), KLFYVERDE (SEQ ID NO:866), KLFYVERDTK (SEQ ID NO:867), KLFYVERDTGE (SEQ ID NO:868), KIERDE (SEQ ID NO:869), KIERDTK (SEQ ID NO:870), KIERDTGE (SEQ ID NO:871), KYIERDE (SEQ ID NO:872), KYIERDTK (SEQ ID NO:873), KYIERDTGE (SEQ ID NO:874),

KFYIERDE (SEQ ID NO:875), KFYIERDTE (SEQ ID NO:876), KFYIERDTGE  
 (SEQ ID NO:877), KLFYIERDE (SEQ ID NO:878), KLFYIERDTE (SEQ ID  
 NO:879), KLFYIERDTGE (SEQ ID NO:880), EKDTG (SEQ ID NO:881), IEKDT  
 (SEQ ID NO:882), IEKDTG (SEQ ID NO:883), YIEKD (SEQ ID NO:884), YIEKDT  
 5 (SEQ ID NO:885), YIEKDTG (SEQ ID NO:886), FYIEKD (SEQ ID NO:887),  
FYIEKDT (SEQ ID NO:888), FYIEKDTG (SEQ ID NO:889), LFYIEKD (SEQ ID  
 NO:890), LFYIEKDT (SEQ ID NO:891), LFYIEKDTG (SEQ ID NO:892), ERDTG  
 (SEQ ID NO:893), VERDT (SEQ ID NO:894), VERDTG (SEQ ID NO:895), YVERD  
 (SEQ ID NO:896), YVERDT (SEQ ID NO:897), YVERDTG (SEQ ID NO:898),  
 10 FYVERD (SEQ ID NO:899), FYVERDT (SEQ ID NO:900), FYVERDTG (SEQ ID  
 NO:901), LFYVERD (SEQ ID NO:902), LFYVERDT (SEQ ID NO:903),  
LFYVERDTG (SEQ ID NO:904), IERDT (SEQ ID NO:905), IERDTG (SEQ ID  
 NO:906), YIERD (SEQ ID NO:907), YIERDT (SEQ ID NO:908), YIERDTG (SEQ ID  
 NO:909), FYIERD (SEQ ID NO:910), FYIERDT (SEQ ID NO:911), FYIERDTG  
 15 (SEQ ID NO:912), LFYIERD (SEQ ID NO:913), LFYIERDT (SEQ ID NO:914) and  
LFYIERDTG (SEQ ID NO:915).

Representative cyclic peptides comprising a desmoglein-1 CAR  
 sequence include: CRALC (SEQ ID NO:916), CRALNC (SEQ ID NO:917),  
CRALNSC (SEQ ID NO:918), CRALNSMC (SEQ ID NO:919), CRALNSLC (SEQ ID  
 NO:920), CRALNSMGC (SEQ ID NO:921), CRALNSLGC (SEQ ID NO:922),  
 20 CCRALC (SEQ ID NO:923), CCRALNC (SEQ ID NO:924), CCRALNSC (SEQ ID  
 NO:925), CCRALNSMC (SEQ ID NO:926), CCRALNSLC (SEQ ID NO:927),  
CCRALNSMGC (SEQ ID NO:928), CCRALNSLGC (SEQ ID NO:929), CYCRALC  
 (SEQ ID NO:930), CYCRALNC (SEQ ID NO:931), CYCRALNSC (SEQ ID NO:932),  
 25 CYCRALNSMC (SEQ ID NO:933), CYCRALNSLC (SEQ ID NO:934),  
CYCRALNSMGC (SEQ ID NO:935), CYCRALNSLGC (SEQ ID NO:936),  
CIYRALC (SEQ ID NO:937), CIYRALNC (SEQ ID NO:938), CIYRALNSC (SEQ ID  
 NO:939), CIYRALNSMC (SEQ ID NO:940), CIYRALNSLC (SEQ ID NO:941),  
CIYRALNSMGC (SEQ ID NO:942), CIYRALNSLGC (SEQ ID NO:943), ERALK  
 30 (SEQ ID NO:944), ERALNK (SEQ ID NO:945), ERALNSK (SEQ ID NO:946),

ERALNSMK (SEQ ID NO:947), ERALNSLK (SEQ ID NO:948), ERALNSMGK  
 (SEQ ID NO:949), ERALNSLGK (SEQ ID NO:950), ECRALK (SEQ ID NO:951),  
ECRALNK (SEQ ID NO:952), ECRALNSK (SEQ ID NO:953), ECRALNSMK (SEQ  
 ID NO:954), ECRALNSLK (SEQ ID NO:955), ECRALNSMGK (SEQ ID NO:956),  
 5 ECRALNSLGK (SEQ ID NO:957), EYCRALK (SEQ ID NO:958), EYCRALNK  
 (SEQ ID NO:959), EYCRALNSK (SEQ ID NO:960), EYCRALNSMK (SEQ ID  
 NO:961), EYCRALNSLK (SEQ ID NO:962), EYCRALNSMGK (SEQ ID NO:963),  
EYCRALNSLGK (SEQ ID NO:964), EYRALNK (SEQ ID NO:965), EYRALNSK  
 (SEQ ID NO:966), EYRALNSMK (SEQ ID  
 10 NO:968), EYRALNSLK (SEQ ID NO:969), EYRALNSMGK (SEQ ID NO:970),  
EYRALNSLGK (SEQ ID NO:971), KRALE (SEQ ID NO:972), KRALNE (SEQ ID  
 NO:973), KRALNSE (SEQ ID NO:974), KRALNSME (SEQ ID NO:975),  
KRALNSLE (SEQ ID NO:976), KRALNSMGE (SEQ ID NO:977), KRALNSLGE  
 (SEQ ID NO:978), KCRALE (SEQ ID NO:979), KCRALNE (SEQ ID NO:980),  
 15 KCRALNSE (SEQ ID NO:981), KCRALNSME (SEQ ID NO:982), KCRALNSLE  
 (SEQ ID NO:983), KCRALNSMGE (SEQ ID NO:984), KCRALNSLGE (SEQ ID  
 NO:985), KYCRALE (SEQ ID NO:986), KYCRALNE (SEQ ID NO:987),  
KYCRALNSE (SEQ ID NO:988), KYCRALNSME (SEQ ID NO:989),  
KYCRALNSLE (SEQ ID NO:990), KYCRALNSMGE (SEQ ID NO:991),  
 20 KYCRALNSLGE (SEQ ID NO:992), KIYRALE (SEQ ID NO:993), KIYRALNE  
 (SEQ ID NO:994), KIYRALNSE (SEQ ID NO:995), KIYRALNSME (SEQ ID  
 NO:996), KIYRALNSLE (SEQ ID NO:997), KIYRALNSMGE (SEQ ID NO:998),  
KIYRALNSLGE (SEQ ID NO:999), DRALK (SEQ ID NO:1000), DRALNK (SEQ ID  
 NO:1001), DRALNSK (SEQ ID NO:1002), DRALNSMK (SEQ ID NO:1003),  
 25 DRALNSLK (SEQ ID NO:1004), DRALNSMGK (SEQ ID NO:1005), DRALNSLGK  
 (SEQ ID NO:1006), DCRALK (SEQ ID NO:1007), DCRALNK (SEQ ID NO:1008),  
DCRALNSK (SEQ ID NO:1009), DCRALNSMK (SEQ ID NO:1010), DCRALNSLK  
 (SEQ ID NO:1011), DCRALNSMGK (SEQ ID NO:1012), DCRALNSLGK (SEQ ID  
 NO:1013), DYCRALK (SEQ ID NO:1014), DYCRALNK (SEQ ID NO:1015),  
 30 DYCRALNSK (SEQ ID NO:1016), DYCRALNSMK (SEQ ID NO:1017),

DYCRALNSLK (SEQ ID NO:1018), DYCRALNSMGK (SEQ ID NO:1019),  
DYCRALNSLGK (SEQ ID NO:1020), DIYRALK (SEQ ID NO:1021), DIYRALNK  
 (SEQ ID NO:1022), DIYRALNSK (SEQ ID NO:1023), DIYRALNSMK (SEQ ID  
 NO:1024), DIYRALNSLK (SEQ ID NO:1025), DIYRALNSMGK (SEQ ID NO:1026),  
 5 DIYRALNSLGK (SEQ ID NO:1027), KRALD (SEQ ID NO:1028), KRALND (SEQ  
 ID NO:1029), KRALNSD (SEQ ID NO:1030), KRALNSMD (SEQ ID NO:1031),  
KRALNSLD (SEQ ID NO:1032), KRALNSMGD (SEQ ID NO:1033), KRALNSLGD  
 (SEQ ID NO:1034), KCRALD (SEQ ID NO:1035), KCRALND (SEQ ID NO:1036),  
KCRALNSD (SEQ ID NO:1037), KCRALNSMD (SEQ ID NO:1038), KCRALNSLD  
 10 (SEQ ID NO:1039), KCRALNSMGD (SEQ ID NO:1040), KCRALNSLGD (SEQ ID  
 NO:1041), KYCRALD (SEQ ID NO:1042), KYCRALND (SEQ ID NO:1043),  
KYCRALNSD (SEQ ID NO:1044), KYCRALNSMD (SEQ ID NO:1045),  
KYCRALNSLD (SEQ ID NO:1046), KYCRALNSMGD (SEQ ID NO:1047),  
KYCRALNSLGD (SEQ ID NO:1048), KIYRALD (SEQ ID NO:1049), KIYRALND  
 15 (SEQ ID NO:1050), KIYRALNSD (SEQ ID NO:1051), KIYRALNSMD (SEQ ID  
 NO:1052), KIYRALNSLD (SEQ ID NO:1053), KIYRALNSMGD (SEQ ID NO:1054),  
KIYRALNSLGD (SEQ ID NO:1055), RALNS (SEQ ID NO:1056), RALNSM (SEQ  
 ID NO:1057), RALNSL (SEQ ID NO:1058), RALNSMG (SEQ ID NO:1059),  
RALNSLG (SEQ ID NO:1060), CRALN (SEQ ID NO:1061), CRALNS (SEQ ID  
 20 NO:1062), CRALNSM (SEQ ID NO:1063), CRALNSL (SEQ ID NO:1064),  
CRALNSMG (SEQ ID NO:1065), CRALNSLG (SEQ ID NO:1066), YCRAL (SEQ ID  
 NO:1067), YCRALN (SEQ ID NO:1068), YCRALNS (SEQ ID NO:1069),  
YCRALNSM (SEQ ID NO:1070), YCRALNSL (SEQ ID NO:1071), YCRALNSMG  
 (SEQ ID NO:1072), YCRALNSLG (SEQ ID NO:1073), IYRAL (SEQ ID NO:1074),  
 25 IYRALN (SEQ ID NO:1075), IYRALNS (SEQ ID NO:1076), IYRALNSM (SEQ ID  
 NO:1077), IYRALNSL (SEQ ID NO:1078), IYRALNSMG (SEQ ID NO:1079) and  
IYRALNSLG (SEQ ID NO:1080).

Representative cyclic peptides comprising a desmoglein-2 CAR  
 sequence include: CYALC (SEQ ID NO:1081), CYALDC (SEQ ID NO:1082),  
 30 CYALDAC (SEQ ID NO:1083), CYALDARC (SEQ ID NO:1084), CYALDARGC

(SEQ ID NO:1085), CGYALC (SEQ ID NO:1086), CGYALDC (SEQ ID NO:1087),  
CGYALDAC (SEQ ID NO:1088), CGYALDARC (SEQ ID NO:1089),  
CGYALDARGC (SEQ ID NO:1090), CTGYALC (SEQ ID NO:1091), CTGYALDC  
5 (SEQ ID NO:1092), CTGYALDAC (SEQ ID NO:1093), CTGYALDARC (SEQ ID  
NO:1094), CTGYALDARGC (SEQ ID NO:1095), CLTGYALC (SEQ ID NO:1096),  
CLTGYALDC (SEQ ID NO:1097), CLTGYALDAC (SEQ ID NO:1098),  
CLTGYALDARC (SEQ ID NO:1099), CLTGYALDARGC (SEQ ID NO:1100),  
EYALK (SEQ ID NO:1101), EYALDK (SEQ ID NO:1102), EYALDAK (SEQ ID  
NO:1103), EYALDARK (SEQ ID NO:1104), EYALDARGK (SEQ ID NO:1105),  
10 EGYALK (SEQ ID NO:1106), EGYALDK (SEQ ID NO:1107), EGYALDAK (SEQ  
ID NO:1108), EGYALDARK (SEQ ID NO:1109), EGYALDARGK (SEQ ID  
NO:1110), ETGYALK (SEQ ID NO:1111), ETGYALDK (SEQ ID NO:1112),  
ETGYALDAK (SEQ ID NO:1113), ETGYALDARK (SEQ ID NO:1114),  
ETGYALDARGK (SEQ ID NO:1115), ELTGYALK (SEQ ID NO:1116),  
15 ELTGYALDK (SEQ ID NO:1117), ELTGYALDAK (SEQ ID NO:1118),  
ELTGYALDARK (SEQ ID NO:1119), ELTGYALDARGK (SEQ ID NO:1120),  
KYALE (SEQ ID NO:1121), KYALDE (SEQ ID NO:1122), KYALDAE (SEQ ID  
NO:1123), KYALDARE (SEQ ID NO:1124), KYALDARGE (SEQ ID NO:1125),  
KGYALE (SEQ ID NO:1126), KGYALDE (SEQ ID NO:1127), KGYALDAE (SEQ  
20 ID NO:1128), KGYALDARE (SEQ ID NO:1129), KGYALDARGE (SEQ ID  
NO:1130), KTGYALE (SEQ ID NO:1131), KTGYALDE (SEQ ID NO:1132),  
KTGYALDAE (SEQ ID NO:1133), KTGYALDARE (SEQ ID NO:1134),  
KTGYALDARGE (SEQ ID NO:1135), KLTGYALE (SEQ ID NO:1136),  
KLTGYALDE (SEQ ID NO:1137), KLTGYALDAE (SEQ ID NO:1138),  
25 KLTGYALDARE (SEQ ID NO:1139), KLTGYALDARGE (SEQ ID NO:1140),  
DYALK (SEQ ID NO:1141), DYALDK (SEQ ID NO:1142), DYALDAK (SEQ ID  
NO:1143), DYALDARK (SEQ ID NO:1144), DYALDARGK (SEQ ID NO:1145),  
DGYALK (SEQ ID NO:1146), DGYALDK (SEQ ID NO:1147), DGYALDAK (SEQ  
ID NO:1148), DGYALDARK (SEQ ID NO:1149), DGYALDARGK (SEQ ID  
30 NO:1150), DTGYALK (SEQ ID NO:1151), DTGYALDK (SEQ ID NO:1152),

DTGYALDAK (SEQ ID NO:1153), DTGYALDARK (SEQ ID NO:1154),  
DTGYALDARGK (SEQ ID NO:1155), DLTGYALK (SEQ ID NO:1156),  
DLTGYALDK (SEQ ID NO:1157), DLTGYALDAK (SEQ ID NO:1158),  
DLTGYALDARK (SEQ ID NO:1159), DLTGYALDARGK (SEQ ID NO:1160),  
5 KYALD (SEQ ID NO:1161), KYALDD (SEQ ID NO:1162), KYALDAD (SEQ ID  
NO:1163), KYALDARD (SEQ ID NO:1164), KYALDARGD (SEQ ID NO:1165),  
KGYALD (SEQ ID NO:1166), KGYALDD (SEQ ID NO:1167), KGYALDAD (SEQ  
ID NO:1168), KGYALDARD (SEQ ID NO:1169), KGYALDARGD (SEQ ID  
NO:1170), KTGYALD (SEQ ID NO:1171), KTGYALDD (SEQ ID NO:1172),  
10 KTGYALDAD (SEQ ID NO:1173), KTGYALDARD (SEQ ID NO:1174),  
KTGYALDARGD (SEQ ID NO:1175), KLTGYALD (SEQ ID NO:1176),  
KLTGYALDD (SEQ ID NO:1177), KLTGYALDAD (SEQ ID NO:1178),  
KLTGYALDARD (SEQ ID NO:1179), KLTGYALDARGD (SEQ ID NO:1180),  
YALDA (SEQ ID NO:1181), YALDAR (SEQ ID NO:1182), YALDARG (SEQ ID  
15 NO:1183), GYALD (SEQ ID NO:1184), GYALDA (SEQ ID NO:1185), GYALDAR  
(SEQ ID NO:1186), GYALDARG (SEQ ID NO:1187), TGYAL (SEQ ID NO:1188),  
TGYALD (SEQ ID NO:1189), TGYALDA (SEQ ID NO:1190), TGYALDAR (SEQ ID  
NO:1191), TGYALDARG (SEQ ID NO:1192), LTGYAL (SEQ ID NO:1193),  
LTGYALD (SEQ ID NO:1194), LTGYALDA (SEQ ID NO:1195), LTGYALDAR  
20 (SEQ ID NO:1196) and LTGYALDARG (SEQ ID NO:1197).

Representative cyclic peptides comprising a desmoglein-3 CAR  
sequence include: CRALC (SEQ ID NO:916), CRALNC (SEQ ID NO:917),  
CRALNAC (SEQ ID NO:1198), CRALNAQC (SEQ ID NO:1199), CRALNALC (SEQ  
ID NO:1200), CRALNAQGC (SEQ ID NO:1201), CRALNALGC (SEQ ID NO:1202),  
25 CCRALC (SEQ ID NO:923), CCRALNC (SEQ ID NO:924), CCRALNAC (SEQ ID  
NO:1203), CCRALNAQC (SEQ ID NO:1204), CCRALNALC (SEQ ID NO:1205),  
CCRALNAQGC (SEQ ID NO:1206), CCRALNALGC (SEQ ID NO:1207),  
CTCRALC (SEQ ID NO:1208), CTCRALNC (SEQ ID NO:1209), CTCRALNAC  
(SEQ ID NO:1210), CTCRALNAQC (SEQ ID NO:1211), CTCRALNALC (SEQ ID  
30 NO:1212), CTCRALNAQGC (SEQ ID NO:1213), CTCRALNALGC (SEQ ID

NO:1214), CITCRALC (SEQ ID NO:1215), CITCRALNC (SEQ ID NO:1216),  
CITCRALNAC (SEQ ID NO:1217), CITCRALNAQC (SEQ ID NO:1218),  
CITCRALNALC (SEQ ID NO:1219), CITCRALNAQGC (SEQ ID NO:1220),  
CITCRALNALGC (SEQ ID NO:1221), ERALK (SEQ ID NO:944), ERALNK (SEQ  
 5 ID NO:945), ERALNAK (SEQ ID NO:1222), ERALNAQK (SEQ ID NO:1223),  
ERALNALK (SEQ ID NO:1224), ERALNAQK (SEQ ID NO:1225), ERALNALGK  
 (SEQ ID NO:1226), ECRALK (SEQ ID NO:951), ECRALNK (SEQ ID NO:952),  
ECRALNAK (SEQ ID NO:1227), ECRALNAQK (SEQ ID NO:1228), ECRALNALK  
 (SEQ ID NO:1229), ECRALNAQK (SEQ ID NO:1230), ECRALNALGK (SEQ ID  
 10 NO:1231), ETCRALK (SEQ ID NO:1232), ETCRALNK (SEQ ID NO:1233),  
ETCRALNAK (SEQ ID NO:1234), ETCRALNAQK (SEQ ID NO:1235),  
ETCRALNALK (SEQ ID NO:1236), ETCRALNAQK (SEQ ID NO:1237),  
ETCRALNALGK (SEQ ID NO:1238), EITCRALK (SEQ ID NO:1239), EITCRALNK  
 (SEQ ID NO:1240), EITCRALNAK (SEQ ID NO:1241), EITCRALNAQK (SEQ ID  
 15 NO:1242), EITCRALNALK (SEQ ID NO:1243), EITCRALNAQK (SEQ ID  
 NO:1244), EITCRALNALGK (SEQ ID NO:1245), KRALE (SEQ ID NO:972);  
KRALNE (SEQ ID NO:973), KRALNAE (SEQ ID NO:1246), KRALNAQE (SEQ ID  
 NO:1247), KRALNALE (SEQ ID NO:1248), KRALNAQGE (SEQ ID NO:1249),  
KRALNALGE (SEQ ID NO:1250), KCRALE (SEQ ID NO:979), KCRALNE (SEQ ID  
 20 NO:980), KCRALNAE (SEQ ID NO:1251), KCRALNAQE (SEQ ID NO:1252),  
KCRALNALE (SEQ ID NO:1253), KCRALNAQGE (SEQ ID NO:1254),  
KCRALNALGE (SEQ ID NO:1255), KTCRALE (SEQ ID NO:1256), KTCRALNE  
 (SEQ ID NO:1257), KTCRALNAE (SEQ ID NO:1258), KTCRALNAQE (SEQ ID  
 NO:1259), KTCRALNALE (SEQ ID NO:1260), KTCRALNAQGE (SEQ ID  
 25 NO:1261), KTCRALNALGE (SEQ ID NO:1262), KITCRALE (SEQ ID NO:1263),  
KITCRALNE (SEQ ID NO:1264), KITCRALNAE (SEQ ID NO:1265),  
KITCRALNAQE (SEQ ID NO:1266), KITCRALNALE (SEQ ID NO:1267),  
KITCRALNAQGE (SEQ ID NO:1268), KITCRALNALGE (SEQ ID NO:1269),  
DRALK (SEQ ID NO:1000), DRALNK (SEQ ID NO:1001), DRALNAK (SEQ ID  
 30 NO:1270), DRALNAQK (SEQ ID NO:1271), DRALNALK (SEQ ID NO:1272),



DRALNAQ GK (SEQ ID NO:1273), DRALNAL GK (SEQ ID NO:1274), DCRALK  
 (SEQ ID NO:1007), DCRALNK (SEQ ID NO:1008), DCRALNAK (SEQ ID  
 NO:1275), DCRALNAQK (SEQ ID NO:1276), DCRALNALK (SEQ ID NO:1277),  
DCRALNAQ GK (SEQ ID NO:1278), DCRALNAL GK (SEQ ID NO:1279),  
 5 DTCRALK (SEQ ID NO:1280), DTCRALNK (SEQ ID NO:1281), DTCRALNAK  
 (SEQ ID NO:1282), DTCRALNAQK (SEQ ID NO:1283), DTCRALNALK (SEQ ID  
 NO:1284), DTCRALNAQ GK (SEQ ID NO:1285), DTCRALNAL GK (SEQ ID  
 NO:1286), DITCRALK (SEQ ID NO:1287), DITCRALNK (SEQ ID NO:1288),  
DITCRALNAK (SEQ ID NO:1289), DITCRALNAQK (SEQ ID NO:1290),  
 10 DITCRALNALK (SEQ ID NO:1291), DITCRALNAQ GK (SEQ ID NO:1292),  
DITCRALNAL GK (SEQ ID NO:1293), KRALD (SEQ ID NO:1028), KRALND (SEQ  
 ID NO:1029), KRALNAD (SEQ ID NO:1294), KRALNAQD (SEQ ID NO:1295),  
KRALNALD (SEQ ID NO:1296), KRALNAQGD (SEQ ID NO:1297), KRALNALGD  
 (SEQ ID NO:1298), KCRALD (SEQ ID NO:1034), KCRALND (SEQ ID NO:1035),  
 15 KCRALNAD (SEQ ID NO:1299), KCRALNAQD (SEQ ID NO:1300), KCRALNALD  
 (SEQ ID NO:1301), KCRALNAQGD (SEQ ID NO:1302), KCRALNALGD (SEQ ID  
 NO:1303), KTCRALD (SEQ ID NO:1304), KTCRALND (SEQ ID NO:1305),  
KTCRALNAD (SEQ ID NO:1306), KTCRALNAQD (SEQ ID NO:1307),  
KTCRALNALD (SEQ ID NO:1308), KTCRALNAQGD (SEQ ID NO:1309),  
 20 KTCRALNALGD (SEQ ID NO:1310), KITCRALD (SEQ ID NO:1311),  
KITCRALND (SEQ ID NO:1312), KITCRALNAD (SEQ ID NO:1313),  
KITCRALNAQD (SEQ ID NO:1314), KITCRALNALD (SEQ ID NO:1315),  
KITCRALNAQGD (SEQ ID NO:1316), KITCRALNALGD (SEQ ID NO:1317),  
RALNA (SEQ ID NO:1318), RALNAQ (SEQ ID NO:1319), RALNAL (SEQ ID  
 25 NO:1320), RALNAQG (SEQ ID NO:1321), RALNALG (SEQ ID NO:1322), CRALN  
 (SEQ ID NO:1323), CRALNA (SEQ ID NO:1324), CRALNAQ (SEQ ID NO:1325),  
CRALNAL (SEQ ID NO:1326), CRALNAQG (SEQ ID NO:1327), CRALNALG (SEQ  
 ID NO:1328), TCRAL (SEQ ID NO:1329), TCRALN (SEQ ID NO:1330), TCRALNA  
 (SEQ ID NO:1331), TCRALNAQ (SEQ ID NO:1332), TCRALNAL (SEQ ID  
 30 NO:1333), TCRALNAQG (SEQ ID NO:1334), TCRALNALG (SEQ ID NO:1335),

ITCRAL (SEQ ID NO:1336), ITCRALN (SEQ ID NO:1337), ITCRALNA (SEQ ID NO:1338), ITCRALNAQ (SEQ ID NO:1339), ITCRALNAL (SEQ ID NO:1340), ITCRALNAQG (SEQ ID NO:1341) and ITCRALNALG (SEQ ID NO:1342).

Representative cyclic peptides comprising a desmocollin-1 CAR  
 5 sequence include: CYATC, CYATTC, CYATTAC, CYATTADC, CYATTADGC,  
CGYATC, CGYATTC, CGYATTAC, CGYATTADC, CGYATTADGC, CAYATC,  
CAYATTC, CAYATTAC, CAYATTADC, CAYATTADGC, CYGYATC,  
CYGYATTC, CYGYATTAC, CYGYATTADC, CYGYATTADGC, CYAYATC,  
CYAYATTC, CYAYATTAC, CYAYATTADC, CYAYATTADGC, CLYGYATC,  
 10 CLYGYATTC, CLYGYATTAC, CLYGYATTADC, CLYGYATTADGC,  
CLYAYATC, CLYAYATTC, CLYAYATTAC, CLYAYATTADC,  
CLYAYATTADGC, CVYGYATC, CVYGYATTC, CVYGYATTAC,  
CVYGYATTADC, CVYGYATTADGC, CVYAYATC, CVYAYATTC,  
CVYAYATTAC, CVYAYATTADC, CVYAYATTADGC, CIYGYATC,  
 15 CIYGYATTC, CIYGYATTAC, CIYGYATTADC, CIYGYATTADGC, CIYAYATC,  
CIYAYATTC, CIYAYATTAC, CIYAYATTADC, CIYAYATTADGC, EYATK,  
EYATTK, EYATTAK, EYATTADK, EYATTADGK, EGYATK, EGYATTK,  
EGYATTAK, EGYATTADK, EGYATTADGK, EAYATK, EAYATTK,  
EAYATTAK, EAYATTADK, EAYATTADGK, EYGYATK, EYGYATTK,  
 20 EYGYATTAK, EYGYATTADK, EYGYATTADGK, EYAYATK, EYAYATTK,  
EYAYATTAK, EYAYATTADK, EYAYATTADGK, ELYGYATK, ELYGYATTK,  
ELYGYATTAK, ELYGYATTADK, ELYGYATTADGK, ELYAYATK,  
ELYAYATTK, ELYAYATTAK, ELYAYATTADK, ELYAYATTADGK,  
EVYGYATK, EVYGYATTK, EVYGYATTAK, EVYGYATTADK,  
 25 EVYGYATTADGK, EVYAYATK, EVYAYATTK, EVYAYATTAK,  
EVYAYATTADK, EVYAYATTADGK, EIYGYATK, EIYGYATTK,  
EIYGYATTAK, EIYGYATTADK, EIYGYATTADGK, EIYAYATK, EIYAYATTK,  
EIYAYATTAK, EIYAYATTADK, EIYAYATTADGK, KYATE, KYATTE,  
KYATTAE, KYATTADK, KYATTADGK, KGYATE, KGYATTE, KGYATTAE,  
 30 KGYATTADK, KGYATTADGK, KAYATE, KAYATTE, KAYATTAE,

KAYATTADE, KAYATTADGE, KYGYATE, KYGYATTE, KYGYATTAE,  
KYGYATTADE, KYGYATTADGE, KYAYATE, KYAYATTE, KYAYATTAE,  
KYAYATTADE, KYAYATTADGE, KLYGYATE, KLYGYATTE, KLYGYATTAE,  
KLYGYATTADE, KLYGYATTADGE, KLYAYATE, KLYAYATTE,  
5 KLYAYATTAE, KLYAYATTADE, KLYAYATTADGE, KVYGYATE,  
KVYGYATTE, KVYGYATTAE, KVYGYATTADE, KVYGYATTADGE,  
KVYAYATE, KVYAYATTE, KVYAYATTAE, KVYAYATTADE,  
KVYAYATTADGE, KIYGYATE, KIYGYATTE, KIYGYATTAE, KIYGYATTADE,  
KIYGYATTADGE, KIYAYATE, KIYAYATTE, KIYAYATTAE, KIYAYATTADE,  
10 KIYAYATTADGE, DYATK, DYATTK, DYATTAK, DYATTADK, DYATTADGK,  
DGYATK, DGYATTK, DGYATTAK, DGYATTADK, DGYATTADGK, DAYATK,  
DAYATTK, DAYATTAK, DAYATTADK, DAYATTADGK, DYGYATK,  
DYGYATTK, DYGYATTAK, DYGYATTADK, DYGYATTADGK, DYAYATK,  
DYAYATTK, DYAYATTAK, DYAYATTADK, DYAYATTADGK, DLYGYATK,  
15 DLYGYATTK, DLYGYATTAK, DLYGYATTADK, DLYGYATTADGK,  
DLYAYATK, DLYAYATTK, DLYAYATTAK, DLYAYATTADK,  
DLYAYATTADGK, DVYGYATK, DVYGYATTK, DVYGYATTAK,  
DVYGYATTADK, DVYGYATTADGK, DVYAYATK, DVYAYATTK,  
DVYAYATTAK, DVYAYATTADK, DVYAYATTADGK, DIYGYATK,  
20 DIYGYATTK, DIYGYATTAK, DIYGYATTADK, DIYGYATTADGK, DIYAYATK,  
DIYAYATTK, DIYAYATTAK, DIYAYATTADK, DIYAYATTADGK, KYATD,  
KYATTD, KYATTAD, KYATTADD, KYATTADGD, KGYATD, KGYATTD,  
KGYATTAD, KGYATTADD, KGYATTADGD, KAYATD, KAYATTD,  
KAYATTAD, KAYATTADD, KAYATTADGD, KYGYATD, KYGYATTD,  
25 KYGYATTAD, KYGYATTADD, KYGYATTADGD, KYAYATD, KYAYATTD,  
KYAYATTAD, KYAYATTADD, KYAYATTADGD, KLYGYATD, KLYGYATTD,  
KLYGYATTAD, KLYGYATTADD, KLYGYATTADGD, KLYAYATD,  
KLYAYATTD, KLYAYATTAD, KLYAYATTADD, KLYAYATTADGD,  
KVYGYATD, KVYGYATTD, KVYGYATTAD, KVYGYATTADD,  
30 KVYGYATTADGD, KVYAYATD, KVYAYATTD, KVYAYATTAD,

KVYAYATTADD, KVYAYATTADGD, KIYGYATD, KIYGYATTD,  
KIYGYATTAD, KIYGYATTADD, KIYGYATTADGD, KIYAYATD, KIYAYATTD,  
KIYAYATTAD, KIYAYATTADD, KIYAYATTADGD, YATTA, YATTAD,  
YATTADG, GYATT, GYATTA, GYATTAD, GYATTADG, AYATT, AYATTA,  
5 AYATTAD, AYATTADG, YGYAT, YGYATT, YGYATTA, YGYATTAD,  
YGYATTADG, YAYAT, YAYATT, YAYATTA, YAYATTAD, YAYATTADG,  
LYGYAT, LYGYATT, LYGYATTA, LYGYATTAD, LYGYATTADG, LYAYAT,  
LYAYATT, LYAYATTA, LYAYATTAD, LYAYATTADG, VYGYAT, VYGYATT,  
VYGYATTA, VYGYATTAD, VYGYATTADG, VYAYAT, VYAYATT,  
10 VYAYATTA, VYAYATTAD, VYAYATTADG, IYGYAT, IYGYATT, IYGYATTA,  
IYGYATTAD, IYGYATTADG, IYAYAT, IYAYATT, IYAYATTA, IYAYATTAD  
and IYAYATTADG.

Representative cyclic peptides comprising a desmocollin-2 CAR  
sequence include: CFATC, CFATTC, CFATTPC, CFATTPDC, CFATTPDGC,  
15 CAFATC, CAFATTC, CAFATTPC, CAFATTPDC, CAFATTPDGC, CIAFATC,  
CIAFATTC, CIAFATTPC, CIAFATTPDC, CIAFATTPDGC, CIIAFATC,  
CIIAFATTC, CIIAFATTPC, CIIAFATTPDC, CIIAFATTPDGC, CLIAFATC,  
CLIAFATTC, CLIAFATTPC, CLIAFATTPDC, CLIAFATTPDGC, EFATK,  
EFATTK, EFATTPK, EFATTPDK, EFATTPDGK, EAFATK, EAFATTK,  
20 EAFATTPK, EAFATTPDK, EAFATTPDGK, EIAFATK, EIAFATTK, EIAFATTPK,  
EIAFATTPDK, EIAFATTPDGK, EIIAFATK, EIIAFATTK, EIIAFATTPK,  
EIIAFATTPDK, EIIAFATTPDGK, ELIAFATK, ELIAFATTK, ELIAFATTPK,  
ELIAFATTPDK, ELIAFATTPDGK, KFATE, KFATTE, KFATTPE, KFATTPDE,  
KFATTPDGE, KAFATE, KAFATTE, KAFATTPE, KAFATTPDE, KAFATTPDGE,  
25 KIAFATE, KIAFATTE, KIAFATTPE, KIAFATTPDE, KIAFATTPDGE, KIIAFATE,  
KIIAFATTE, KIIAFATTPE, KIIAFATTPDE, KIIAFATTPDGE, KLIAFATE,  
KLIAFATTE, KLIAFATTPE, KLIAFATTPDE, KLIAFATTPDGE, DFATK,  
DFATTK, DFATTPK, DFATTPDK, DFATTPDGK, DAFATK, DAFATTK,  
DAFATTPK, DAFATTPDK, DAFATTPDGK, DIAFATK, DIAFATTK,  
30 DIAFATTPK, DIAFATTPDK, DIAFATTPDGK, DIIAFATK, DIIAFATTK,

DIIAFATTPK, DIIAFATTPDK, DIIAFATTPDGK, DLIAFATK, DLIAFATTK,  
DLIAFATTPK, DLIAFATTPDK, DLIAFATTPDGK, KFATD, KFATTD, KFATTPD,  
KFATTPDD, KFATTPDGD, KAFATD, KAFATTD, KAFATTPD, KAFATTPDD,  
KAFATTPDGD, KIAFATD, KIAFATTD, KIAFATTPD, KIAFATTPDD,  
5 KIAFATTPDGD, KIIAFATD, KIIAFATTD, KIIAFATTPD, KIIAFATTPDD,  
KIIAFATTPDGD, KLIAFATD, KLIAFATTD, KLIAFATTPD, KLIAFATTPDD,  
KLIAFATTPDGD, FATTP, FATTPD, FATTPDG, AFATT, AFATTP, AFATTPD,  
AFATTPDG, IAFAT, IAFATT, IAFATTP, IAFATTPD, IAFATTPDG, IIAFAT,  
IIAFATT, IIAFATTP, IIAFATTPD, IIAFATTPDG, LIAFAT, LIAFATT, LIAFATTP,  
10 LIAFATTPD, LIAFATTPDG.

Representative cyclic peptides comprising a desmocollin-3 or  
desmocollin-4 CAR sequence include: CYASC, CYASTC, CYASTAC, CYASTADC,  
CYASTADGC, CAYASC, CAYASTC, CAYASTAC, CAYASTADC,  
CAYASTADGC, CIAYASC, CIAYASTC, CIAYASTAC, CIAYASTADC,  
15 CIAYASTADGC, CLIAYASC, CLIAYASTC, CLIAYASTAC, CLIAYASTADC,  
CLIAYASTADGC, EYASK, EYASTK, EYASTAK, EYASTADK, EYASTADGK,  
EAYASK, EAYASTK, EAYASTAK, EAYASTADK, EAYASTADGK, EIAYASK,  
EIAYASTK, EIAYASTAK, EIAYASTADK, EIAYASTADGK, ELIAYASK,  
ELIAYASTK, ELIAYASTAK, ELIAYASTADK, ELIAYASTADGK, KYASE,  
20 KYASTE, KYASTAE, KYASTADE, KYASTADGE, KAYASE, KAYASTE,  
KAYASTAE, KAYASTADE, KAYASTADGE, KIAYASE, KIAYASTE,  
KIAYASTAE, KIAYASTADE, KIAYASTADGE, KLIAYASE, KLIAYASTE,  
KLIAYASTAE, KLIAYASTADE, KLIAYASTADGE, DYASK, DYASTK,  
DYASTAK, DYASTADK, DYASTADGK, DAYASK, DAYASTK, DAYASTAK,  
25 DAYASTADK, DAYASTADGK, DIAYASK, DIAYASTK, DIAYASTAK,  
DIAYASTADK, DIAYASTADGK, DLIAYASK, DLIAYASTK, DLIAYASTAK,  
DLIAYASTADK, DLIAYASTADGK, KYASD, KYASTD, KYASTAD,  
KYASTADD, KYASTADGD, KAYASD, KAYASTD, KAYASTAD, KAYASTADD,  
KAYASTADGD, KIAYASD, KIAYASTD, KIAYASTAD, KIAYASTADD,  
30 KIAYASTADGD, KLIAYASD, KLIAYASTD, KLIAYASTAD, KLIAYASTADD.

KLIAYASTADGD, YASTA, YASTAD, YASTADG, AYAST, AYASTA,  
AYASTAD, AYASTADG, IAYAS, IAYAST, IAYASTA, IAYASTAD,  
IAYASTADG, LIAYAS, LIAYAST, LIAYASTA, LIAYASTAD and LIAYASTADG.

As noted above, certain preferred modulating agents comprise a peptide  
5 (containing a desmosomal cadherin CAR sequence or an analogue thereof) in which at  
least one terminal amino acid residue is modified (*e.g.*, the N-terminal amino group is  
modified by, for example, acetylation or alkoxybenzylation and/or an amide or ester is  
formed at the C-terminus). It has been found, within the context of the present  
invention, that the addition of at least one such group to a linear or cyclic peptide  
10 modulating agent may improve the ability of the agent to modulate a desmosomal  
cadherin-mediated function. Certain preferred modulating agents contain modifications  
at the N- and C-terminal residues, such as N-terminal acetylation and/or C-terminal  
amidation.

The present invention further contemplates desmosomal cadherin CAR  
15 sequences from other organisms. Such CAR sequences may be identified based upon  
similarity to the sequences provided herein, and the ability to modulate a desmosomal  
cadherin-mediated function such as may be confirmed as described herein.

Within certain embodiments, as discussed below, cyclic peptides that  
contain small CAR sequences (*e.g.*, three residues without significant flanking  
20 sequences) are preferred for modulating desmosomal cadherin-mediated functions.  
Such peptides may contain an N-acetyl group and a C-amide group. Small cyclic  
peptides may generally be used to specifically modulate adhesion of cancer and/or other  
cell types by topical administration or by systemic administration, with or without  
linking a targeting agent to the peptide, as discussed below.

25 Within embodiments in which inhibition of a desmosomal cadherin-  
interaction is desired, a modulating agent may contain one desmosomal cadherin CAR  
sequence, or multiple CAR sequences that are adjacent to one another (*i.e.*, without  
intervening sequences) or in close proximity (*i.e.*, separated by peptide and/or non-  
peptide linkers to give a distance between the desmosomal CAR sequences that ranges  
30 from about 0.1 to 400 nm). A linker may be any molecule (including peptide and/or

non-peptide sequences) that does not contain a CAR sequence and that can be covalently linked to at least two peptide sequences. Using a linker, CAR sequence-containing peptides and other peptide or protein sequences may be joined end-to-end (*i.e.*, the linker may be covalently attached to the carboxyl or amino group of each peptide sequence), and/or via side chains. One linker that can be used for such purposes is  $(H_2N(CH_2)_nCO_2H)$ , or derivatives thereof, where  $n$  ranges from 1 to 4. Other linkers that may be used will be apparent to those of ordinary skill in the art. Peptide and non-peptide linkers may generally be incorporated into a modulating agent using any appropriate method known in the art.

Within embodiments in which enhancement of cell adhesion mediated by a desmosomal cadherin is desired, a modulating agent may contain multiple desmosomal cadherin CAR sequences, or antibodies that specifically bind to such sequences, joined by linkers as described above. For enhancers of cadherin function, the linker distance should generally be 400-10,000 nm. One linker that can be used for such purposes is  $(H_2N(CH_2)_nCO_2H)_m$ , or derivatives thereof, where  $n$  ranges from 1 to 10 and  $m$  ranges from 1 to 4000. For example, if glycine ( $H_2NCH_2CO_2H$ ) or a multimer thereof is used as a linker, each glycine unit corresponds to a linking distance of 2.45 angstroms, or 0.245 nm, as determined by calculation of its lowest energy conformation when linked to other amino acids using molecular modeling techniques. Similarly, aminopropanoic acid corresponds to a linking distance of 3.73 angstroms, aminobutanoic acid to 4.96 angstroms, aminopentanoic acid to 6.30 angstroms and amino hexanoic acid to 6.12 angstroms. Enhancement of cell adhesion may also be achieved by attachment of multiple modulating agents to a support material, as discussed further below.

Certain preferred modulating agents may comprise a desmoglein CAR sequence in combination with a desmocollin CAR sequence. When the CAR sequences are joined by linkers as described above, such agents may be used, for example, to promote cell adhesion in a variety of contexts. By way of example, such an agent could comprise the dsg-2 CAR sequence YAL in combination with the dsc-2 CAR sequence FAT.

A modulating agent as described herein may additionally comprise one or more CAR sequences for one or more different adhesion molecules (including, but not limited to, nonclassical cadherins and other CAMs) and/or one or more substances, such as antibodies or fragments thereof, that bind to such sequences. Linkers may, but need not, be used to separate such CAR sequence(s) and/or antibody sequence(s) from the CAR sequence(s) and/or each other. Such modulating agents may generally be used within methods in which it is desirable to simultaneously disrupt a function mediated by multiple adhesion molecules. As used herein, an "adhesion molecule" is any molecule that mediates cell adhesion via a receptor on a cell's surface. Adhesion molecules include cell adhesion proteins (*e.g.*, other members of the cadherin gene superfamily, such as N-cadherin and E-cadherin); integrins; extracellular matrix proteins such as laminin, fibronectin, collagens, vitronectin, entactin and tenascin; and members of the immunoglobulin supergene family, such as N-CAM. Preferred CAR sequences for inclusion within a modulating agent include the classical cadherin CAR sequence His-Ala-Val (HAV); Arg-Gly-Asp (RGD), which is bound by integrins (*see* Cardarelli et al., *J. Biol. Chem.* 267:23159-64, 1992); Tyr-Ile-Gly-Ser-Arg (YIGSR; SEQ ID NO:\_\_), which is bound by  $\alpha 6 \beta 1$  integrin; KYSFNYDGSE (SEQ ID NO:\_\_), which is bound by N-CAM; the N-CAM heparin sulfate-binding site IWKHKGRDVILKKDVRF (SEQ ID NO:\_\_); the putative claudin CAR sequence IYSY (SEQ ID NO:\_\_) and/or the occludin CAR sequence LYHY (SEQ ID NO:\_\_).

Other preferred CAR sequences for inclusion within a modulating agent include OB-cadherin CAR sequences such as DDK, IDDK (SEQ ID NO:\_\_) DDKS (SEQ ID NO:\_\_), VIDDK (SEQ ID NO:\_\_), IDDKS (SEQ ID NO:\_\_), VIDDKS (SEQ ID NO:\_\_), DDKSG (SEQ ID NO:\_\_), IDDKSG (SEQ ID NO:\_\_), VIDDKSG (SEQ ID NO:\_\_), FVIDDK (SEQ ID NO:\_\_), FVIDDKS (SEQ ID NO:\_\_), FVIDDKSG (SEQ ID NO:\_\_), IFVIDDK (SEQ ID NO:\_\_), IFVIDDKS (SEQ ID NO:\_\_), IFVIDDKSG (SEQ ID NO:\_\_), EEY, IEY (SEQ ID NO:\_\_), EEYT (SEQ ID NO:\_\_), VIEY (SEQ ID NO:\_\_), IEY (SEQ ID NO:\_\_), VIEYT (SEQ ID NO:\_\_), EEYTG (SEQ ID NO:\_\_), IEYTG (SEQ ID NO:\_\_), VIEYTG (SEQ ID NO:\_\_), FVIEY (SEQ ID NO:\_\_), FVIEYT (SEQ ID NO:\_\_), FVIEYTG (SEQ ID NO:\_\_),



FFVIEEY (SEQ ID NO:\_\_\_), FFVIEEYT (SEQ ID NO:\_\_\_), FFVIEEYTG (SEQ ID NO:\_\_\_), EAQ, VEAQ (SEQ ID NO:\_\_\_), EAQT (SEQ ID NO:\_\_\_), SVEAQ (SEQ ID NO:\_\_\_), VEAQT (SEQ ID NO:\_\_\_), SVEAQT (SEQ ID NO:\_\_\_), EAQTG (SEQ ID NO:\_\_\_), VEAQTG (SEQ ID NO:\_\_\_), SVEAQTG (SEQ ID NO:\_\_\_), FSVEAQ (SEQ ID NO:\_\_\_), FSVEAQT (SEQ ID NO:\_\_\_), FSVEAQTG (SEQ ID NO:\_\_\_), YFSVEAQ (SEQ ID NO:\_\_\_), YFSVEAQT (SEQ ID NO:\_\_\_) and YFSVEAQTG (SEQ ID NO:\_\_\_).

Still further CAR sequences that may be included in a modulating agent are cadherin-5 CAR sequences such as DAE, VDAE (SEQ ID NO:\_\_\_), DAET (SEQ ID NO:\_\_\_), RVDAE (SEQ ID NO:\_\_\_), VDAET (SEQ ID NO:\_\_\_), RVDAET (SEQ ID NO:\_\_\_), DAETG (SEQ ID NO:\_\_\_), VDAETG (SEQ ID NO:\_\_\_), RVDAETG (SEQ ID NO:\_\_\_), FRVDAE (SEQ ID NO:\_\_\_), FRVDAET (SEQ ID NO:\_\_\_), FRVDAETG (SEQ ID NO:\_\_\_), VFRVDAE (SEQ ID NO:\_\_\_), VFRVDAET (SEQ ID NO:\_\_\_) and VFRVDAETG (SEQ ID NO:\_\_\_).

Using linkers, such modulating agents may form linear or branched structures. Within one embodiment, modulating agents having a branched structure comprise four different CAR sequences, such as IFVIDDKSG (SEQ ID NO:\_\_\_), RGD, YIGSR (SEQ ID NO:\_\_\_) and HAV. Bi-functional modulating agents that comprise a desmosomal cadherin CAR sequence joined via a linker to a classical cadherin CAR sequence are also preferred for certain embodiments. As noted above, linkers preferably produce a distance between CAR sequences ranging from 0.1 to 10,000 nm, more preferably ranging from 0.1-400 nm. A separation distance between recognition sites may generally be determined according to the desired function of the modulating agent.

The total number of CAR sequences (including the desmosomal cadherin CAR sequence, with or without other CAR sequences derived from one or more different adhesion molecules) present within a modulating agent may range from 1 to a large number, such as 100, preferably from 1 to 10, and more preferably from 1 to 5. Peptide modulating agents comprising multiple CAR sequences typically contain from 6 to about 1000 amino acid residues, preferably from 6 to 50 residues. When non-

peptide linkers are employed, each CAR sequence of the modulating agent is present within a peptide that generally ranges in size from 3 to 50 residues in length, preferably from 4 to 25 residues, and more preferably from 5 to 15 residues.

As noted above, modulating agents may be polypeptides or salts thereof, containing only amino acid residues linked by peptide bonds, or may contain non-peptide regions, such as linkers. Peptide regions of a modulating agent may comprise residues of L-amino acids, D-amino acids, or any combination thereof. Amino acids may be from natural or non-natural sources, provided that at least one amino group and at least one carboxyl group are present in the molecule;  $\alpha$ - and  $\beta$ -amino acids are generally preferred. The 20 L-amino acids commonly found in proteins are identified herein by the conventional three-letter or one-letter abbreviations, and the corresponding D-amino acids are designated by a lower case one letter symbol.

A modulating agent may also contain rare amino acids (such as 4-hydroxyproline or hydroxylysine), organic acids or amides and/or derivatives of common amino acids, such as amino acids having the C-terminal carboxylate esterified (*e.g.*, benzyl, methyl or ethyl ester) or amidated and/or having modifications of the N-terminal amino group (*e.g.*, acetylation or alkoxycarbonylation), with or without any of a wide variety of side-chain modifications and/or substitutions (*e.g.*, methylation, benzylation, t-butylation, tosylation, alkoxycarbonylation, and the like). Preferred derivatives include amino acids having a C-terminal amide group. Residues other than common amino acids that may be present with a modulating agent include, but are not limited to, 2-mercaptoaniline, 2-mercaptoproline, ornithine, diaminobutyric acid,  $\alpha$ -aminoadipic acid, m-aminomethylbenzoic acid and  $\alpha,\beta$ -diaminopropionic acid.

Peptide modulating agents (and peptide portions of modulating agents) as described herein may be synthesized by methods well known in the art, including chemical synthesis and recombinant DNA methods. For modulating agents up to about 50 residues in length, chemical synthesis may be performed using solution or solid phase peptide synthesis techniques, in which a peptide linkage occurs through the direct condensation of the  $\alpha$ -amino group of one amino acid with the  $\alpha$ -carboxy group of the other amino acid with the elimination of a water molecule. Peptide bond synthesis by

direct condensation, as formulated above, requires suppression of the reactive character of the amino group of the first and of the carboxyl group of the second amino acid. The masking substituents must permit their ready removal, without inducing breakdown of the labile peptide molecule.

5                   In solution phase synthesis, a wide variety of coupling methods and protecting groups may be used (*see* Gross and Meienhofer, eds., "The Peptides: Analysis, Synthesis, Biology," Vol. 1-4 (Academic Press, 1979); Bodansky and Bodansky, "The Practice of Peptide Synthesis," 2d ed. (Springer Verlag, 1994)). In addition, intermediate purification and linear scale up are possible. Those of ordinary  
10 skill in the art will appreciate that solution synthesis requires consideration of main chain and side chain protecting groups and activation method. In addition, careful segment selection is necessary to minimize racemization during segment condensation. Solubility considerations are also a factor.

                  Solid phase peptide synthesis uses an insoluble polymer for support  
15 during organic synthesis. The polymer-supported peptide chain permits the use of simple washing and filtration steps instead of laborious purifications at intermediate steps. Solid-phase peptide synthesis may generally be performed according to the method of Merrifield et al., *J. Am. Chem. Soc.* 85:2149, 1963, which involves assembling a linear peptide chain on a resin support using protected amino acids. Solid  
20 phase peptide synthesis typically utilizes either the Boc or Fmoc strategy. The Boc strategy uses a 1% cross-linked polystyrene resin. The standard protecting group for  $\alpha$ -amino functions is the tert-butyloxycarbonyl (Boc) group. This group can be removed with dilute solutions of strong acids such as 25% trifluoroacetic acid (TFA). The next Boc-amino acid is typically coupled to the amino acyl resin using  
25 dicyclohexylcarbodiimide (DCC). Following completion of the assembly, the peptide-resin is treated with anhydrous HF to cleave the benzyl ester link and liberate the free peptide. Side-chain functional groups are usually blocked during synthesis by benzyl-derived blocking groups, which are also cleaved by HF. The free peptide is then extracted from the resin with a suitable solvent, purified and characterized. Newly  
30 synthesized peptides can be purified, for example, by gel filtration, HPLC, partition

chromatography and/or ion-exchange chromatography, and may be characterized by, for example, mass spectrometry or amino acid sequence analysis. In the Boc strategy, C-terminal amidated peptides can be obtained using benzhydrylamine or methylbenzhydrylamine resins, which yield peptide amides directly upon cleavage with  
5 HF.

In the procedures discussed above, the selectivity of the side-chain blocking groups and of the peptide-resin link depends upon the differences in the rate of acidolytic cleavage. Orthogonal systems have been introduced in which the side-chain blocking groups and the peptide-resin link are completely stable to the reagent used to  
10 remove the  $\alpha$ -protecting group at each step of the synthesis. The most common of these methods involves the 9-fluorenylmethyloxycarbonyl (Fmoc) approach. Within this method, the side-chain protecting groups and the peptide-resin link are completely stable to the secondary amines used for cleaving the N- $\alpha$ -Fmoc group. The side-chain protection and the peptide-resin link are cleaved by mild acidolysis. The repeated  
15 contact with base makes the Merrifield resin unsuitable for Fmoc chemistry, and p-alkoxybenzyl esters linked to the resin are generally used. Deprotection and cleavage are generally accomplished using TFA.

Those of ordinary skill in the art will recognize that, in solid phase synthesis, deprotection and coupling reactions must go to completion and the side-chain  
20 blocking groups must be stable throughout the entire synthesis. In addition, solid phase synthesis is generally most suitable when peptides are to be made on a small scale.

Acetylation of the N-terminus can be accomplished by reacting the final peptide with acetic anhydride before cleavage from the resin. C-amidation is accomplished using an appropriate resin such as methylbenzhydrylamine resin using the  
25 Boc technology.

Following synthesis of a linear peptide, with or without N-acetylation and/or C-amidation, cyclization may be achieved if desired by any of a variety of techniques well known in the art. Within one embodiment, a bond may be generated between reactive amino acid side chains. For example, a disulfide bridge may be  
30 formed from a linear peptide comprising two thiol-containing residues by oxidizing the

peptide using any of a variety of methods. Within one such method, air oxidation of thiols can generate disulfide linkages over a period of several days using either basic or neutral aqueous media. The peptide is used in high dilution to minimize aggregation and intermolecular side reactions. This method suffers from the disadvantage of being  
 5 slow but has the advantage of only producing H<sub>2</sub>O as a side product. Alternatively, strong oxidizing agents such as I<sub>2</sub> and K<sub>3</sub>Fe(CN)<sub>6</sub> can be used to form disulfide linkages. Those of ordinary skill in the art will recognize that care must be taken not to oxidize the sensitive side chains of Met, Tyr, Trp or His. Cyclic peptides produced by this method require purification using standard techniques, but this oxidation is  
 10 applicable at acid pHs. Oxidizing agents also allow concurrent deprotection/oxidation of suitable S-protected linear precursors to avoid premature, nonspecific oxidation of free cysteine.

DMSO, unlike I<sub>2</sub> and K<sub>3</sub>Fe(CN)<sub>6</sub>, is a mild oxidizing agent which does not cause oxidative side reactions of the nucleophilic amino acids mentioned above.  
 15 DMSO is miscible with H<sub>2</sub>O at all concentrations, and oxidations can be performed at acidic to neutral pHs with harmless byproducts. Methyltrichlorosilane-diphenylsulfoxide may alternatively be used as an oxidizing agent, for concurrent deprotection/oxidation of S-Acm, S-Tacm or S-t-Bu of cysteine without affecting other nucleophilic amino acids. There are no polymeric products resulting from  
 20 intermolecular disulfide bond formation. Suitable thiol-containing residues for use in such oxidation methods include, but are not limited to, cysteine, β,β-dimethyl cysteine (penicillamine or Pen), β,β-tetramethylene cysteine (Tmc), β,β-pentamethylene cysteine (Pmc), β-mercaptopropionic acid (Mpr), β,β-pentamethylene-β-mercaptopropionic acid (Pmp), 2-mercaptobenzene, 2-mercaptoaniline and 2-mercaptoproline. Peptides  
 25 containing such residues are illustrated by the following representative formulas, in which the desmosomal cadherin is a desmoglein, the underlined portion is cyclized, N-acetyl groups are indicated by N-Ac and C-terminal amide groups are represented by -NH<sub>2</sub>:

30 i) N-Ac-Cys-Asn-Gln-Lys-Cys-NH<sub>2</sub> (SEQ ID NO:     )

- 5
- ii) N-Ac-Cys-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- iii) N-Ac-Cys-Ile-Asn-Gln-Lys-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- iv) N-Ac-Cys-Asn-Gln-Lys-Thr-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- v) N-Ac-Cys-Ile-Asn-Gln-Lys-Thr-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- 10 vi) N-Ac-Cys-Asn-Gln-Lys-Thr-Cys-OH (SEQ ID NO: \_\_)
- vii) H-Cys-Ile-Asn-Gln-Lys-Thr-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- viii) N-Ac-Cys-Asn-Gln-Lys-Pen-NH<sub>2</sub> (SEQ ID NO: \_\_)
- 15 ix) N-Ac-Cys-Phe-Val-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub>  
(SEQ ID NO: \_\_)
- x) N-Ac-Cys-Ile-Phe-Val-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub>  
20 (SEQ ID NO: \_\_)
- xi) N-Ac-Ile-Tmc-Val-Ile-Asn-Gln-Lys-Thr-Cys-Glu-NH<sub>2</sub> (SEQ ID NO: \_\_)
- xii) N-Ac-Ile-Pmc-Val-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- 25 xiii) Mpr-Val-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
- xiv) Pmp-Val-Ile-Asn-Gln-Lys-Thr-Gly-Cys-NH<sub>2</sub> (SEQ ID NO: \_\_)
-

It will be readily apparent to those of ordinary skill in the art that, within each of these representative formulas, any of the above thiol-containing residues may be employed in place of one or both of the thiol-containing residues recited. Similar formulas comprising different nonclassical cadherin CAR sequences may be generated  
5 by those of ordinary skill in the art, based on the CAR sequences provided herein.


Within another embodiment, cyclization may be achieved by amide bond formation. For example, a peptide bond may be formed between terminal functional groups (*i.e.*, the amino and carboxy termini of a linear peptide prior to cyclization). One such cyclic peptide comprising a desmoglein CAR sequence is INQKTG (SEQ ID  
10 NO: \_\_) with or without an N-terminal acetyl group and/or a C-terminal amide. Within another such embodiment, the linear peptide comprises a D-amino acid (*e.g.*, NQKtS; SEQ ID NO: \_\_). Alternatively, cyclization may be accomplished by linking one terminus and a residue side chain or using two side chains, as in KNQKD (SEQ ID NO: \_\_) or KINQKTGD (SEQ ID NO: \_\_), with or without an N-terminal acetyl group  
15 and/or a C-terminal amide. Residues capable of forming a lactam bond include lysine, ornithine (Orn),  $\alpha$ -amino adipic acid, m-aminomethylbenzoic acid,  $\alpha,\beta$ -diaminopropionic acid, glutamate or aspartate.

Methods for forming amide bonds are well known in the art and are based on well established principles of chemical reactivity. Within one such method,  
20 carbodiimide-mediated lactam formation can be accomplished by reaction of the carboxylic acid with DCC, DIC, EDAC or DCCI, resulting in the formation of an O-acylurea that can be reacted immediately with the free amino group to complete the cyclization. The formation of the inactive N-acylurea, resulting from O $\rightarrow$ N migration, can be circumvented by converting the O-acylurea to an active ester by reaction with an  
25 N-hydroxy compound such as 1-hydroxybenzotriazole, 1-hydroxysuccinimide, 1-hydroxynorbornene carboxamide or ethyl 2-hydroximino-2-cyanoacetate. In addition to minimizing O $\rightarrow$ N migration, these additives also serve as catalysts during cyclization and assist in lowering racemization. Alternatively, cyclization can be performed using the azide method, in which a reactive azide intermediate is generated from an alkyl ester  
30 via a hydrazide. Hydrazinolysis of the terminal ester necessitates the use of a t-butyl

group for the protection of side chain carboxyl functions in the acylating component. This limitation can be overcome by using diphenylphosphoryl acid (DPPA), which furnishes an azide directly upon reaction with a carboxyl group. The slow reactivity of azides and the formation of isocyanates by their disproportionation restrict the usefulness of this method. The mixed anhydride method of lactam formation is widely used because of the facile removal of reaction by-products. The anhydride is formed upon reaction of the carboxylate anion with an alkyl chloroformate or pivaloyl chloride. The attack of the amino component is then guided to the carbonyl carbon of the acylating component by the electron donating effect of the alkoxy group or by the steric bulk of the pivaloyl chloride t-butyl group, which obstructs attack on the wrong carbonyl group. Mixed anhydrides with phosphoric acid derivatives have also been successfully used. Alternatively, cyclization can be accomplished using activated esters. The presence of electron withdrawing substituents on the alkoxy carbon of esters increases their susceptibility to aminolysis. The high reactivity of esters of p-nitrophenol, N-hydroxy compounds and polyhalogenated phenols has made these "active esters" useful in the synthesis of amide bonds. The last few years have witnessed the development of benzotriazolyloxytris-(dimethylamino)phosphonium hexafluorophosphate (BOP) and its congeners as advantageous coupling reagents. Their performance is generally superior to that of the well established carbodiimide amide bond formation reactions.

Within a further embodiment, a thioether linkage may be formed between the side chain of a thiol-containing residue and an appropriately derivatized  $\alpha$ -amino acid. By way of example, a lysine side chain can be coupled to bromoacetic acid through the carbodiimide coupling method (DCC, EDAC) and then reacted with the side chain of any of the thiol containing residues mentioned above to form a thioether linkage. In order to form dithioethers, any two thiol containing side-chains can be reacted with dibromoethane and diisopropylamine in DMF. Examples of thiol-containing linkages are shown below:





Cyclization may also be achieved using  $\delta_1, \delta_1$ -Ditryptophan (e.g., Ac-  
 5 Trp-Asn-Gln-Lys-Trp-OMe) (SEQ ID NO:     ).

The structures and formulas recited herein are provided solely for the purpose of illustration, and are not intended to limit the scope of the cyclic peptides described herein.

For longer modulating agents, recombinant methods are preferred for  
 10 synthesis. Within such methods, all or part of a modulating agent can be synthesized in living cells, using any of a variety of expression vectors known to those of ordinary skill in the art to be appropriate for the particular host cell. Suitable host cells may include bacteria, yeast cells, mammalian cells, insect cells, plant cells, algae and other animal cells (e.g., hybridoma, CHO, myeloma). The DNA sequences expressed in this  
 15 manner may encode portions of a desmosomal cadherin or other adhesion molecule, or may encode a peptide comprising a desmosomal cadherin analogue or an antibody fragment that specifically binds to a desmosomal cadherin CAR sequence. Such DNA sequences may be prepared based on known cDNA or genomic sequences, or from sequences isolated by screening an appropriate library with probes designed based on  
 20 the sequences of known desmosomal cadherins. Such screens may generally be performed as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989 (and references cited therein). Polymerase chain reaction (PCR) may also be employed, using oligonucleotide primers in methods well known in the art, to isolate nucleic acid  
 25 molecules encoding all or a portion of an endogenous adhesion molecule. To generate a

nucleic acid molecule encoding a desired modulating agent, an endogenous cadherin sequence may be modified using well known techniques. For example, portions encoding one or more CAR sequences may be joined, with or without separation by nucleic acid regions encoding linkers, as discussed above. Alternatively, portions of the desired nucleic acid sequences may be synthesized using well known techniques, and then ligated together to form a sequence encoding the modulating agent.

As noted above, polynucleotides may also function as modulating agents. In general, such polynucleotides should be formulated to permit expression of a polypeptide modulating agent following administration to a mammal. Such formulations are particularly useful for therapeutic purposes, as described below. Those of ordinary skill in the art will appreciate that there are many ways to achieve expression of a polynucleotide within a mammal, and any suitable method may be employed. For example, a polynucleotide may be incorporated into a viral vector such as, but not limited to, adenovirus, adeno-associated virus, retrovirus, or vaccinia or other pox virus (*e.g.*, avian pox virus). Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transfected cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an antibody, by methods known to those of ordinary skill in the art. Other formulations for polynucleotides for therapeutic purposes include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome (*i.e.*, an artificial membrane vesicle). The preparation and use of such systems is well known in the art.

As noted above, a modulating agent may additionally, or alternatively, comprise a substance such as an antibody or antigen-binding fragment thereof, that specifically binds to a desmosomal cadherin CAR sequence. As used herein, a substance is said to "specifically bind" to a desmosomal cadherin CAR sequence (with

or without flanking amino acids) if it reacts at a detectable level with a peptide containing that sequence, and does not react detectably with peptides containing a different CAR sequence or a sequence in which the order of amino acid residues in the cadherin CAR sequence and/or flanking sequence is altered. Such antibody binding  
5 properties may generally be assessed using an ELISA, which may be readily performed by those of ordinary skill in the art and is described, for example, by Newton et al., *Develop. Dynamics* 197:1-13, 1993.

Polyclonal and monoclonal antibodies may be raised against a desmosomal cadherin CAR sequence using conventional techniques. See, e.g., Harlow  
10 and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the CAR sequence is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep or goats). The smaller immunogens (i.e., less than about 20 amino acids) should be joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. Following one  
15 or more injections, the animals are bled periodically. Polyclonal antibodies specific for the CAR sequence may then be purified from such antisera by, for example, affinity chromatography using the modulating agent or antigenic portion thereof coupled to a suitable solid support.

Monoclonal antibodies specific for a desmosomal cadherin sequence  
20 may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity from spleen cells obtained from an animal immunized as described above. The spleen cells are immortalized by, for example, fusion with a myeloma cell  
25 fusion partner, preferably one that is syngeneic with the immunized animal. Single colonies are selected and their culture supernatants tested for binding activity against the modulating agent or antigenic portion thereof. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing  
30 hybridoma colonies, with or without the use of various techniques known in the art to

enhance the yield. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation and extraction. Antibodies having the desired activity may generally be identified using immunofluorescence analyses of tissue sections, cell or other samples where the target  
5 cadherin is localized.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane,  
10 *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988; see especially page 309) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns (Harlow and Lane, 1988, pages 628-29).

#### 15 EVALUATION OF MODULATING AGENT ACTIVITY

Modulating agents as described above are capable of modulating one or more desmosomal cadherin-mediated functions. An initial screen for such activity may be performed by evaluating the ability of a modulating agent to bind to a desmosomal  
20 cadherin using any binding assay known to those of ordinary skill in the art. For example, a Pharmacia Biosensor machine may be used, as discussed in Jonsson et al., *Biotechniques* 11:520-27, 1991. For example, a modulating agent may comprise a CAR sequence that binds to a desmosomal cadherin. A specific example of a technology that measures the interaction of peptides with molecules can be found in Williams et al., *J.*  
25 *Biol. Chem.* 272, 22349-22354, 1997. Alternatively, real-time BIA (Biomolecular Interaction Analysis) uses the optical phenomenon surface plasmon resonance to monitor biomolecular interactions. The detection depends upon changes in the mass concentration of macromolecules at the biospecific interface, which in turn depends upon the immobilization of test molecule or peptide (referred to as the ligand) to the  
30 surface of a Biosensor chip, followed by binding of the interacting molecule (referred to

as the analyte) to the ligand. Binding to the chip is measured in real-time in arbitrary units of resonance (RU).

By way of example, surface plasmon resonance experiments may be carried out using a BIAcore X™ Biosensor (Pharmacia Ltd., BIAcore, Uppsala, Sweden). Parallel flow cells of CM 5 sensor chips may be derivatized, using the amine coupling method, with streptavidin (200µg/ml) in 10mM Sodium Acetate, pH 4.0, according to the manufacturer's protocol. Approximately 2100-2600 resonance units (RU) of ligand may be immobilized, corresponding to a concentration of about 2.1-2.6 ng/mm<sup>2</sup>. The chips may then coated be with desmosomal cadherin derivatized to biotin.

Any non-specifically bound protein is removed.

To determine binding, test analytes (*e.g.*, peptides containing a desmosomal cadherin CAR sequence) may be placed in running buffer and passed simultaneously over test and control flow cells. After a period of free buffer flow, any analyte remaining bound to the surface may be removed with, for example, a pulse of 0.1% SDS bringing the signal back to baseline. Specific binding to the derivatized sensor chips may be determined automatically by the system by subtraction of test from control flow cell responses. In general, a modulating agent binds to a desmosomal cadherin at a detectable level within such as assay. The level of binding is preferably at least that observed for the full length desmosomal cadherin under similar conditions.

The ability to modulate a desmosomal cadherin-mediated function may be evaluated using any of a variety of *in vitro* assays designed to measure the effect of the peptide on a response that is generally mediated by the desmosomal cadherin. As noted above, modulating agents may be capable of enhancing or inhibiting a desmosomal cadherin-mediated function.

Certain desmosomal cadherins are associated with adhesion of particular cell types (*e.g.*, epithelial cells). The ability of an agent to modulate cell adhesion may generally be evaluated *in vitro* by assaying the effect on adhesion between appropriate cells. In general, a modulating agent is an inhibitor of cell adhesion if contact of the test cells with the modulating agent results in a discernible disruption of cell adhesion.

Modulating agents that enhance cell adhesion (*e.g.*, agents comprising multiple

desmosomal cadherin CAR sequences and/or desmosomal cadherin CAR sequences linked to a support material) are considered to be modulators of cell adhesion if they are capable of promoting cell adhesion, as judged by plating assays to assess cell adhesion to a modulating agent attached to a support material, such as tissue culture plastic.

5                Within certain cell adhesion assays, the addition of a modulating agent to cells that express a desmosomal cadherin results in disruption of cell adhesion. A "desmosomal cadherin-expressing cell," as used herein, may be any type of cell that expresses a desmosomal cadherin at a detectable level, using standard techniques such as immunocytochemical protocols (*e.g.*, Blaschuk and Farookhi, *Dev. Biol.* 136:564-  
10 567, 1989). For example, such cells may be plated under standard conditions that, in the absence of modulating agent, permit cell adhesion. In the presence of modulating agent (*e.g.*, 1 mg/mL), disruption of cell adhesion may be determined visually within 24 hours, by observing retraction of the cells from one another and the substratum.

              Suitable cells for use within such assays may be any of a variety of cells  
15 that express the desmosomal cadherin of interest. Certain cells express one or more cadherins endogenously. In general, MDCK cells or keratinocytes may be used to evaluate desmocollin- or desmoglein-mediated cell adhesion. It will be apparent that other cells may also be used within such assays, provided that the cells express the desmosomal cadherin of interest.

20                Alternatively, cells that do not naturally express a cadherin may be used within such assays. Such cells may be stably transfected with a polynucleotide (*e.g.*, a cDNA) encoding one or more cadherins of interest, such that the cadherin or cadherins are expressed on the surface of the cell. For example, as noted above, both a desmoglein and a desmocollin may be required for optimal cell adhesion, and such  
25 assays may be performed using cells transformed with polynucleotides encoding both of these desmosomal cadherins. Expression of the cadherin(s) may be confirmed by assessing adhesion of the transfected cells, in conjunction with immunocytochemical techniques using antibodies directed against the cadherin of interest. The stably transfected cells that aggregate, as judged by light microscopy, following transfection  
30 express sufficient levels of the desmosomal cadherin(s). Preferred cells for use in such

assays include L cells, which do not detectably adhere and do not express any cadherin (Nagafuchi et al., *Nature* 329:341-343, 1987). Following transfection of L cells with a cDNA encoding one or more cadherins, aggregation is observed. Modulating agents that detectably inhibit such aggregation may be used to modulate functions mediated by the cadherin. Such assays have been used for numerous nonclassical cadherins, including OB-cadherin (Okazaki et al., *J. Biol. Chem.* 269:12092-98, 1994), cadherin-5 (Breier et al., *Blood* 87:630-641, 1996), cadherin-6 (Mbalaviele et al., *J. Cell. Biol.* 141:1467-1476, 1998), cadherin-8 (Kido et al., *Genomics* 48:186-194, 1998), cadherin-15 (Shimoyama et al., *J. Biol. Chem.* 273:10011-10018, 1998), PB-cadherin (Sugimoto et al., *J. Biol. Chem.* 271:11548-11556, 1996), LI-cadherin (Kreft et al., *J. Cell. Biol.* 136:1109-1121, 1997), protocadherin 42 and 43 (Sano et al., *EMBO J.* 12:2249-2256, 1993) and desmosomal cadherins (Marcozzi et al., *J. Cell. Sci.* 111:495-509, 1998; Tselepis et al., *Proc. Natl. Acad. Sci. USA* 95:8064-8069, 1998). It will be apparent to those of ordinary skill in the art that assays may be performed in a similar manner for other cadherins.

Transfection of cells for use in cell adhesion assays may be performed using standard techniques and published cadherin sequences. For example, sequences of desmosomal cadherins may be found within references cited herein and in the GenBank database. GenBank accession numbers for certain desmosomal cadherins include: X56654 (human desmoglein 1); Z26317 and S64273 (human desmoglein 2); X72925 (human desmocollin 1); X56807 (human desmocollin 2); X83929 (human desmocollin 3); and D17427 (human desmocollin 4).

By way of example, an assay for evaluating a modulating agent for the ability to inhibit a desmosomal cadherin mediated function may evaluate the effect of a modulating agent on the electrical resistance across a monolayer of cells. For example, Madin Darby canine kidney (MDCK) cells can be exposed to the modulating agent dissolved in medium (e.g., at a final concentration of 0.5 mg/ml for a period of 24 hours). The effect on electrical resistance can be measured using standard techniques. This assay evaluates the effect of a modulating agent on tight junction formation in

epithelial cells. In general, the presence of 500  $\mu\text{g/mL}$  modulating agent should result in a statistically significant decrease in electrical resistance after 24 hours.

Other assays involve visual evaluation of cell adhesion, using standard methods. In general, a modulating agent that is derived from a particular desmosomal cadherin CAR sequence (*i.e.*, comprises such a CAR sequence, or an analog or mimetic thereof, or an antibody that specifically recognizes such a CAR sequence) and that detectably modulates adhesion of cells that express the same desmosomal cadherin is considered to modulate a function mediated by the desmosomal cadherin.

Other assays may be used to assess the effect of a modulating agent on specific desmosomal cadherin-mediated functions. For example, modulating agents that inhibit interactions of desmogleins and desmocollins may enhance skin permeability. This ability may be assessed by evaluating, for example, the effect of a modulating agent on permeability of adherent epithelial and/or endothelial cell layers (*e.g.*, human skin). Such skin may be derived from a natural source or may be synthetic. Human abdominal skin for use in such assays may generally be obtained from humans at autopsy within 24 hours of death. Briefly, a modulating agent (*e.g.*, 500  $\mu\text{g/mL}$ ) and a test marker (*e.g.*, the fluorescent markers Oregon Green™ and Rhodamine Green™ Dextran) may be dissolved in a sterile buffer (*e.g.*, phosphate buffer, pH 7.2), and the ability of the marker to penetrate through the skin and into a receptor fluid (*e.g.*, phosphate buffer) may be measured using a Franz Cell apparatus (Franz, *Curr. Prob. Dermatol.* 7:58-68, 1978; Franz, *J. Invest. Dermatol.* 64:190-195, 1975). The penetration of the markers through the skin may be assessed at, for example, 6, 12, 24, 36, and 48 hours after the start of the experiment. In general, a modulating agent that enhances the permeability of human skin results in a statistically significant increase in the amount of marker in the receptor compartment after 6-48 hours in the presence of 500  $\mu\text{g/mL}$  modulating agent.

#### MODULATING AGENT MODIFICATION AND FORMULATIONS

A modulating agent as described herein may, but need not, be linked to one or more additional molecules. In particular, as discussed below, it may be



beneficial for certain applications to link multiple modulating agents (which may, but need not, be identical) to a support material, such as a single molecule (*e.g.*, keyhole limpet hemocyanin) or a solid support, such as a polymeric matrix (which may be formulated as a membrane or microstructure, such as an ultra thin film), a container surface (*e.g.*, the surface of a tissue culture plate or the interior surface of a bioreactor), or a bead or other particle, which may be prepared from a variety of materials including glass, plastic or ceramics. For certain applications, biodegradable support materials are preferred, such as cellulose and derivatives thereof, collagen, spider silk or any of a variety of polyesters (*e.g.*, those derived from hydroxy acids and/or lactones) or sutures (see U.S. Patent No. 5,245,012). Within certain embodiments, modulating agents and molecules comprising other CAR sequence(s) (*e.g.*, an HAV or RGD sequence, or an OB-cadherin or cadherin-5 CAR sequence as described above) may be attached to a support such as a polymeric matrix, preferably in an alternating pattern.

Suitable methods for linking a modulating agent to a support material will depend upon the composition of the support and the intended use, and will be readily apparent to those of ordinary skill in the art. Attachment may generally be achieved through noncovalent association, such as adsorption or affinity or, preferably, via covalent attachment (which may be a direct linkage between a modulating agent and functional groups on the support, or may be a linkage by way of a cross-linking agent). Attachment of a modulating agent by adsorption may be achieved by contact, in a suitable buffer, with a solid support for a suitable amount of time. The contact time varies with temperature, but is generally between about 5 seconds and 1 day, and typically between about 10 seconds and 1 hour.

Covalent attachment of a modulating agent to a molecule or solid support may generally be achieved by first reacting the support material with a bifunctional reagent that will also react with a functional group, such as a hydroxyl or amino group, on the modulating agent. For example, a modulating agent may be bound to an appropriate polymeric support or coating using benzoquinone, by condensation of an aldehyde group on the support with an amine and an active hydrogen on the modulating agent or by condensation of an amino group on the support with a

carboxylic acid on the modulating agent. A preferred method of generating a linkage is via amino groups using glutaraldehyde. A modulating agent may be linked to cellulose via ester linkages. Similarly, amide linkages may be suitable for linkage to other molecules such as keyhole limpet hemocyanin or other support materials. Multiple  
5 modulating agents and/or molecules comprising other CAR sequences may be attached, for example, by random coupling, in which equimolar amounts of such molecules are mixed with a matrix support and allowed to couple at random.

Although modulating agents as described herein may preferentially bind to specific tissues or cells, and thus may be sufficient to target a desired site *in vivo*, it  
10 may be beneficial for certain applications to include an additional targeting agent. Accordingly, a targeting agent may also, or alternatively, be linked to a modulating agent to facilitate targeting to one or more specific tissues. As used herein, a "targeting agent," may be any substance (such as a compound or cell) that, when linked to a modulating agent enhances the transport of the modulating agent to a target tissue,  
15 thereby increasing the local concentration of the modulating agent. Targeting agents include antibodies or fragments thereof, receptors, ligands and other molecules that bind to cells of, or in the vicinity of, the target tissue. Known targeting agents include serum hormones, antibodies against cell surface antigens, lectins, adhesion molecules, tumor cell surface binding ligands, steroids, cholesterol, lymphokines, fibrinolytic enzymes  
20 and those drugs and proteins that bind to a desired target site. Among the many monoclonal antibodies that may serve as targeting agents are anti-TAC, or other interleukin-2 receptor antibodies; 9.2.27 and NR-ML-05, reactive with the 250 kilodalton human melanoma-associated proteoglycan; and NR-LU-10, reactive with a pancarcinoma glycoprotein. An antibody targeting agent may be an intact (whole)  
25 molecule, a fragment thereof, or a functional equivalent thereof. Examples of antibody fragments are F(ab')<sub>2</sub>, -Fab', Fab and F[v] fragments, which may be produced by conventional methods or by genetic or protein engineering. Linkage is generally covalent and may be achieved by, for example, direct condensation or other reactions, or by way of bi- or multi-functional linkers.

For certain embodiments, it may be beneficial to also, or alternatively, link a drug to a modulating agent. As used herein, the term "drug" refers to any bioactive agent intended for administration to a mammal to prevent or treat a disease or other undesirable condition. Drugs include hormones, growth factors, proteins, peptides and other compounds. The use of certain specific drugs within the context of the present invention is discussed below.

Modulating agents as described herein may be present within a pharmaceutical composition. A pharmaceutical composition comprises one or more modulating agents in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such compositions may comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide) and/or preservatives. Within yet other embodiments, compositions of the present invention may be formulated as a lyophilizate. One or more modulating agents (alone or in combination with a targeting agent and/or drug) may, but need not, be encapsulated within liposomes using well known technology. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous, or intramuscular administration.

For certain embodiments, as discussed below, a pharmaceutical composition may further comprise a modulator of cell adhesion that is mediated by one or more molecules other than the particular desmosomal cadherin. Such modulators may generally be prepared as described above, using one or more CAR sequences and/or antibodies thereto. Such compositions are particularly useful for situations in which it is desirable to inhibit cell adhesion mediated by multiple cell adhesion molecules, such as other members of the cadherin gene superfamily such as the classical cadherins (*e.g.*, N-cadherin and E-cadherin); nonclassical cadherins, integrins; occludin;

claudins; N-CAM and/or extracellular matrix proteins such as laminin, fibronectin, collagens, vitronectin, entactin and tenascin.

A pharmaceutical composition may also, or alternatively, contain one or more drugs, which may be linked to a modulating agent or may be free within the composition. Virtually any drug may be administered in combination with a modulating agent as described herein, for a variety of purposes as described below. Examples of types of drugs that may be administered with a modulating agent include analgesics, anesthetics, antianginals, antifungals, antibiotics, anticancer drugs (*e.g.*, taxol or mitomycin C), antiinflammatories (*e.g.*, ibuprofen and indomethacin), anthelmintics, antidepressants, antidotes, antiemetics, antihistamines, antihypertensives, antimalarials, antimicrotubule agents (*e.g.*, colchicine or vinca alkaloids), antimigraine agents, antimicrobials, antipsychotics, antipyretics, antiseptics, anti-signaling agents (*e.g.*, protein kinase C inhibitors or inhibitors of intracellular calcium mobilization), antiarthritics, antithrombin agents, antituberculotics, antitussives, antivirals, appetite suppressants, cardioactive drugs, chemical dependency drugs, cathartics, chemotherapeutic agents, coronary, cerebral or peripheral vasodilators, contraceptive agents, depressants, diuretics, expectorants, growth factors, hormonal agents, hypnotics, immunosuppression agents, narcotic antagonists, parasympathomimetics, sedatives, stimulants, sympathomimetics, toxins (*e.g.*, cholera toxin), tranquilizers and urinary antiinfectives.

For imaging purposes, any of a variety of diagnostic agents may be incorporated into a pharmaceutical composition, either linked to a modulating agent or free within the composition. Diagnostic agents include any substance administered to illuminate a physiological function within a patient, while leaving other physiological functions generally unaffected. Diagnostic agents include metals, radioactive isotopes and radioopaque agents (*e.g.*, gallium, technetium, indium, strontium, iodine, barium, bromine and phosphorus-containing compounds), radiolucent agents, contrast agents, dyes (*e.g.*, fluorescent dyes and chromophores) and enzymes that catalyze a colorimetric or fluorometric reaction. In general, such agents may be attached using a variety of techniques as described above, and may be present in any orientation.

The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of modulating agent following administration). Such formulations may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a modulating agent dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane (*see, e.g.*, European Patent Application 710,491 A). Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of modulating agent release. The amount of modulating agent contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Pharmaceutical compositions of the present invention may be administered in a manner appropriate to the disease to be treated (or prevented). Appropriate dosages and a suitable duration and frequency of administration will be determined by such factors as the condition of the patient, the type and severity of the patient's disease and the method of administration. In general, an appropriate dosage and treatment regimen provides the modulating agent(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Within particularly preferred embodiments of the invention, a modulating agent or pharmaceutical composition as described herein may be administered at a dosage ranging from 0.001 to 50 mg/kg body weight, preferably from 0.1 to 20 mg/kg, on a regimen of single or multiple daily doses. For topical administration, a cream typically comprises an amount of modulating agent ranging from 0.00001% to 1%, preferably 0.0001% to 0.002%. Fluid compositions typically contain about 10 ng/ml to 5 mg/ml, preferably from about 10 µg to 2 mg/mL modulating agent. Appropriate dosages may generally be determined using experimental models and/or clinical trials. In general, the use of the minimum dosage that is sufficient to provide effective therapy is preferred. Patients may generally be

monitored for therapeutic effectiveness using assays suitable for the condition being treated or prevented, which will be familiar to those of ordinary skill in the art.

#### MODULATING AGENT METHODS OF USE

5                   In general, the modulating agents and compositions described herein may be used for modulating a function, such as cell adhesion, of desmosomal cadherin-expressing cells. Such modulation may be performed *in vitro* and/or *in vivo*, preferably in a mammal such as a human, using any method that contacts the desmosomal cadherin-expressing cell with the modulating agent. As noted above, modulating agents  
10 for purposes that involve the disruption of desmosomal cadherin-mediated cell adhesion may comprise a desmosomal cadherin CAR sequence, multiple desmosomal cadherin CAR sequences in close proximity and/or a substance (such as an antibody or an antigen-binding fragment thereof) that recognizes a desmosomal cadherin CAR sequence. When it is desirable to also disrupt cell adhesion mediated by other adhesion  
15 molecules, a modulating agent may additionally comprise one or more CAR sequences bound by such adhesion molecules (and/or antibodies or fragments thereof that bind such sequences), preferably separated from each other and from the desmosomal cadherin CAR sequence by linkers. As noted above, such linkers may or may not comprise one or more amino acids. For enhancing cell adhesion, a modulating agent  
20 may contain multiple desmosomal cadherin CAR sequences derived from one or more desmosomal cadherins or antibodies (or fragments), preferably separated by linkers, and/or may be linked to a single molecule or to a support material as described above. When it is desirable to also enhance cell adhesion mediated by other adhesion molecules, a modulating agent may additionally comprise one or more CAR sequences  
25 bound by such adhesion molecules (and/or antibodies or fragments thereof that bind such sequences), preferably separated from each other and from the desmosomal cadherin CAR sequence by linkers.

Certain methods involving the disruption of cell adhesion as described herein have an advantage over prior techniques in that they block tumor cell adhesion.  
30 As described in greater detail below, modulating agents as described herein may also be

used to disrupt or enhance cell adhesion in a variety of other contexts. Within each of the methods described herein, one or more modulating agents may generally be administered alone, or within a pharmaceutical composition. In each specific method described herein, as noted above, a targeting agent may be employed to increase the  
5 local concentration of modulating agent at the target site.

Within one aspect, methods are provided in which cell adhesion is diminished. In one such aspect, the present invention provides methods for reducing unwanted cellular adhesion in a mammal by administering a modulating agent as described herein. Unwanted cellular adhesion can occur, for example, between tumor  
10 cells, between tumor cells and normal cells or between normal cells as a result of surgery, injury, chemotherapy, disease, inflammation or other condition jeopardizing cell viability or function. In one particularly preferred embodiment, a modulating agent is further capable of disrupting cell adhesion mediated by multiple adhesion molecules. Such an agent may comprise, in addition to one or more desmosomal cadherin CAR  
15 sequences, CAR sequences such as the classical cadherin CAR sequence HAV sequence, an RGD sequence, which is bound by integrins, an OB-cadherin or cadherin-5 CAR sequence as described above, the occludin CAR sequence LYHY (SEQ ID NO:\_\_); and/or the claudin CAR sequence IYSY (SEQ ID NO:\_\_), preferably separated from the desmosomal cadherin CAR sequence via a linker. Alternatively, separate  
20 modulators of cell adhesion mediated by other adhesion molecules may be administered in conjunction with the modulating agent(s), either within the same pharmaceutical composition or separately.

Topical administration of the modulating agent(s) is generally preferred, but other means may also be employed. Preferably, a fluid composition for topical  
25 administration (comprising, for example, physiological saline) comprises an amount of modulating agent as described above, and more preferably from 10µg/mL to 1mg/mL. Creams may generally be formulated as described above. Topical administration in the surgical field may be given once at the end of surgery by irrigation of the wound or as an intermittent or continuous irrigation with the use of surgical drains in the post-  
30 operative period or by the use of drains specifically inserted in an area of inflammation,

injury or disease in cases where surgery does not need to be performed. Alternatively, parenteral or transcutaneous administration may be used to achieve similar results.

Certain modulating agents as provided herein may be used to facilitate transdermal drug delivery. Transdermal delivery of drugs is a convenient and non-invasive method that can be used to maintain relatively constant blood levels of a drug. In general, to facilitate drug delivery via the skin, it is necessary to perturb adhesion between the epithelial cells (keratinocytes) and the endothelial cells of the microvasculature. Using currently available techniques, only small, uncharged molecules may be delivered across skin *in vivo*. The methods described herein are not subject to the same degree of limitation. Accordingly, a wide variety of drugs may be transported across the epithelial and endothelial cell layers of skin, for systemic or topical administration. Such drugs may be delivered to melanomas or may enter the blood stream of the mammal for delivery to other sites within the body.

To enhance the delivery of a drug through the skin, a modulating agent as described herein and a drug are contacted with the skin surface. Multifunctional modulating agents comprising multiple nonclassical cadherin CAR sequences (*e.g.*, a desmosomal cadherin CAR sequence in combination with a CAR sequence derived from OB-cadherin or cadherin-5) may also be used. Such modulating agents may also, or alternatively, comprise the classical cadherin CAR sequence HAV, the fibronectin CAR sequence RGD, which is recognized by integrins, and/or the occludin CAR sequence LYHY (SEQ ID NO:\_\_). Alternatively, a separate modulator of cell adhesion may be administered in conjunction with the modulating agent(s), either within the same pharmaceutical composition or separately.

Contact may be achieved by direct application of the modulating agent, generally within a composition formulated as a cream or gel, or using any of a variety of skin contact devices for transdermal application (such as those described in European Patent Application No. 566,816 A; U.S. Patent No. 5,613,958; U.S. Patent No. 5,505,956). A skin patch provides a convenient method of administration (particularly for slow-release formulations). Such patches may contain a reservoir of modulating agent and drug separated from the skin by a membrane through which the drug diffuses.



Within other patch designs, the modulating agent and drug may be dissolved or suspended in a polymer or adhesive matrix that is then placed in direct contact with the patient's skin. The modulating agent and drug may then diffuse from the matrix into the skin. Modulating agent(s) and drug(s) may be contained within the same composition or skin patch, or may be separately administered, although administration at the same time and site is preferred. In general, the amount of modulating agent administered via the skin varies with the nature of the condition to be treated or prevented, but may vary as described above. Such levels may be achieved by appropriate adjustments to the device used, or by applying a cream formulated as described above. Transfer of the drug across the skin and to the target tissue may be predicted based on *in vitro* studies using, for example, a Franz cell apparatus, and evaluated *in vivo* by appropriate means that will be apparent to those of ordinary skill in the art. As an example, monitoring of the serum level of the administered drug over time provides an easy measure of the drug transfer across the skin.

Transdermal drug delivery as described herein is particularly useful in situations in which a constant rate of drug delivery is desired, to avoid fluctuating blood levels of a drug. For example, morphine is an analgesic commonly used immediately following surgery. When given intermittently in a parenteral form (intramuscular, intravenous), the patient usually feels sleepy during the first hour, is well during the next 2 hours and is in pain during the last hour because the blood level goes up quickly after the injection and goes down below the desirable level before the 4 hour interval prescribed for re-injection is reached. Transdermal administration as described herein permits the maintenance of constant levels for long periods of time (*e.g.*, days), which allows adequate pain control and mental alertness at the same time. Insulin provides another such example. Many diabetic patients need to maintain a constant baseline level of insulin which is different from their needs at the time of meals. The baseline level may be maintained using transdermal administration of insulin, as described herein. Antibiotics may also be administered at a constant rate, maintaining adequate bactericidal blood levels, while avoiding the high levels that are often responsible for

the toxicity (*e.g.*, levels of gentamycin that are too high typically result in renal toxicity).

Drug delivery by the methods of the present invention also provides a more convenient method of drug administration. For example, it is often particularly difficult to administer parenteral drugs to newborns and infants because of the difficulty associated with finding veins of acceptable caliber to catheterize. However, newborns and infants often have a relatively large skin surface as compared to adults. Transdermal drug delivery permits easier management of such patients and allows certain types of care that can presently be given only in hospitals to be given at home. Other patients who typically have similar difficulties with venous catheterization are patients undergoing chemotherapy or patients on dialysis. In addition, for patients undergoing prolonged therapy, transdermal administration as described herein is more convenient than parenteral administration.

Transdermal administration as described herein also allows the gastrointestinal tract to be bypassed in situations where parenteral uses would not be practical. For example, there is a growing need for methods suitable for administration of therapeutic small peptides and proteins, which are typically digested within the gastrointestinal tract. The methods described herein permit administration of such compounds and allow easy administration over long periods of time. Patients who have problems with absorption through their gastrointestinal tract because of prolonged ileus or specific gastrointestinal diseases limiting drug absorption may also benefit from drugs formulated for transdermal application as described herein.

Further, there are many clinical situations where it is difficult to maintain compliance. For example, patients with mental problems (*e.g.*, patients with Alzheimer's disease or psychosis) are easier to manage if a constant delivery rate of drug is provided without having to rely on their ability to take their medication at specific times of the day. Also patients who simply forget to take their drugs as prescribed are less likely to do so if they merely have to put on a skin patch periodically (*e.g.*, every 3 days). Patients with diseases that are without symptoms, like patients

with hypertension, are especially at risk of forgetting to take their medication as prescribed.

For patients taking multiple drugs, devices for transdermal application such as skin patches may be formulated with combinations of drugs that are frequently  
5 used together. For example, many heart failure patients are given digoxin in combination with furosemide. The combination of both drugs into a single skin patch facilitates administration, reduces the risk of errors (taking the correct pills at the appropriate time is often confusing to older people), reduces the psychological strain of taking "so many pills," reduces skipped dosage because of irregular activities and  
10 improves compliance.

The methods described herein are particularly applicable to humans, but also have a variety of veterinary uses, such as the administration of growth factors or hormones (*e.g.*, for fertility control) to an animal.

As noted above, a wide variety of drugs may be administered according  
15 to the methods provided herein. Some examples of drug categories that may be administered transdermally include anti-inflammatory drugs (*e.g.*, in arthritis and in other condition) such as all NSAID, indomethacin, prednisone, etc.; analgesics (especially when oral absorption is not possible, such as after surgery, and when parenteral administration is not convenient or desirable), including morphine, codeine,  
20 Demerol, acetaminophen and combinations of these (*e.g.*, codeine plus acetaminophen); antibiotics such as Vancomycin (which is not absorbed by the GI tract and is frequently given intravenously) or a combination of INH and Rifampicin (*e.g.*, for tuberculosis); anticoagulants such as heparin (which is not well absorbed by the GI tract and is generally given parenterally, resulting in fluctuation in the blood levels with an  
25 increased risk of bleeding at high levels and risks of inefficacy at lower levels) and Warfarin (which is absorbed by the GI tract but cannot be administered immediately after abdominal surgery because of the normal ileus following the procedure); antidepressants (*e.g.*, in situations where compliance is an issue as in Alzheimer's disease or when maintaining stable blood levels results in a significant reduction of anti-  
30 cholinergic side effects and better tolerance by patients), such as amitriptylin,

---

imipramin, prozac, etc.; antihypertensive drugs (*e.g.*, to improve compliance and reduce side effects associated with fluctuating blood levels), such as diuretics and beta-blockers (which can be administered by the same patch; *e.g.*, furosemide and propranolol); antipsychotics (*e.g.*, to facilitate compliance and make it easier for care giver and family members to make sure that the drug is received), such as haloperidol and chlorpromazine; and anxiolytics or sedatives (*e.g.*, to avoid the reduction of alertness related to high blood levels after oral administration and allow a continual benefit throughout the day by maintaining therapeutic levels constant).

Numerous other drugs may be administered as described herein, including naturally occurring and synthetic hormones, growth factors, proteins and peptides. For example, insulin and human growth hormone, growth factors like erythropoietin, interleukins and inteferons may be delivered via the skin.

Kits for administering a drug via the skin of a mammal are also provided within the present invention. Such kits generally comprise a device for transdermal application (*e.g.*, a skin patch) in combination with, or impregnated with, one or more modulating agents. A drug may additionally be included within such kits.

Within a related aspect, modulating agents as described herein may be used to increase the permeability of endothelial and epithelial cell layers, thereby facilitating sampling of the blood compartment by passive diffusion. Such methods permit the detection and/or measurement of the levels of specific molecules circulating in the blood. In general, to sample the blood compartment, it is necessary to perturb adhesion between the epithelial cells (keratinocytes) and the endothelial cells of the microvasculature. Using currently available techniques, only small, uncharged molecules may be detected across skin *in vivo*. The methods described herein are not subject to the same degree of limitation. Accordingly, a wide variety of blood components may be sampled across epithelial and endothelial cell layers. Such sampling may be achieved across any such cell layers, including skin and gums.

For example, application of one or more modulating agents to the skin, via a skin patch as described herein, permits the patch to function like a sponge to accumulate a small quantity of fluid containing a representative sample of the serum.

The patch is then removed after a specified amount of time and analyzed by suitable techniques for the compound of interest (*e.g.*, a medication, hormone, growth factor, metabolite or marker). Alternatively, a patch may be impregnated with reagents to permit a color change if a specific substance (*e.g.*, an enzyme) is detected. Substances  
5 that can be detected in this manner include, but are not limited to, illegal drugs such as cocaine, HIV enzymes, glucose and PSA. This technology is of particular benefit for home testing kits.

To facilitate sampling of blood in a patient, a modulating agent as described above for enhancing drug delivery is contacted with the skin surface.  
10 Modulating agent(s) and reagents for assaying blood components may, but need not, be contained within the same composition or skin patch. In general, the amount of modulating agent administered via the skin may vary as described above. Such levels may be achieved by appropriate adjustments to the device used, or by applying a cream formulated as described above. Transfer of the blood component across the skin may be  
15 predicted based on *in vitro* studies using, for example, a Franz cell apparatus, and evaluated *in vivo* by appropriate means that will be apparent to those of ordinary skill in the art.

Kits for sampling blood component via, for example, the skin or gums of a mammal, are also provided within the present invention. Such kits generally comprise  
20 a device for transdermal application (*i.e.*, skin patch) in combination with, or impregnated with, one or more modulating agents. A reagent for detection of a blood component may additionally be included within such kits.

Within a further aspect, methods are provided for enhancing delivery of a drug to a tumor in a mammal, comprising administering a modulating agent in  
25 combination with a drug to a tumor-bearing mammal. Such a modulating agent may comprise only a desmosomal cadherin CAR sequence, or may further comprise one or more other nonclassical or classical cadherin CAR sequences, such as a CAR sequence derived from OB-cadherin, cadherin-5, cadherin-6, E-cadherin and/or N-cadherin (*e.g.*, HAV, SHA VSS (SEQ ID NO: \_\_), AHAVDI (SEQ ID NO: \_\_) or a analogue of such a  
30 sequence). Bi-functional modulating agents that comprise the desmosomal cadherin

CAR sequence with either flanking E-cadherin-specific sequences or flanking N-cadherin-specific sequences joined via a linker to the desmosomal cadherin CAR sequence are also preferred. Preferably, the peptide portion(s) of a modulating agent comprises 6-16 amino acids, since longer peptides are difficult to dissolve in aqueous solution and are more likely to be degraded by peptidases.

In one particularly preferred embodiment, a modulating agent is capable of disrupting cell adhesion mediated by multiple adhesion molecules. For example, a single branched modulating agent (or multiple agents linked to a single molecule or support material) may disrupt adhesion mediated by a desmosomal cadherin, as well as E-cadherin, N-cadherin, occludin, claudin and integrin mediated cell adhesion. Such agents serve as multifunctional disrupters of cell adhesion. Alternatively, a separate modulator may be administered in conjunction with the modulating agent(s), either within the same pharmaceutical composition or separately. Preferred antibody modulating agents include Fab fragments directed against a nonclassical or classical cadherin CAR sequence, as described above. A Fab fragment may be incorporated into a modulating agent or may be present within a separate modulator that is administered concurrently.

Preferably, the modulating agent and the drug are formulated within the same composition or drug delivery device prior to administration. In general, a modulating agent may enhance drug delivery to any tumor (*e.g.*, breast tumor, stomach tumor, ovarian tumor or kidney tumor), and the method of administration may be chosen based on the type of target tumor. For example, injection or topical administration as described above may be preferred for melanomas and other accessible tumors (*e.g.*, metastases from primary ovarian tumors may be treated by flushing the peritoneal cavity with the composition). Other tumors (*e.g.*, breast tumors) may be treated by injection of the modulating agent and the drug (such as mitomycin C) into the site of the tumor. In other instances, the composition may be administered systemically, and targeted to the tumor using any of a variety of specific targeting agents. Suitable drugs may be identified by those of ordinary skill in the art based upon the type of cancer to be treated (*e.g.*, taxol for breast cancer). In general, the amount of

modulating agent administered varies with the method of administration and the nature of the tumor, within the typical ranges provided above, preferably ranging from about 1 $\mu$ g/mL to about 2 mg/mL, and more preferably from about 10 $\mu$ g/mL to 1mg/mL. Transfer of the drug to the target tumor may be evaluated by appropriate means that will  
5 be apparent to those of ordinary skill in the art. Drugs may also be labeled (*e.g.*, using radionuclides) to permit direct observation of transfer to the target tumor using standard imaging techniques.

Within a related aspect, the present invention provides methods for treating cancer in a mammal. Such a modulating agent may comprise only a  
10 desmosomal cadherin CAR sequence, or may further comprise one or more other nonclassical or classical cadherin CAR sequences, such as CAR sequences derived from OB-cadherin, cadherin-5, cadherin-6, E-cadherin and/or N-cadherin. CAR sequences involved in integrin-mediated cell adhesion (*e.g.*, RGD) may also be used. For example, such a modulating agent may further comprise a sequence such as HAV,  
15 SHAVSS (SEQ ID NO:\_\_), AHAVDI (SEQ ID NO:\_\_), RGD, YIGSR (SEQ ID NO:\_\_), or a derivative of such a sequence. Preferably, the peptide portion(s) of such modulating agents comprise 6-16 amino acids. Preferred antibody modulating agents include Fab fragments directed against a nonclassical or classical cadherin CAR sequence. The Fab fragments may be either incorporated into a modulating agent or  
20 may be present within a separate modulator that is administered concurrently.

A modulating agent may be administered alone (*e.g.*, via the skin) or within a pharmaceutical composition. For melanomas and certain other accessible tumors, injection or topical administration as described above may be preferred. Other tumors (*e.g.*, bladder tumors, bronchial tumors or tracheal tumors) may be treated by  
25 injection of the modulating agent into the cavity. In other instances, the composition may be administered systemically, and targeted to the tumor using any of a variety of specific targeting agents, as described above. Preferably, the tumor is a breast tumor, stomach tumor or kidney tumor. In general, the amount of modulating agent administered varies depending upon the method of administration and the nature of the  
30 cancer, but may vary within the ranges identified above. The effectiveness of the cancer

treatment may be evaluated using well known clinical observations, such as monitoring the level of serum tumor markers (*e.g.*, CEA or PSA).

The addition of a targeting agent as described above may be beneficial, particularly when the administration is systemic. Suitable modes of administration and dosages depend upon the condition to be prevented or treated but, in general, administration by injection is appropriate. Dosages may vary as described above. The effectiveness of the inhibition may be evaluated grossly by assessing the inability of the tumors to maintain their growth and microscopically by observing an absence of nerves at the periphery of the tumor.

In yet another related aspect, the present invention provides methods for inducing apoptosis in a desmosomal cadherin-expressing cell. Such a modulating agent may comprise only a desmosomal cadherin CAR sequence, or may further comprise one or more other nonclassical or classical cadherin CAR sequences, such as HAV, SHAVSS (SEQ ID NO:\_\_), AHAVDI (SEQ ID NO:\_\_), RGD, YIGSR (SEQ ID NO:\_\_) or an analogue of such a sequence. Preferably, the peptide portion(s) of such modulating agents comprise 6-16 amino acids. Preferred antibody modulating agents include Fab fragments directed against a nonclassical or classical cadherin CAR sequence. The Fab fragments may be either incorporated into a modulating agent or within a separate modulator that is administered concurrently. Administration may be topical, via injection or by other means, and the addition of a targeting agent may be beneficial, particularly when the administration is systemic. Suitable modes of administration and dosages depend upon the location and nature of the cells for which induction of apoptosis is desired but, in general, dosages may vary as described above. A biopsy may be performed to evaluate the level of induction of apoptosis.

In certain other aspects, the present invention provides methods for enhancing adhesion of desmosomal cadherin-expressing cells. Within certain embodiments, a modulating agent may be linked to a solid support, resulting in a matrix that comprises multiple modulating agents. Within one such embodiment, the support is a polymeric matrix to which modulating agents and molecules comprising other CAR sequence(s) are attached (*e.g.*, modulating agents and molecules comprising either HAV



or RGD sequences may be attached to the same matrix, preferably in an alternating pattern). Such matrices may be used in contexts in which it is desirable to enhance adhesion mediated by multiple cell adhesion molecules. Alternatively, the modulating agent itself may comprise multiple desmosomal cadherin CAR sequences or antibodies  
5 (or fragments thereof), separated by linkers as described above. Either way, the modulating agent(s) function as a "biological glue" to bind multiple desmosomal cadherin-expressing cells within a variety of contexts.

Within one such aspect, modulating agents comprising the desmosomal cadherin CAR sequence and/or multiple modulating agents linked to a single molecule  
10 or support material may be used to facilitate wound healing and/or reduce scar tissue in a mammal. Such a modulating agent may comprise one desmosomal cadherin CAR sequence, or may comprise multiple such sequences (*e.g.*, a desmoglein CAR sequence and a desmocollin CAR sequence). Optionally, a modulating agent may further comprise one or more other nonclassical cadherin CAR sequences, such as CAR  
15 sequences derived from OB-cadherin or cadherin-5 and/or one or more classical cadherin CAR sequences, including HAV, SHAVSS (SEQ ID NO: \_\_), AHAVDI (SEQ ID NO: \_\_), or an analogue of such a sequence. Preferred antibody modulating agents include Fab fragments directed against either the nonclassical cadherin or E-cadherin CAR sequences. Modulating agents that are linked to a biocompatible and  
20 biodegradable matrix such as cellulose or collagen are particularly preferred. For use within such methods, a modulating agent should have a free amino or hydroxyl group. The modulating agents are generally administered topically to the wound, where they may facilitate closure of the wound and may augment, or even replace, stitches. Similarly, administration of matrix-linked modulating agents may facilitate cell  
25 adhesion in skin grafting and prosthetic implants, and may prolong the duration and usefulness of collagen injection. In general, the amount of matrix-linked modulating agent administered to a wound, graft or implant site varies with the severity of the wound and/or the nature of the wound, graft, or implant, but may vary as discussed above. Multi-functional modulating agents comprising a desmosomal cadherin  
30 sequence, a classical cadherin CAR sequence (HAV), and the CAR sequence bound by

certain integrins (RGD) may also be used as potent stimulators of wound healing and/or to reduce scar tissue. Alternatively, one or more separate modulators of classical cadherin- or integrin-mediated cell adhesion may be administered in conjunction with the modulating agent(s), either within the same pharmaceutical composition or  
5 separately.

Within further aspects, modulating agents may be used for the treatment of autoimmune blistering disorders, such as pemphigus vulgaris, pemphigus foliaceus and intercellular IgA dermatosis. These disorders are pathological conditions in which a patient begins to develop antibodies against one or more desmosomal cadherins. In  
10 consequence, adhesion of skin cells begins to fail, resulting in various kinds of blistering. The use of modulating agents to enhance desmosomal cadherin-mediated cell adhesion may alleviate symptoms of such disorders. Optionally, such methods would be performed in conjunction with a method for eliminating the anti-cadherin antibodies. Modulating agents for use in such therapies may comprise one desmosomal  
15 cadherin CAR sequence or, preferably, comprise a desmoglein CAR sequence and a desmocollin CAR sequence. Optionally, a modulating agent further comprises one or more other nonclassical cadherin CAR sequences, such as CAR sequences derived from OB-cadherin or cadherin-5 and/or one or more classical cadherin CAR sequences, including HAV, SHAVSS (SEQ ID NO:\_\_), AHAVDI (SEQ ID NO:\_\_), or an  
20 analogue of such a sequence. Preferred antibody modulating agents include Fab fragments directed against either the nonclassical cadherin or E-cadherin CAR sequences. Modulating agents that are linked to a biocompatible and biodegradable matrix such as cellulose or collagen are particularly preferred. For use within such methods, a modulating agent should have a free amino or hydroxyl group. The  
25 modulating agents are generally administered topically to the site of blisters. In general, the amount of matrix-linked modulating agent administered to a blister site varies with the severity of the condition, and generally ranges as discussed above. Multi-functional modulating agents comprising one or more desmosomal cadherin sequences, a classical cadherin CAR sequence (HAV), and the CAR sequence bound by certain integrins  
30 (RGD) may also be used. Alternatively, one or more separate modulators of classical

cadherin- or integrin-mediated cell adhesion may be administered in conjunction with the modulating agent(s), either within the same pharmaceutical composition or separately.

Within another aspect, one or more modulating agents may be linked to  
5 the interior surface of a tissue culture plate or other cell culture support, such as for use in a bioreactor. Such linkage may be performed by any suitable technique, as described above. Modulating agents linked in this fashion may generally be used to immobilize cadherin-expressing cells. For example, dishes or plates coated with one or more modulating agents may be used to immobilize cadherin-expressing cells within a variety  
10 of assays and screens. Within bioreactors (*i.e.*, systems for large scale production of cells or organoids), modulating agents may generally be used to improve cell attachment and stabilize cell growth. Modulating agents may also be used within bioreactors to support the formation and function of highly differentiated organoids derived, for example, from dispersed populations of fetal mammalian cells. Bioreactors  
15 containing biomatrices of modulating agent(s) may also be used to facilitate the production of specific proteins.

Modulating agents as described herein may be used within a variety of bioreactor configurations. In general, a bioreactor is designed with an interior surface area sufficient to support large numbers of adherent cells. This surface area can be  
20 provided using membranes, tubes, microtiter wells, columns, hollow fibers, roller bottles, plates, dishes, beads or a combination thereof. A bioreactor may be compartmentalized. The support material within a bioreactor may be any suitable material known in the art; preferably, the support material does not dissolve or swell in water. Preferred support materials include, but are not limited to, synthetic polymers  
25 such as acrylics, vinyls, polyethylene, polypropylene, polytetrafluoroethylene, nylons, polyurethanes, polyamides, polysulfones and poly(ethylene terephthalate); ceramics; glass and silica.

Other aspects of the present invention provide methods that employ antibodies raised against a desmosomal CAR sequence for diagnostic and assay  
30 purposes. Assays typically involve using an antibody to detect the presence or absence

of a desmosomal cadherin (free or on the surface of a cell), or proteolytic fragments containing one or more EC domains in a suitable biological sample, such as tumor or normal tissue biopsies, blood, lymph node, serum or urine samples, or other tissue, homogenate, or extract thereof obtained from a patient.

5                   There are a variety of assay formats known to those of ordinary skill in the art for using an antibody to detect a target molecule in a sample. *See, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, the assay may be performed in a Western blot format, wherein a protein preparation from the biological sample is submitted to gel electrophoresis, transferred to  
10 a suitable membrane and allowed to react with the antibody. The presence of the antibody on the membrane may then be detected using a suitable detection reagent, as described below.

                  In another embodiment, the assay involves the use of antibody immobilized on a solid support to bind to the target cadherin, or a proteolytic fragment  
15 containing an extracellular domain and encompassing a CAR sequence, and remove it from the remainder of the sample. The bound cadherin may then be detected using a second antibody or reagent that contains a reporter group. Alternatively, a competitive assay may be utilized, in which a cadherin is labeled with a reporter group and allowed to bind to the immobilized antibody after incubation of the antibody with the sample.  
20 The extent to which components of the sample inhibit the binding of the labeled cadherin to the antibody is indicative of the reactivity of the sample with the immobilized antibody, and as a result, indicative of the level of the cadherin in the sample.

                  The solid support may be any material known to those of ordinary skill  
25 in the art to which the antibody may be attached, such as a test well in a microtiter plate, a nitrocellulose filter or another suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic such as polystyrene or polyvinylchloride. The antibody may be immobilized on the solid support using a variety of techniques known to those in the art, which are amply described in the patent  
30 and scientific literature.

In certain embodiments, the assay for detection of a desmosomal cadherin in a sample is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the biological sample, such that the desmosomal cadherin within the sample is allowed to bind to the immobilized antibody (a 30 minute incubation time at room temperature is generally sufficient). Unbound sample is then removed from the immobilized cadherin-antibody complexes and a second antibody (containing a reporter group such as an enzyme, dye, radionuclide, luminescent group, fluorescent group or biotin) capable of binding to a different site on the cadherin is added. The amount of second antibody that remains bound to the solid support is then determined using a method appropriate for the specific reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products. Standards and standard additions may be used to determine the level of cadherin in a sample, using well known techniques.

Such immunoassays may be used, for example, in the diagnosis and monitoring of conditions associated with abnormal desmosomal cadherin expression. As noted above, such conditions include autoimmune blistering disorders, such as pemphigus vulgaris, pemphigus foliaceus and intercellular IgA dermatosis. In general, the presence or absence of such a condition in a patient may be determined by (a) contacting a biological sample obtained from a patient with an antibody or antigen-binding fragment thereof; (b) detecting in the sample a level or pattern of polypeptide that binds to the antibody or fragment thereof; and (c) comparing the level or pattern of polypeptide with a predetermined cut-off value or a normal pattern. In general, these diseases are associated with an autoimmune response against desmosomal cadherins,

which causes the internalization of these cadherins. Such internalization may be visualized as patchy cytoplasmic granules in stained biopsy sections of lesional and perilesional keratinocytes (see Iwatsuki et al., *Br. J. Dermatol.* 140:35-43). These granules are not detected in normal patients.

5                   Immunoassays may further be used to differentiate between benign skin lesions, such as keratoacanthoma, and squamous cell carcinoma. The location of desmoglein staining, in particular, may be used to differentiate between a keratoacanthoma and a carcinoma (see Krunic et al., *Acta Derm. Venereol.* 76:394-398, 1996). In general, extensive pericellular staining indicates a keratoacanthoma. Focal  
10 pericellular staining indicates a keratoacanthoma or a well-differentiated squamous cell carcinoma. No staining or juxtanuclear staining indicates a poorly differentiated squamous cell carcinoma. Thus, a reduction of pericellular staining (as determined visually or quantitatively, as a reduction of 50%) is indicative of squamous cell carcinoma.

15                   The present invention also provides kits for use in such immunoassays. Such kits generally comprise one or more antibodies, as described above. In addition, one or more additional compartments or containers of a kit generally enclose elements, such as reagents, buffers and/or wash solutions, to be used in the immunoassay.

                  Within further aspects, modulating agents or antibodies (or fragments  
20 thereof) may be used to facilitate cell identification and sorting *in vitro* or imaging *in vivo*, permitting the selection of cells expressing the desmosomal cadherin (or different desmosomal cadherin levels). Preferably, the modulating agent(s) or antibodies for use in such methods are linked to a detectable marker. Suitable markers are well known in the art and include radionuclides, luminescent groups, fluorescent groups, enzymes,  
25 dyes, constant immunoglobulin domains and biotin. Within one preferred embodiment, a modulating agent linked to a fluorescent marker, such as fluorescein, is contacted with the cells, which are then analyzed by fluorescence activated cell sorting (FACS).

                  Antibodies or fragments thereof may also be used within screens of combinatorial or other nonpeptide-based libraries to identify other compounds capable  
30 of modulating desmosomal cadherin-mediated cell adhesion. Such screens may

generally be performed using an ELISA or other method well known to those of ordinary skill in the art that detect compounds with a shape and structure similar to that of the modulating agent. In general, such screens may involve contacting an expression library producing test compounds with an antibody, and detecting the level of antibody  
5 bound to the candidate compounds. Compounds for which the antibody has a higher affinity may be further characterized as described herein, to evaluate the ability to modulate OB-cadherin-mediated cell adhesion.

The following Example is offered by way of illustration and not by way of limitation.

## EXAMPLE

Preparation of Representative Modulating Agents

5                   This Example illustrates the solid phase synthesis of representative peptide modulating agents.

                  The peptides were synthesized on a 431A Applied Biosystems peptide synthesizer using p-Hydroxymethylphenoxymethyl polystyrene (HMP) resin and standard Fmoc chemistry. After synthesis and deprotection, the peptides were de-salted  
10                   on a Sephadex G-10 column and lyophilized. The peptides were analyzed for purity by analytical HPLC, and in each case a single peak was observed. Peptides were made as stock solutions at 10 to 25 mg/mL in dimethylsulfoxide (DMSO) or water and stored at -20°C before use.

15                   From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

---



What is claimed is:

1. A modulating agent that:

- (a) comprises a desmosomal cadherin CAR sequence; and
- (b) contains 3-16 amino acid residues linked by peptide bonds.

2. A modulating agent that:

- (a) comprises at least five consecutive amino acid residues of a desmosomal cadherin CAR sequence having the formula:

Aaa-Phe-Baa-Ile/Leu/Val-Asp/Asn/Glu-Caa-Daa-Ser/Thr/Asn-Gly

(SEQ ID NO:3)

wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine and asparagine; and

- (b) contains no more than 50 consecutive amino acid residues present within the desmosomal cadherin.

3. A modulating agent that:

- (a) comprises a desmosomal cadherin CAR sequence having the formula:

Aaa-Phe-Baa-Ile/Leu/Val-Asp/Asn/Glu-Caa-Daa-Ser/Thr/Asn-Gly

(SEQ ID NO:3)

wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine or asparagine; and

---

(b) contains no more than 50 consecutive amino acid residues present within the desmosomal cadherin.

4. A modulating agent that:

(a) comprises at least nine consecutive amino acid residues of a desmosomal cadherin, wherein the nine consecutive amino acids comprise a desmosomal cadherin CAR sequence having the formula:

Aaa-Phe-Baa-Ile/Leu/Val-Asp/Asn/Glu-Caa-Daa-Ser/Thr/Asn-Gly

(SEQ ID NO:3)

wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine and asparagine; and

(b) contains no more than 50 consecutive amino acid residues present within the desmosomal cadherin.


5. A modulating agent that comprises a desmosomal cadherin CAR sequence selected from the group consisting of NQK, NRN, NKD, EKD, ERD, RAL, YAL, YAT, FAT and YAS, wherein the agent contains no more than 50 consecutive amino acid residues present within a desmosomal cadherin.

6. A modulating agent according to any one of claims 2-5, wherein the agent is a peptide ranging in size from 3 to 50 amino acid residues.

7. A modulating agent according to any one of claims 1-5, wherein the agent is a peptide ranging in size from 4 to 16 amino acid residues.

8. A modulating agent according to any one of claims 1-5, wherein the CAR sequence is present within a cyclic peptide.

9. A modulating agent according to claim 8, wherein the cyclic peptide has the formula:



wherein W is a tripeptide selected from the group consisting of NQK, NRN, NKD, EKD, ERD, RAL, YAL, YAT, FAT and YAS;

wherein  $X_1$ , and  $X_2$  are optional, and if present, are independently selected from the group consisting of amino acid residues and combinations thereof in which the residues are linked by peptide bonds, and wherein  $X_1$  and  $X_2$  independently range in size from 0 to 10 residues, such that the sum of residues contained within  $X_1$  and  $X_2$  ranges from 1 to 12;

wherein  $Y_1$  and  $Y_2$  are independently selected from the group consisting of amino acid residues, and wherein a covalent bond is formed between residues  $Y_1$  and  $Y_2$ ; and


wherein  $Z_1$  and  $Z_2$  are optional, and if present, are independently selected from the group consisting of amino acid residues and combinations thereof in which the residues are linked by peptide bonds.

10. A polynucleotide encoding a modulating agent according to any one of claims 1-5.

11. An expression vector comprising a polynucleotide according to claim 10.

12. A host cell transformed or transfected with an expression vector according to claim 11.


13. A modulating agent comprising an antibody or antigen-binding fragment thereof that specifically binds to a desmosomal cadherin CAR sequence and modulates a desmosomal cadherin-mediated function, wherein the desmosomal cadherin CAR sequence has the formula:



wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine and asparagine; and

wherein the modulating agent inhibits or enhances a function mediated by the desmosomal cadherin.

14. A modulating agent comprising a mimetic of a desmosomal cadherin CAR sequence that comprises at least three consecutive amino acid residues of a desmosomal cadherin CAR sequence having the formula



wherein Aaa, Baa, Caa and Daa are independently selected amino acid residues; Ile/Leu/Val is an amino acid that is selected from the group consisting of isoleucine, leucine and valine, Asp/Asn/Glu is an amino acid that is selected from the group consisting of aspartate, asparagine and glutamate; and Ser/Thr/Asn is an amino acid that is selected from the group consisting of serine, threonine and asparagine;

wherein the mimetic is capable of modulating a desmosomal cadherin-mediated function.

15. A modulating agent according to any one of claims 1-5, wherein the agent comprises one or more desmoglein CAR sequences selected from the group consisting NQK, NQKT (SEQ ID NO:63), NQKTG (SEQ ID NO:64), INQK (SEQ ID

NO:65), INQKT (SEQ ID NO:66), INQKTG (SEQ ID NO:67), VINQK (SEQ ID NO:68), VINQKT (SEQ ID NO:69), VINQKTG (SEQ ID NO:70), FVINQK (SEQ ID NO:71), FVINQKT (SEQ ID NO:72), FVINQKTG (SEQ ID NO:73), IFVINQK (SEQ ID NO:74), IFVINQKT (SEQ ID NO:75), IFVINQKTG (SEQ ID NO:76), NRN, NRNT (SEQ ID NO:77), NRNTG (SEQ ID NO:78), INRN (SEQ ID NO:79), INRNT (SEQ ID NO:80), INRNTG (SEQ ID NO:81), IINRN (SEQ ID NO:82), IINRNT (SEQ ID NO:83), IINRNTG (SEQ ID NO:84), FIINRN (SEQ ID NO:85), FIINRNT (SEQ ID NO:86), FIINRNTG (SEQ ID NO:87), MFIINRN (SEQ ID NO:88), MFIINRNT (SEQ ID NO:89), MFIINRNTG (SEQ ID NO:90), NKD, NKDT (SEQ ID NO:91), NKDTG (SEQ ID NO:92), LNKD (SEQ ID NO:93), LNKDT (SEQ ID NO:94), LNKDTG (SEQ ID NO:95), YLNKD (SEQ ID NO:96), YLNKDT (SEQ ID NO:97), YLNKDTG (SEQ ID NO:98), FYLNKD (SEQ ID NO:99), FYLNKDT (SEQ ID NO:100), FYLNKDTG (SEQ ID NO:101), VFYLNKD (SEQ ID NO:102), VFYLNKDT (SEQ ID NO:103), VFYLNKDTG (SEQ ID NO:104), RAL, RALN (SEQ ID NO:151), RALNS (SEQ ID NO:152), RALNSM (SEQ ID NO:153), RALNSL (SEQ ID NO:154), RALNSMG (SEQ ID NO:155), RALNSLG (SEQ ID NO:156), CRAL (SEQ ID NO:157), CRALN (SEQ ID NO:158), CRALNS (SEQ ID NO:159), CRALNSM (SEQ ID NO:160), CRALNSL (SEQ ID NO:161), CRALNSMG (SEQ ID NO:162), CRALNSLG (SEQ ID NO:163), YCRAL (SEQ ID NO:164), YCRALN (SEQ ID NO:165), YCRALNS (SEQ ID NO:166), YCRALNSM (SEQ ID NO:167), YCRALNSL (SEQ ID NO:168), YCRALNSMG (SEQ ID NO:169), YCRALNSLG (SEQ ID NO:170), IYRAL (SEQ ID NO:171), IYRALN (SEQ ID NO:172), IYRALNS (SEQ ID NO:173), IYRALNSM (SEQ ID NO:174), IYRALNSL (SEQ ID NO:175), IYRALNSMG (SEQ ID NO:176), IYRALNSLG (SEQ ID NO:177), YAL, YALD (SEQ ID NO:180), YALDA (SEQ ID NO:181), YALDAR (SEQ ID NO:182), YALDARG (SEQ ID NO:183), GYAL (SEQ ID NO:184), GYALD (SEQ ID NO:185), GYALDA (SEQ ID NO:186), GYALDAR (SEQ ID NO:187), GYALDARG (SEQ ID NO:188), TGYAL (SEQ ID NO:189), TGYALD (SEQ ID NO:190), TGYALDA (SEQ ID NO:191), TGYALDAR (SEQ ID NO:192), TGYALDARG (SEQ ID NO:193), LTGYAL (SEQ ID NO:194), LTGYALD (SEQ ID NO:195), LTGYALDA (SEQ ID NO:196), LTGYALDAR (SEQ ID NO:197),

LTGYALDARG (SEQ ID NO:198), RALNA (SEQ ID NO:200), RALNAQ (SEQ ID NO:201), RALNAL (SEQ ID NO:202), RALNAQG (SEQ ID NO:203), RALNALG (SEQ ID NO:204), CRALNA (SEQ ID NO:205), CRALNAQ (SEQ ID NO:206), CRALNAL (SEQ ID NO:207), CRALNAQG (SEQ ID NO:208), CRALNALG (SEQ ID NO:209), TCRAL (SEQ ID NO:210), TCRALN (SEQ ID NO:211), TCRALNA (SEQ ID NO:212), TCRALNAQ (SEQ ID NO:213), TCRALNAL (SEQ ID NO:214), TCRALNAQG (SEQ ID NO:215), TCRALNALG (SEQ ID NO:216), ITCRAL (SEQ ID NO:217), ITCRALN (SEQ ID NO:218), ITCRALNA (SEQ ID NO:219), ITCRALNAQ (SEQ ID NO:220), ITCRALNAL (SEQ ID NO:221), ITCRALNAQG (SEQ ID NO:222) and ITCRALNALG (SEQ ID NO:223).

16. A modulating agent according to claim 15, wherein the agent comprises a linear peptide having the sequence N-Ac-IFVINQKTG-NH<sub>2</sub> (SEQ ID NO:105), N-Ac-MFIINRNTG-NH<sub>2</sub> (SEQ ID NO:106), N-Ac-VFYLNKDTG-NH<sub>2</sub> (SEQ ID NO:107), N-Ac-IYRALNSMG-NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-IYRALNSLG-NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-LTGYALDARG-NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-ITCRALNAQG-NH<sub>2</sub> (SEQ ID NO:\_\_\_) and N-Ac-ITCRALNALG-NH<sub>2</sub> (SEQ ID NO:\_\_\_).

17. A modulating agent according to claim 15, wherein a desmoglein CAR sequence is present within a cyclic peptide.

18. A modulating agent according to claim 17, wherein the cyclic peptide comprises a sequence selected from the group consisting of CNQKC (SEQ ID NO:3066), CNQKTC (SEQ ID NO:3067), CNQKTGC (SEQ ID NO:3068), CINQKC (SEQ ID NO:3069), CINQKTC (SEQ ID NO:3070), CINQKTGC (SEQ ID NO:3071), CVINQKC (SEQ ID NO:3072), CVINQKTC (SEQ ID NO:3073), CVINQKTGC (SEQ ID NO:3074), CFVINQKC (SEQ ID NO:3075), CFVINQKTC (SEQ ID NO:3076), CFVINQKTGC (SEQ ID NO:3077), CIFVINQKC (SEQ ID NO:3078), CIFVINQKTC (SEQ ID NO:3079), CIFVINQKTGC (SEQ ID NO:3080), CNRNC (SEQ ID NO:3081), CNRNTC (SEQ ID NO:3082), CNRNTGC (SEQ ID NO:3083), CINRNC

(SEQ ID NO:3084), CINRNTC (SEQ ID NO:3085), CINRNTGC (SEQ ID NO:3086), CIINRNC (SEQ ID NO:3087), CIINRNTC (SEQ ID NO:3088), CIINRNTGC (SEQ ID NO:3089), CFIINRNC (SEQ ID NO:3090), CFIINRNTC (SEQ ID NO:3091), CFIINRNTGC (SEQ ID NO:3092), CMFIINRNC (SEQ ID NO:3093), CMFIINRNTC (SEQ ID NO:3094), CMFIINRNTGC (SEQ ID NO:3095), CNKDC (SEQ ID NO:3096), CNKDTC (SEQ ID NO:3097), CNKDTGC (SEQ ID NO:3098), CLNKDC (SEQ ID NO:3099), CLNKDTC (SEQ ID NO:3100), CLNKDTGC (SEQ ID NO:3101), CYLNKDC (SEQ ID NO:3102), CYLNKDTGC (SEQ ID NO:3103), CYLNKDTGC (SEQ ID NO:3104), CFYLNKDC (SEQ ID NO:3105), CFYLNKDTGC (SEQ ID NO:3106), CFYLNKDTGC (SEQ ID NO:3107), CVFYLNKDC (SEQ ID NO:3108), CVFYLNKDTGC (SEQ ID NO:3109), CVFYLNKDTGC (SEQ ID NO:3110), ENQKK (SEQ ID NO:3111), ENQKTK (SEQ ID NO:3112), ENQKTGK (SEQ ID NO:3113), EINQKK (SEQ ID NO:3114), EINQKTK (SEQ ID NO:3115), EINQKTGK (SEQ ID NO:3116), EVINQKK (SEQ ID NO:3117), EVINQKTK (SEQ ID NO:3118), EVINQKTGK (SEQ ID NO:3119), EFVINQKK (SEQ ID NO:3120), EFVINQKTK (SEQ ID NO:3121), EFVINQKTGK (SEQ ID NO:3122), EIFVINQKK (SEQ ID NO:3123), EIFVINQKTK (SEQ ID NO:3124), EIFVINQKTGK (SEQ ID NO:3125), ENRNK (SEQ ID NO:3126), ENRNTK (SEQ ID NO:3127), ENRNTGK (SEQ ID NO:3128), EINRNK (SEQ ID NO:3129), EINRNTK (SEQ ID NO:3130), EINRNTGK (SEQ ID NO:3131), EIINRNK (SEQ ID NO:3132), EIINRNTK (SEQ ID NO:3133), EIINRNTGK (SEQ ID NO:3134), EFIINRNK (SEQ ID NO:3135), EFIINRNTK (SEQ ID NO:3136), EFIINRNTGK (SEQ ID NO:3137), EMFIINRNK (SEQ ID NO:3138), EMFIINRNTK (SEQ ID NO:3139), EMFIINRNTGK (SEQ ID NO:3140), ENKDK (SEQ ID NO:3141), ENKDTK (SEQ ID NO:3142), ENKDTGK (SEQ ID NO:3143), ELNKDK (SEQ ID NO:3144), ELNKDTK (SEQ ID NO:3145), ELNKDTGK (SEQ ID NO:3146), EYLNKDK (SEQ ID NO:3147), EYLNKDTK (SEQ ID NO:3148), EYLNKDTGK (SEQ ID NO:3149), EFYLNKDK (SEQ ID NO:3150), EFYLNKDTK (SEQ ID NO:3151), EFYLNKDTGK (SEQ ID NO:3152), EVFYLNKDK (SEQ ID NO:3153), EVFYLNKDTK (SEQ ID NO:3154), EVFYLNKDTGK (SEQ ID NO:3155), KNQKD (SEQ ID NO:3156), KNQKTD (SEQ

ID NO:3157), KNQKTGD (SEQ ID NO:3158), KINQKD (SEQ ID NO:3159),  
KINQKTD (SEQ ID NO:3160), KINQKTGD (SEQ ID NO:3161), KVINQKD (SEQ  
 ID NO:3162), KVINQKTD (SEQ ID NO:3163), KVINQKTGD (SEQ ID NO:3164),  
KFVINQKD (SEQ ID NO:3165), KFVINQKTD (SEQ ID NO:3166), KFVINQKTGD  
 (SEQ ID NO:3167), KIFVINQKD (SEQ ID NO:3168), KIFVINQKTD (SEQ ID  
 NO:3169), KIFVINQKTGD (SEQ ID NO:3170), KNRND (SEQ ID NO:3171),  
KNRNTD (SEQ ID NO:3172), KNRNTGD (SEQ ID NO:3173), KINRND (SEQ ID  
 NO:3174), KINRNTD (SEQ ID NO:3175), KINRNTGD (SEQ ID NO:3176),  
KIINRND (SEQ ID NO:3177), KIINRNTD (SEQ ID NO:3178), KIINRNTGD (SEQ  
 ID NO:3179), KFIINRND (SEQ ID NO:3180), KFIINRNTD (SEQ ID NO:3181),  
KFIINRNTGD (SEQ ID NO:3182), KMFIINRND (SEQ ID NO:3183), KMFIINRNTD  
 (SEQ ID NO:3184), KMFIINRNTGD (SEQ ID NO:3185), KNKDD (SEQ ID  
 NO:3186), KNKDTD (SEQ ID NO:3187), KNKDTGD (SEQ ID NO:3188), KLNKDD  
 (SEQ ID NO:3189), KLNKDTD (SEQ ID NO:3190) KLNKDTGD (SEQ ID NO:3191),  
KYLNKDD (SEQ ID NO:3192), KYLNKDTD (SEQ ID NO:3193), KYLNKDTGD  
 (SEQ ID NO:3194), KFYLNKDD (SEQ ID NO:3195), KFYLNKDTD (SEQ ID  
 NO:3196), KFYLNKDTGD (SEQ ID NO:3197), KVFYLNKDD (SEQ ID NO:3198),  
KVFYLNKDTD (SEQ ID NO:3199), KVFYLNKDTGD (SEQ ID NO:3200), DNQKK  
 (SEQ ID NO:3201), DNQKTK (SEQ ID NO:3202), DNQKTGK (SEQ ID NO:3203),  
DINQKK (SEQ ID NO:3204), DINQKTK (SEQ ID NO:3205), DINQKTGK (SEQ ID  
 NO:3206), DVINQKK (SEQ ID NO:3207), DVINQKTK (SEQ ID NO:3208),  
DVINQKTGK (SEQ ID NO:3209), DFVINQKK (SEQ ID NO:3210), DFVINQKTK  
 (SEQ ID NO:3211), DFVINQKTGK (SEQ ID NO:3212), DIFVINQKK (SEQ ID  
 NO:3213), DIFVINQKTK (SEQ ID NO:3214), DIFVINQKTGK (SEQ ID NO:3215),  
DNRNK (SEQ ID NO:3216), DNRNTK (SEQ ID NO:3217), DNRNTGK (SEQ ID  
 NO:3218), DINRNK (SEQ ID NO:3219), DINRNTK (SEQ ID NO:3220), DINRNTGK  
 (SEQ ID NO:3221), DIINRNK (SEQ ID NO:3222), DIINRNTK (SEQ ID NO:3223),  
DIINRNTGK (SEQ ID NO:3224), DFIINRNK (SEQ ID NO:3225), DFIINRNTK (SEQ  
 ID NO:3226), DFIINRNTGK (SEQ ID NO:3227), DMFIINRNK (SEQ ID NO:3228),  
DMFIINRNTK (SEQ ID NO:3229), DMFIINRNTGK (SEQ ID NO:3230), DNKDK



(SEQ ID NO:3231), DNKDTK (SEQ ID NO:3232), DNKDTGK (SEQ ID NO:3233), DLNKDK (SEQ ID NO:3234), DLNKDTK (SEQ ID NO:3235), DLNKDTGK (SEQ ID NO:3236), DYLNKDK (SEQ ID NO:3237), DYLNKDTK (SEQ ID NO:3238), DYLNKDTGK (SEQ ID NO:3239), DFYLNKDK (SEQ ID NO:3240), DFYLNKDTK (SEQ ID NO:3241), DFYLNKDTGK (SEQ ID NO:3242), DVFYLNKDK (SEQ ID NO:3243), DVFYLNKDTK (SEQ ID NO:3244), DVFYLNKDTGK (SEQ ID NO:3245), KKNQKE (SEQ ID NO:3246), KNQKTE (SEQ ID NO:3247), KNQKTGE (SEQ ID NO:3248), KINQKE (SEQ ID NO:3249), KINQKTE (SEQ ID NO:3250), KINQKTGE (SEQ ID NO:3251), KVINQKE (SEQ ID NO:3252), KVINQKTE (SEQ ID NO:3253), KVINQKTGE (SEQ ID NO:3254), KFVINQKE (SEQ ID NO:3255), KFVINQKTE (SEQ ID NO:3256), KFVINQKTGE (SEQ ID NO:3257), KIFVINQKE (SEQ ID NO:3258), KIFVINQKTE (SEQ ID NO:3259), KIFVINQKTGE (SEQ ID NO:3260), KNRNE (SEQ ID NO:3261), KNRNTE (SEQ ID NO:3262), KNRNTGE (SEQ ID NO:3263), KINRNE (SEQ ID NO:3264), KINRNTE (SEQ ID NO:3265), KINRNTGE (SEQ ID NO:3266), KIINRNE (SEQ ID NO:3267), KIINRNTE (SEQ ID NO:3268), KIINRNTGE (SEQ ID NO:3269), KFIINRNE (SEQ ID NO:3270), KFIINRNTE (SEQ ID NO:3271), KFIINRNTGE (SEQ ID NO:3272), KMFIINRNE (SEQ ID NO:3273), KMFIINRNTE (SEQ ID NO:3274), KMFIINRNTGE (SEQ ID NO:3275), KNKDE (SEQ ID NO:3276), KNKDE (SEQ ID NO:3277), KNKDTGE (SEQ ID NO:3278), KLNKDE (SEQ ID NO:3279), KLNKDE (SEQ ID NO:3280), KLNKDTGE (SEQ ID NO:3281), KYLNKDE (SEQ ID NO:3282), KYLNKDE (SEQ ID NO:3283), KYLNKDTGE (SEQ ID NO:3284), KFYLNKDE (SEQ ID NO:3285), KFYLNKDE (SEQ ID NO:3286), KFYLNKDTGE (SEQ ID NO:3287), KVFYLNKDE (SEQ ID NO:3288), KVFYLNKDE (SEQ ID NO:3289), KVFYLNKDTGE (SEQ ID NO:3290), NQKTG (SEQ ID NO:3291), INQKT (SEQ ID NO:3292), INQKTG (SEQ ID NO:3293), VINQK (SEQ ID NO:3294), VINQKT (SEQ ID NO:3295), VINQKTG (SEQ ID NO:3296), FVINQK (SEQ ID NO:3297), FVINQKT (SEQ ID NO:3298), FVINQKTG (SEQ ID NO:3299), IFVINQK (SEQ ID NO:3300), IFVINQKT (SEQ ID NO:3301), IFVINQKTG (SEQ ID NO:3302), NRNTG (SEQ ID NO:3303), INRNT (SEQ ID NO:3304), INRNTG (SEQ ID NO:3305), IINRN

(SEQ ID NO:3306), IINRNT (SEQ ID NO:3307), IINRNTG (SEQ ID NO:3308), FIINRN (SEQ ID NO:3309), FIINRNT (SEQ ID NO:3310), FIINRNTG (SEQ ID NO:3311), MFIINRN (SEQ ID NO:3312), MFIINRNT (SEQ ID NO:3313), MFIINRNTG (SEQ ID NO:3314), NKDTG (SEQ ID NO:3315), LNKDT (SEQ ID NO:3316), LNKDTG (SEQ ID NO:3317), YLNKD (SEQ ID NO:3318), YLNKDT (SEQ ID NO:3319), YLNKDTG (SEQ ID NO:3320), FYLNKD (SEQ ID NO:3321), FYLNKDT (SEQ ID NO:3322), FYLNKDTG (SEQ ID NO:3323), VFYLNKD (SEQ ID NO:3324), VFYLNKDT (SEQ ID NO:3325) and VFYLNKDTG (SEQ ID NO:3326), CRALC, CRALNC, CRALNSC, CRALNSMC, CRALNSLC, CRALNSMGC, CRALNSLGC, CCRALC, CCRALNC, CCRALNSC, CCRALNSMC, CCRALNSLC, CCRALNSMGC, CCRALNSLGC, CYCRALC, CYCRALNC, CYCRALNSC, CYCRALNSMC, CYCRALNSLC, CYCRALNSMGC, CYCRALNSLGC, CIYRALC, CIYRALNC, CIYRALNSC, CIYRALNSMC, CIYRALNSLC, CIYRALNSMGC, CIYRALNSLGC, ERALK, ERALNK, ERALNSK, ERALNSMK, ERALNSLK, ERALNSMGK, ERALNSLGK, ECRALK, ECRALNK, ECRALNSK, ECRALNSMK, ECRALNSLK, ECRALNSMGK, ECRALNSLGK, EYCRALK, EYCRALNK, EYCRALNSK, EYCRALNSMK, EYCRALNSLK, EYCRALNSMGK, EYCRALNSLGK, EIYRALK, EIYRALNK, EIYRALNSK, EIYRALNSMK, EIYRALNSLK, EIYRALNSMGK, EIYRALNSLGK, KRALE, KRALNE, KRALNSE, KRALNSME, KRALNSLE, KRALNSMGE, KRALNSLGE, KCRALE, KCRALNE, KCRALNSE, KCRALNSME, KCRALNSLE, KCRALNSMGE, KCRALNSLGE, KYCRALC, KYCRALNE, KYCRALNSE, KYCRALNSME, KYCRALNSLE, KYCRALNSMGE, KYCRALNSLGE, KIYRALC, KIYRALNE, KIYRALNSE, KIYRALNSME, KIYRALNSLE, KIYRALNSMGE, KIYRALNSLGE, DRALK, DRALNK, DRALNSK, DRALNSMK, DRALNSLK, DRALNSMGK, DRALNSLGK, DCRALK, DCRALNK, DCRALNSK, DCRALNSMK, DCRALNSLK, DCRALNSMGK, DCRALNSLGK, DYCRALK, DYCRALNK, DYCRALNSK, DYCRALNSMK, DYCRALNSLK, DYCRALNSMGK, DYCRALNSLGK, DIYRALC, DIYRALNK, DIYRALNSK, DIYRALNSMK, DIYRALNSLK, DIYRALNSMGK, DIYRALNSLGK, KRALD,

KRALND, KRALNSD, KRALNSMD, KRALNSLD, KRALNSMGD, KRALNSLGD,  
KCRALD, KCRALND, KCRALNSD, KCRALNSMD, KCRALNSLD,  
KCRALNSMGD, KCRALNSLGD, KYCRALED, KYCRALEDND, KYCRALEDNSD,  
KYCRALEDNSMD, KYCRALEDNSLD, KYCRALEDNSMGD, KYCRALEDNSLGD,  
KIYRALD, KIYRALND, KIYRALNSD, KIYRALNSMD, KIYRALNSLD,  
KIYRALNSMGD, KIYRALNSLGD, RALNS, RALNSM, RALNSL, RALNSMG,  
RALNSLG, CRALN, CRALNS, CRALNSM, CRALNSL, CRALNSMG,  
CRALNSLG, YCRAL, YCRALN, YCRALNS, YCRALNSM, YCRALNSL,  
YCRALNSMG, YCRALNSLG, IYRAL, IYRALN, IYRALNS, IYRALNSM,  
IYRALNSL, IYRALNSMG, IYRALNSLG, CYALC, CYALDC, CYALDAC,  
CYALDARC, CYALDARGC, CGYALC, CGYALDC, CGYALDAC, CGYALDARC,  
CGYALDARGC, CTGYALC, CTGYALDC, CTGYALDAC, CTGYALDARC,  
CTGYALDARGC, CLTGYALC, CLTGYALDC, CLTGYALDAC,  
CLTGYALDARC, CLTGYALDARGC, EYALK, EYALDK, EYALDAK,  
EYALDARK, EYALDARGK, EGYALK, EGYALDK, EGYALDAK, EGYALDARK,  
EGYALDARGK, ETGYALK, ETGYALDK, ETGYALDAK, ETGYALDARK,  
ETGYALDARGK, ELTGYALK, ELTGYALDK, ELTGYALDAK, ELTGYALDARK,  
ELTGYALDARGK, KYALE, KYALDE, KYALDAE, KYALDARE, KYALDARGE,  
KGYALE, KGYALDE, KGYALDAE, KGYALDARE, KGYALDARGE, KTGYALE,  
KTGYALDE, KTGYALDAE, KTGYALDARE, KTGYALDARGE, KLTGYALE,  
KLTGYALDE, KLTGYALDAE, KLTGYALDARE, KLTGYALDARGE, DYALK,  
DYALDK, DYALDAK, DYALDARK, DYALDARGK, DGYALK, DGYALDK,  
DGYALDAK, DGYALDARK, DGYALDARGK, DTGYALK, DTGYALDK,  
DTGYALDAK, DTGYALDARK, DTGYALDARGK, DLTGYALK, DLTGYALDK,  
DLTGYALDAK, DLTGYALDARK, DLTGYALDARGK, KYALD, KYALDD,  
KYALDAD, KYALDARD, KYALDARGD, KGYALD, KGYALDD, KGYALDAD,  
KGYALDARD, KGYALDARGD, KTGYALD, KTGYALDD, KTGYALDAD,  
KTGYALDARD, KTGYALDARGD, KLTGYALD, KLTGYALDD, KLTGYALDAD,  
KLTGYALDARD, KLTGYALDARGD, YALDA, YALDAR, YALDARG, GYALD,  
GYALDA, GYALDAR, GYALDARG, TGYAL, TGYALD, TGYALDA

TGYALDAR, TGYALDARG, LTGYAL, LTGYALD, LTGYALDA, LTGYALDAR,  
LTGYALDARG, CRALC, CRALNC, CRALNAC, CRALNAQC, CRALNALC,  
CRALNAQGC, CRALNALGC, CCRALC, CCRALNC, CCRALNAC,  
CCRALNAQC, CCRALNALC, CCRALNAQGC, CCRALNALGC, CTCRALC,  
CTCRALNC, CTCRALNAC, CTCRALNAQC, CTCRALNALC, CTCRALNAQGC,  
CTCRALNALGC, CITCRALC, CITCRALNC, CITCRALNAC, CITCRALNAQC,  
CITCRALNALC, CITCRALNAQGC, CITCRALNALGC, ERALK, ERALNK,  
ERALNAK, ERALNAQK, ERALNALK, ERALNAQK, ERALNALGK, ECRALK,  
ECRALNK, ECRALNAK, ECRALNAQK, ECRALNALK, ECRALNAQK,  
ECRALNALGK, ETCRALK, ETCRALNK, ETCRALNAK, ETCRALNAQK,  
ETCRALNALGK, ETCRALNAQK, ETCRALNALGK, EITCRALK, EITCRALNK,  
EITCRALNAK, EITCRALNAQK, EITCRALNALK, EITCRALNAQK,  
EITCRALNALGK, KRALE, KRALNE, KRALNAE, KRALNAQE, KRALNALE,  
KRALNAQGE, KRALNALGE, KCRALE, KCRALNE, KCRALNAE,  
KCRALNAQE, KCRALNALE, KCRALNAQGE, KCRALNALGE, KTCRALE,  
KTCRALNE, KTCRALNAE, KTCRALNAQE, KTCRALNALE, KTCRALNAQGE,  
KTCRALNALGE, KITCRALE, KITCRALNE, KITCRALNAE, KITCRALNAQE,  
KITCRALNALE, KITCRALNAQGE, KITCRALNALGE, DRALK, DRALNK,  
DRALNAK, DRALNAQK, DRALNALK, DRALNAQK, DRALNALGK,  
DCRALK, DCRALNK, DCRALNAK, DCRALNAQK, DCRALNALGK,  
DCRALNAQK, DCRALNALGK, DTCRALK, DTCRALNK, DTCRALNAK,  
DTCRALNAQK, DTCRALNALK, DTCRALNAQK, DTCRALNALGK,  
DITCRALK, DITCRALNK, DITCRALNAK, DITCRALNAQK, DITCRALNALGK,  
DITCRALNAQK, DITCRALNALGK, KRALD, KRALND, KRALNAD,  
KRALNAQD, KRALNALD, KRALNAQGD, KRALNALGD, KCRALD, KCRALND,  
KCRALNAD, KCRALNAQD, KCRALNALD, KCRALNAQGD, KCRALNALGD,  
KTCRALD, KTCRALND, KTCRALNAD, KTCRALNAQD, KTCRALNALD,  
KTCRALNAQGD, KTCRALNALGD, KITCRALD, KITCRALND, KITCRALNAD,  
KITCRALNAQD, KITCRALNALD, KITCRALNAQGD, KITCRALNALGD,  
RALNA, RALNAQ, RALNAL, RALNAQ, RALNALG, CRALN, CRALNA,

CRALNAQ, CRALNAL, CRALNAQG, CRALNALG, TCRAL, TCRALN, TCRALNA, TCRALNAQ, TCRALNAL, TCRALNAQG, TCRALNALG, ITCRAL, ITCRALN, ITCRALNA, ITCRALNAQ, ITCRALNAL, ITCRALNAQG and ITCRALNALG.

19. A modulating agent comprising an antibody or antigen-binding fragment thereof that:

(a) specifically binds to a desmoglein CAR sequence selected from the group consisting of N-Ac-IFVINQKTG-NH<sub>2</sub> (SEQ ID NO:105), N-Ac-MFIINRNTG-NH<sub>2</sub> (SEQ ID NO:106), N-Ac-VFYLNKDTG-NH<sub>2</sub> (SEQ ID NO:107), N-Ac-IYRALNSMG-NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-IYRALNSLG -NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-LTGYALDARG -NH<sub>2</sub> (SEQ ID NO:\_\_\_), N-Ac-ITCRALNAQG-NH<sub>2</sub> (SEQ ID NO:\_\_\_) and N-Ac-ITCRALNALG-NH<sub>2</sub> (SEQ ID NO:\_\_\_); and

(b) modulates a desmoglein-mediated function.

20. A modulating agent according to any one of claims 1-5, wherein the agent comprises one or more desmocollin CAR sequences selected from the group consisting of EKD, EKDT (SEQ ID NO:108), EKDTG (SEQ ID NO:109), IEKD (SEQ ID NO:110), IEKDT (SEQ ID NO:111), IEKDTG (SEQ ID NO:112), YIEKD (SEQ ID NO:113), YIEKDT (SEQ ID NO:114), YIEKDTG (SEQ ID NO:115), FYIEKD (SEQ ID NO:116), FYIEKDT (SEQ ID NO:117), FYIEKDTG (SEQ ID NO:118), LFYIEKD (SEQ ID NO:119), LFYIEKDT (SEQ ID NO:120), LFYIEKDTG (SEQ ID NO:121), ERD, ERDT (SEQ ID NO:122), ERDTG (SEQ ID NO:123), VERD (SEQ ID NO:124), VERDT (SEQ ID NO:125), VERDTG (SEQ ID NO:126), YVERD (SEQ ID NO:127), YVERDT (SEQ ID NO:128), YVERDTG (SEQ ID NO:129), FYVERD (SEQ ID NO:130), FYVERDT (SEQ ID NO:131), FYVERDTG (SEQ ID NO:132), LFYVERD (SEQ ID NO:133), LFYVERDT (SEQ ID NO:134), LFYVERDTG (SEQ ID NO:135), IERD (SEQ ID NO:136), IERDT (SEQ ID NO:137), IERDTG (SEQ ID NO:138), YIERD (SEQ ID NO:139), YIERDT (SEQ ID NO:140), YIERDTG (SEQ ID NO:141), FYIERD (SEQ ID NO:142), FYIERDT (SEQ ID NO:143), FYIERDTG (SEQ ID NO:144), LFYIERD (SEQ ID NO:145), LFYIERDT (SEQ ID NO:146) and

LFYIERDTG (SEQ ID NO:147), YAT, YATT (SEQ ID NO:226), YATTA (SEQ ID NO:227), YATTAD (SEQ ID NO:228), YATTADG (SEQ ID NO:229), GYAT (SEQ ID NO:230), GYATT (SEQ ID NO:231), GYATTA (SEQ ID NO:232), GYATTAD (SEQ ID NO:233), GYATTADG (SEQ ID NO:234), AYAT (SEQ ID NO:235), AYATT (SEQ ID NO:236), AYATTA (SEQ ID NO:237), AYATTAD (SEQ ID NO:238), AYATTADG (SEQ ID NO:239), YGYAT (SEQ ID NO:240), YGYATT (SEQ ID NO:241), YGYATTA (SEQ ID NO:242), YGYATTAD (SEQ ID NO:243), YGYATTADG (SEQ ID NO:244), YAYAT (SEQ ID NO:245), YAYATT (SEQ ID NO:246), YAYATTA (SEQ ID NO:247), YAYATTAD (SEQ ID NO:248), YAYATTADG (SEQ ID NO:249), LYGYAT (SEQ ID NO:250), LYGYATT (SEQ ID NO:251), LYGYATTA (SEQ ID NO:252), LYGYATTAD (SEQ ID NO:253), LYGYATTADG (SEQ ID NO:254), LYAYAT (SEQ ID NO:255), LYAYATT (SEQ ID NO:256), LYAYATTA (SEQ ID NO:257), LYAYATTAD (SEQ ID NO:258), LYAYATTADG (SEQ ID NO:259), VYGYAT (SEQ ID NO:260), VYGYATT (SEQ ID NO:261), VYGYATTA (SEQ ID NO:262), VYGYATTAD (SEQ ID NO:263), VYGYATTADG (SEQ ID NO:264), VYAYAT (SEQ ID NO:265), VYAYATT (SEQ ID NO:266), VYAYATTA (SEQ ID NO:267), VYAYATTAD (SEQ ID NO:268), VYAYATTADG (SEQ ID NO:269), IYGYAT (SEQ ID NO:270), IYGYATT (SEQ ID NO:271), IYGYATTA (SEQ ID NO:272), IYGYATTAD (SEQ ID NO:273), IYGYATTADG (SEQ ID NO:274), IYAYAT (SEQ ID NO:275), IYAYATT (SEQ ID NO:276), IYAYATTA (SEQ ID NO:277), IYAYATTAD (SEQ ID NO:278) or IYAYATTADG (SEQ ID NO:279), FAT, FATT (SEQ ID NO:286), FATTP (SEQ ID NO:287), FATTPD (SEQ ID NO:288), FATTPDG (SEQ ID NO:289), AFAT (SEQ ID NO:290), AFATT (SEQ ID NO:291), AFATTP (SEQ ID NO:292), AFATTPD (SEQ ID NO:293), AFATTPDG (SEQ ID NO:294), IAFAT (SEQ ID NO:295), IAFATT (SEQ ID NO:296), IAFATTP (SEQ ID NO:297), IAFATTPD (SEQ ID NO:298), IAFATTPDG (SEQ ID NO:299), IIAFAT (SEQ ID NO:300), IIAFATT (SEQ ID NO:301), IIAFATTP (SEQ ID NO:302), IIAFATTPD (SEQ ID NO:303), IIAFATTPDG (SEQ ID NO:304), LIAFAT (SEQ ID NO:305), LIAFATT (SEQ ID NO:306), LIAFATTP (SEQ ID NO:307), LIAFATTPD (SEQ ID NO:308),

LIAFATTPDG (SEQ ID NO:309), YAS, YAST (SEQ ID NO:312), YASTA (SEQ ID NO:313), YASTAD (SEQ ID NO:314), YASTADG (SEQ ID NO:315), AYAS (SEQ ID NO:316), AYAST (SEQ ID NO:317), AYASTA (SEQ ID NO:318), AYASTAD (SEQ ID NO:319), AYASTADG (SEQ ID NO:320), IAYAS (SEQ ID NO:321), IAYAST (SEQ ID NO:322), IAYASTA (SEQ ID NO:323), IAYASTAD (SEQ ID NO:324), IAYASTADG (SEQ ID NO:325), LIAYAS (SEQ ID NO:326), LIAYAST (SEQ ID NO:327), LIAYASTA (SEQ ID NO:328), LIAYASTAD (SEQ ID NO:329), LIAYASTADG (SEQ ID NO:330).

21. A modulating agent according to claim 93, wherein the agent comprises a linear peptide having the sequence N-Ac-LFYIEKDTG-NH<sub>2</sub> (SEQ ID NO:148), N-Ac-LFYVERDTG-NH<sub>2</sub> (SEQ ID NO:149), N-Ac-LFYIERDTG-NH<sub>2</sub> (SEQ ID NO:150), N-Ac-LYGYATTADG-NH<sub>2</sub> (SEQ ID NO:280), N-Ac-LYAYATTADG-NH<sub>2</sub> (SEQ ID NO:281), N-Ac-VYGYATTADG-NH<sub>2</sub> (SEQ ID NO:282), N-Ac-VYAYATTADG-NH<sub>2</sub> (SEQ ID NO:283), N-Ac-IYGYATTADG-NH<sub>2</sub> (SEQ ID NO:284), N-Ac-IYAYATTADG-NH<sub>2</sub> (SEQ ID NO:285), IIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:310), N-Ac-LIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:311) or N-Ac-LIAYASTADG-NH<sub>2</sub> (SEQ ID NO:331).

22. A modulating agent according to claim 20, wherein a desmocollin CAR sequence is present within a cyclic peptide.

23. A modulating agent according to claim 22, wherein the cyclic peptide comprises a sequence selected from the group consisting of CEKDC (SEQ ID NO:3327), CEKDTG (SEQ ID NO:3328), CEKDTGC (SEQ ID NO:3329), CIEKDC (SEQ ID NO:3330), CIEKDTG (SEQ ID NO:3331), CIEKDTGC (SEQ ID NO:3332), CYIEKDC (SEQ ID NO:3333), CYIEKDTG (SEQ ID NO:3334), CYIEKDTGC (SEQ ID NO:3335), CFYIEKDC (SEQ ID NO:3336), CFYIEKDTG (SEQ ID NO:3337), CFYIEKDTGC (SEQ ID NO:3338), CLFYIEKDC (SEQ ID NO:3339), CLFYIEKDTG (SEQ ID NO:3340), CLFYIEKDTGC (SEQ ID NO:3341), CERDC (SEQ ID NO:3342), CERDTG (SEQ ID NO:3343), CERDTGC (SEQ ID NO:3344), CVERDC

(SEQ ID NO:3345), CVERDTC (SEQ ID NO:3346), CVERDTGC (SEQ ID NO:3347), CYVERDC (SEQ ID NO:3348), CYVERDTC (SEQ ID NO:3349), CYVERDTGC (SEQ ID NO:3350), CFYVERDC (SEQ ID NO:3351), CFYVERDTC (SEQ ID NO:3352), CFYVERDTGC (SEQ ID NO:3353), CLFYVERDC (SEQ ID NO:3354), CLFYVERDTC (SEQ ID NO:3355), CLFYVERDTGC (SEQ ID NO:3356), CIERDC (SEQ ID NO:3357), CIERDTC (SEQ ID NO:3358), CIERDTGC (SEQ ID NO:3359), CYIERDC (SEQ ID NO:3360), CYIERDTC (SEQ ID NO:3361), CYIERDTGC (SEQ ID NO:3362), CFYIERDC (SEQ ID NO:3363), CFYIERDTC (SEQ ID NO:3364), CFYIERDTGC (SEQ ID NO:3365), CLFYIERDC (SEQ ID NO:3366), CLFYIERDTC (SEQ ID NO:3367), CLFYIERDTGC (SEQ ID NO:3368), EEKDK (SEQ ID NO:3369), EEKDTK (SEQ ID NO:3370), EEKDTGK (SEQ ID NO:3371), EIEKDK (SEQ ID NO:3372), EIEKDTK (SEQ ID NO:3373), EIEKDTGK (SEQ ID NO:3374), EYIEKDK (SEQ ID NO:3375), EYIEKDTK (SEQ ID NO:3376), EYIEKDTGK (SEQ ID NO:3377), EFYIEKDK (SEQ ID NO:3378), EFYIEKDTK (SEQ ID NO:3379), EFYIEKDTGK (SEQ ID NO:3380), ELFYIEKDK (SEQ ID NO:3381), ELFYIEKDTK (SEQ ID NO:3382), ELFYIEKDTGK (SEQ ID NO:3383), EERDK (SEQ ID NO:3384), EERDTK (SEQ ID NO:3385), EERDTGK (SEQ ID NO:3386), EVERDK (SEQ ID NO:3387), EVERDTK (SEQ ID NO:3388), EVERDTGK (SEQ ID NO:3389), EYVERDK (SEQ ID NO:3390), YVERDTK (SEQ ID NO:3391), EYVERDTGK (SEQ ID NO:3392), EFYVERDK (SEQ ID NO:3393), EFYVERDTK (SEQ ID NO:3394), EFYVERDTGK (SEQ ID NO:3395), ELFYVERDK (SEQ ID NO:3396), ELFYVERDTK (SEQ ID NO:3397), ELFYVERDTGK (SEQ ID NO:3398), EIERDK (SEQ ID NO:3399), EIERDTK (SEQ ID NO:3400), EIERDTGK (SEQ ID NO:3401), EYIERDK (SEQ ID NO:3402), EYIERDTK (SEQ ID NO:3403), EYIERDTGK (SEQ ID NO:3404), EFYIERDK (SEQ ID NO:3405), EFYIERDTK (SEQ ID NO:3406), EFYIERDTGK (SEQ ID NO:3407), ELFYIERDK (SEQ ID NO:3408), ELFYIERDTK (SEQ ID NO:3409), ELFYIERDTGK (SEQ ID NO:3410), KEKDD (SEQ ID NO:3411), KEKDTD (SEQ ID NO:3412), KEKDTGD (SEQ ID NO:3413), KIEKDD (SEQ ID NO:3414), KIEKDTD (SEQ ID NO:3415), KIEKDTGD (SEQ ID NO:3416), KYIEKDD (SEQ ID NO:3417), KYIEKDTD (SEQ ID NO:3418), KYIEKDTGD (SEQ



ID NO:3419), KFYIEKDD (SEQ ID NO:3420), KFYIEKDTD (SEQ ID NO:3421), KFYIEKDTGD (SEQ ID NO:3422), KLFYIEKDD (SEQ ID NO:3423), KLFYIEKDTD (SEQ ID NO:3424), KLFYIEKDTGD (SEQ ID NO:3425), KERDD (SEQ ID NO:3426), KERDTD (SEQ ID NO:3427), KERDTGD (SEQ ID NO:3428), KVERDD (SEQ ID NO:3429), KVERDTD (SEQ ID NO:3430), KVERDTGD (SEQ ID NO:3431), KYVERDD (SEQ ID NO:3432), KYVERDTD (SEQ ID NO:3433), KYVERDTGD (SEQ ID NO:3434), KFYVERDD (SEQ ID NO:3435), KFYVERDTD (SEQ ID NO:3436), KFYVERDTGD (SEQ ID NO:3437), KLFYVERDD (SEQ ID NO:3438), KLFYVERDTD (SEQ ID NO:3439), KLFYVERDTGD (SEQ ID NO:3440), KIERDD (SEQ ID NO:3441), KIERDTD (SEQ ID NO:3442), KIERDTGD (SEQ ID NO:3443), KYIERD (SEQ ID NO:3444), KYIERDTD (SEQ ID NO:3445), KYIERDTGD (SEQ ID NO:3446), KFYIERDD (SEQ ID NO:3447), KFYIERDTD (SEQ ID NO:3448), KFYIERDTGD (SEQ ID NO:3449), KLFYIERDD (SEQ ID NO:3450), KLFYIERDTD (SEQ ID NO:3451), KLFYIERDTGD (SEQ ID NO:3452), DEKDK (SEQ ID NO:3453), DEKDTK (SEQ ID NO:3454), DEKDTGK (SEQ ID NO:3455), DIEKDK (SEQ ID NO:3456), DIEKDTK (SEQ ID NO:3457), DIEKDTGK (SEQ ID NO:3458), DYIEKDK (SEQ ID NO:3459), DYIEKDTK (SEQ ID NO:3460), DYIEKDTGK (SEQ ID NO:3461), DFYIEKDK (SEQ ID NO:3462), DFYIEKDTK (SEQ ID NO:3463), DFYIEKDTGK (SEQ ID NO:3464), DLFYIEKDK (SEQ ID NO:3465), DLFYIEKDTK (SEQ ID NO:3466), DLFYIEKDTGK (SEQ ID NO:3467), DERDK (SEQ ID NO:3468), DERDTK (SEQ ID NO:3469), DERDTGK (SEQ ID NO:3470), DVERDK (SEQ ID NO:3471), DVERDTK (SEQ ID NO:3472), DVERDTGK (SEQ ID NO:3473), DYVERDK (SEQ ID NO:3474), DYVERDTK (SEQ ID NO:3475), DYVERDTGK (SEQ ID NO:3476), DFYVERDK (SEQ ID NO:3477), DFYVERDTK (SEQ ID NO:3478), DFYVERDTGK (SEQ ID NO:3479), DLFYVERDK (SEQ ID NO:3480), DLFYVERDTK (SEQ ID NO:3481), DLFYVERDTGK (SEQ ID NO:3482), DIERDK (SEQ ID NO:3483), DIERDTK (SEQ ID NO:3484), DIERDTGK (SEQ ID NO:3485), DYIERDK (SEQ ID NO:3486), DYIERDTK (SEQ ID NO:3487), DYIERDTGK (SEQ ID NO:3488), DFYIERDK (SEQ ID NO:3489), DFYIERDTK (SEQ ID NO:3490), DFYIERDTGK (SEQ ID

---

NO:3491), DLFYIERDK (SEQ ID NO:3492), DLFYIERDTK (SEQ ID NO:3493), DLFYIERDTGK (SEQ ID NO:3494), KEKDE (SEQ ID NO:3495), KEKDT (SEQ ID NO:3496), KEKDTGE (SEQ ID NO:3497), KIEKDE (SEQ ID NO:3498), KIEKDT (SEQ ID NO:3499), KIEKDTGE (SEQ ID NO:3500), KYIEKDE (SEQ ID NO:3501), KYIEKDT (SEQ ID NO:3502), KYIEKDTGE (SEQ ID NO:3503), KFYIEKDE (SEQ ID NO:3504), KFYIEKDT (SEQ ID NO:3505), KFYIEKDTGE (SEQ ID NO:3506), KLFYIEKDE (SEQ ID NO:3507), KLFYIEKDT (SEQ ID NO:3508), KLFYIEKDTGE (SEQ ID NO:3509), KERDE (SEQ ID NO:3510), KERDT (SEQ ID NO:3511), KERDTGE (SEQ ID NO:3512), KVERDE (SEQ ID NO:3513), KVERDT (SEQ ID NO:3514), KVERDTGE (SEQ ID NO:3515), KYVERDE (SEQ ID NO:3516), KYVERDT (SEQ ID NO:3517), KYVERDTGE (SEQ ID NO:3518), KFYVERDE (SEQ ID NO:3519), KFYVERDT (SEQ ID NO:3520), KFYVERDTGE (SEQ ID NO:3521), KLFYVERDE (SEQ ID NO:3522), KLFYVERDT (SEQ ID NO:3523), KLFYVERDTGE (SEQ ID NO:3524), KIERDE (SEQ ID NO:3525), KIERDT (SEQ ID NO:3526), KIERDTGE (SEQ ID NO:3527), KYIERDE (SEQ ID NO:3528), KYIERDT (SEQ ID NO:3529), KYIERDTGE (SEQ ID NO:3530), KFYIERDE (SEQ ID NO:3531), KFYIERDT (SEQ ID NO:3532), KFYIERDTGE (SEQ ID NO:3533), KLFYIERDE (SEQ ID NO:3534), KLFYIERDT (SEQ ID NO:3535), KLFYIERDTGE (SEQ ID NO:3536), EKDTG (SEQ ID NO:3537), IEKDT (SEQ ID NO:3538), IEKDTG (SEQ ID NO:3539), YIEKD (SEQ ID NO:3540), YIEKDT (SEQ ID NO:3541), YIEKDTG (SEQ ID NO:3542), FYIEKD (SEQ ID NO:3543), FYIEKDT (SEQ ID NO:3544), FYIEKDTG (SEQ ID NO:3545), LFYIEKD (SEQ ID NO:3546), LFYIEKDT (SEQ ID NO:3547), LFYIEKDTG (SEQ ID NO:3548), ERDTG (SEQ ID NO:3549), VERDT (SEQ ID NO:3550), VERDTG (SEQ ID NO:3551), YVERD (SEQ ID NO:3552), YVERDT (SEQ ID NO:3553), YVERDTG (SEQ ID NO:3554), FYVERD (SEQ ID NO:3555), FYVERDT (SEQ ID NO:3556), FYVERDTG (SEQ ID NO:3557), LFYVERD (SEQ ID NO:3558), LFYVERDT (SEQ ID NO:3559), LFYVERDTG (SEQ ID NO:3560), IERDT (SEQ ID NO:3561), IERDTG (SEQ ID NO:3562), YIERD (SEQ ID NO:3563), YIERDT (SEQ ID NO:3564), YIERDTG (SEQ ID NO:3565), FYIERD (SEQ ID NO:3566), FYIERDT

(SEQ ID NO:3567), FYIERDTG (SEQ ID NO:3568), LFYIERD (SEQ ID NO:3569), LFYIERDT (SEQ ID NO:3570), LFYIERDTG (SEQ ID NO:3571), CYATC, CYATTC, CYATTAC, CYATTADC, CYATTADGC, CGYATC, CGYATTC, CGYATTAC, CGYATTADC, CGYATTADGC, CAYATC, CAYATTC, CAYATTAC, CAYATTADC, CAYATTADGC, CYGYATC, CYGYATTC, CYGYATTAC, CYGYATTADC, CYGYATTADGC, CYAYATC, CYAYATTC, CYAYATTAC, CYAYATTADC, CYAYATTADGC, CLYGYATC, CLYGYATTC, CLYGYATTAC, CLYGYATTADC, CLYGYATTADGC, CLYAYATC, CLYAYATTC, CLYAYATTAC, CLYAYATTADC, CLYAYATTADGC, CVYGYATC, CVYGYATTC, CVYGYATTAC, CVYGYATTADC, CVYGYATTADGC, CVYAYATC, CVYAYATTC, CVYAYATTAC, CVYAYATTADC, CVYAYATTADGC, CIYGYATC, CIYGYATTC, CIYGYATTAC, CIYGYATTADC, CIYGYATTADGC, CIYAYATC, CIYAYATTC, CIYAYATTAC, CIYAYATTADC, CIYAYATTADGC, EYATK, EYATTK, EYATTAK, EYATTADK, EYATTADGK, EGYATK, EGYATTK, EGYATTAK, EGYATTADK, EGYATTADGK, EAYATK, EAYATTK, EAYATTAK, EAYATTADK, EAYATTADGK, EYGYATK, EYGYATTK, EYGYATTAK, EYGYATTADK, EYGYATTADGK, EYAYATK, EYAYATTK, EYAYATTAK, EYAYATTADK, EYAYATTADGK, ELYGYATK, ELYGYATTK, ELYGYATTAK, ELYGYATTADK, ELYGYATTADGK, ELYAYATK, ELYAYATTK, ELYAYATTAK, ELYAYATTADK, ELYAYATTADGK, EVYGYATK, EVYGYATTK, EVYGYATTAK, EVYGYATTADK, EVYGYATTADGK, EVYAYATK, EVYAYATTK, EVYAYATTAK, EVYAYATTADK, EVYAYATTADGK, EIYGYATK, EIYGYATTK, EIYGYATTAK, EIYGYATTADK, EIYGYATTADGK, EIYAYATK, EIYAYATTK, EIYAYATTAK, EIYAYATTADK, EIYAYATTADGK, KYATE, KYATTE, KYATTAE, KYATTADE, KYATTADGE, KGYATE, KGYATTE, KGYATTAE, KGYATTADE, KGYATTADGE, KAYATE, KAYATTE, KAYATTAE, KAYATTADE, KAYATTADGE, KYGYATE, KYGYATTE, KYGYATTAE, KYGYATTADE, KYGYATTADGE, KYAYATE, KYAYATTE, KYAYATTAE, KYAYATTADE, KYAYATTADGE, KLYGYATE,

KLYGYATTE, KLYGYATTAE, KLYGYATTADE, KLYGYATTADGE,  
KLYAYATE, KLYAYATTE, KLYAYATTAE, KLYAYATTADE,  
KLYAYATTADGE, KVYGYATE, KVYGYATTE, KVYGYATTAE,  
KVYGYATTADE, KVYGYATTADGE, KVYAYATE, KVYAYATTE,  
KVYAYATTAE, KVYAYATTADE, KVYAYATTADGE, KIYGYATE,  
KIYGYATTE, KIYGYATTAE, KIYGYATTADE, KIYGYATTADGE, KIYAYATE,  
KIYAYATTE, KIYAYATTAE, KIYAYATTADE, KIYAYATTADGE, DYATK,  
DYATTK, DYATTAK, DYATTADK, DYATTADGK, DGYATK, DGYATTK,  
DGYATTAK, DGYATTADK, DGYATTADGK, DAYATK, DAYATTK,  
DAYATTAK, DAYATTADK, DAYATTADGK, DYGYATK, DYGYATTK,  
DYGYATTAK, DYGYATTADK, DYGYATTADGK, DYAYATK, DYAYATTK,  
DYAYATTAK, DYAYATTADK, DYAYATTADGK, DLYGYATK, DLYGYATTK,  
DLYGYATTAK, DLYGYATTADK, DLYGYATTADGK, DLYAYATK,  
DLYAYATTK, DLYAYATTAK, DLYAYATTADK, DLYAYATTADGK,  
DVYGYATK, DVYGYATTK, DVYGYATTAK, DVYGYATTADK,  
DVYGYATTADGK, DVYAYATK, DVYAYATTK, DVYAYATTAK,  
DVYAYATTADK, DVYAYATTADGK, DIYGYATK, DIYGYATTK,  
DIYGYATTAK, DIYGYATTADK, DIYGYATTADGK, DIYAYATK, DIYAYATTK,  
DIYAYATTAK, DIYAYATTADK, DIYAYATTADGK, KYATD, KYATTD,  
KYATTAD, KYATTADD, KYATTADGD, KGYATD, KGYATTD, KGYATTAD,  
KGYATTADD, KGYATTADGD, KAYATD, KAYATTD, KAYATTAD,  
KAYATTADD, KAYATTADGD, KYGYATD, KYGYATTD, KYGYATTAD,  
KYGYATTADD, KYGYATTADGD, KYAYATD, KYAYATTD, KYAYATTAD,  
KYAYATTADD, KYAYATTADGD, KLYGYATD, KLYGYATTD,  
KLYGYATTAD, KLYGYATTADD, KLYGYATTADGD, KLYAYATD,  
KLYAYATTD, KLYAYATTAD, KLYAYATTADD, KLYAYATTADGD,  
KVYGYATD, KVYGYATTD, KVYGYATTAD, KVYGYATTADD,  
KVYGYATTADGD, KVYAYATD, KVYAYATTD, KVYAYATTAD,  
KVYAYATTADD, KVYAYATTADGD, KIYGYATD, KIYGYATTD,  
KIYGYATTAD, KIYGYATTADD, KIYGYATTADGD, KIYAYATD, KIYAYATTD,

KIYAYATTAD, KIYAYATTADD, KIYAYATTADGD, YATTA, YATTAD,  
YATTADG, GYATT, GYATTA, GYATTAD, GYATTADG, AYATT, AYATTA,  
AYATTAD, AYATTADG, YGYAT, YGYATT, YGYATTA, YGYATTAD,  
YGYATTADG, YAYAT, YAYATT, YAYATTA, YAYATTAD, YAYATTADG,  
LYGYAT, LYGYATT, LYGYATTA, LYGYATTAD, LYGYATTADG, LYAYAT,  
LYAYATT, LYAYATTA, LYAYATTAD, LYAYATTADG, VYGYAT, VYGYATT,  
VYGYATTA, VYGYATTAD, VYGYATTADG, VYAYAT, VYAYATT,  
VYAYATTA, VYAYATTAD, VYAYATTADG, IYGYAT, IYGYATT, IYGYATTA,  
IYGYATTAD, IYGYATTADG, IYAYAT, IYAYATT, IYAYATTA, IYAYATTAD,  
IYAYATTADG, CFATC, CFATTC, CFATTPC, CFATTPDC, CFATTPDGC,  
CAFATC, CAFATTC, CAFATTPC, CAFATTPDC, CAFATTPDGC, CIAFATC,  
CIAFATTC, CIAFATTPC, CIAFATTPDC, CIAFATTPDGC, CIIAFATC,  
CIIAFATTC, CIIAFATTPC, CIIAFATTPDC, CIIAFATTPDGC, CLIAFATC,  
CLIAFATTC, CLIAFATTPC, CLIAFATTPDC, CLIAFATTPDGC, EFATK,  
EFATTK, EFATTPK, EFATTPDK, EFATTPDGK, EAFATK, EAFATTK,  
EAFATTPK, EAFATTPDK, EAFATTPDGK, EIAFATK, EIAFATTK, EIAFATTPK,  
EIAFATTPDK, EIAFATTPDGK, EIIAFATK, EIIAFATTK, EIIAFATTPK,  
EIIAFATTPDK, EIIAFATTPDGK, ELIAFATK, ELIAFATTK, ELIAFATTPK,  
ELIAFATTPDK, ELIAFATTPDGK, KFATE, KFATTE, KFATTPE, KFATTPDE,  
KFATTPDGE, KAFATE, KAFATTE, KAFATTPE, KAFATTPDE, KAFATTPDGE,  
KIAFATE, KIAFATTE, KIAFATTPE, KIAFATTPDE, KIAFATTPDGE, KIIAFATE,  
KIIAFATTE, KIIAFATTPE, KIIAFATTPDE, KIIAFATTPDGE, KLIAFATE,  
KLIAFATTE, KLIAFATTPE, KLIAFATTPDE, KLIAFATTPDGE, DFATK,  
DFATTK, DFATTPK, DFATTPDK, DFATTPDGK, DAFATK, DAFATTK,  
DAFATTPK, DAFATTPDK, DAFATTPDGK, DIAFATK, DIAFATTK,  
DIAFATTPK, DIAFATTPDK, DIAFATTPDGK, DIIAFATK, DIIAFATTK,  
DIIAFATTPK, DIIAFATTPDK, DIIAFATTPDGK, DLIAFATK, DLIAFATTK,  
DLIAFATTPK, DLIAFATTPDK, DLIAFATTPDGK, KFATD, KFATTD, KFATTPD,  
KFATTPDD, KFATTPDGD, KAFATD, KAFATTD, KAFATTPD, KAFATTPDD,  
KAFATTPDGD, KIAFATD, KIAFATTD, KIAFATTPD, KIAFATTPDD,

KIAFATTPDGD, KIIAFATD, KIIAFATTD, KIIAFATTPD, KIIAFATTPDD,  
KIIAFATTPDGD, KLIAFATD, KLIAFATTD, KLIAFATTPD, KLIAFATTPDD,  
KLIAFATTPDGD, FATTP, FATTPD, FATTPDG, AFATT, AFATTP, AFATTPD,  
AFATTPDG, IAFAT, IAFATT, IAFATTP, IAFATTPD, IAFATTPDG, IIAFAT,  
IIAFATT, IIAFATTP, IIAFATTPD, IIAFATTPDG, LIAFAT, LIAFATT, LIAFATTP,  
LIAFATTPD, LIAFATTPDG, CYASC, CYASTC, CYASTAC, CYASTADC,  
CYASTADGC, CAYASC, CAYASTC, CAYASTAC, CAYASTADC,  
CAYASTADGC, CIAYASC, CIAYASTC, CIAYASTAC, CIAYASTADC,  
CIAYASTADGC, CLIAYASC, CLIAYASTC, CLIAYASTAC, CLIAYASTADC,  
CLIAYASTADGC, EYASK, EYASTK, EYASTAK, EYASTADK, EYASTADGK,  
EAYASK, EAYASTK, EAYASTAK, EAYASTADK, EAYASTADGK, EIAYASK,  
EIAYASTK, EIAYASTAK, EIAYASTADK, EIAYASTADGK, ELIAYASK,  
ELIAYASTK, ELIAYASTAK, ELIAYASTADK, ELIAYASTADGK, KYASE,  
KYASTE, KYASTAE, KYASTADE, KYASTADGE, KAYASE, KAYASTE,  
KAYASTAE, KAYASTADE, KAYASTADGE, KIAYASE, KIAYASTE,  
KIAYASTAE, KIAYASTADE, KIAYASTADGE, KLIAYASE, KLIAYASTE,  
KLIAYASTAE, KLIAYASTADE, KLIAYASTADGE, DYASK, DYASTK,  
DYASTAK, DYASTADK, DYASTADGK, DAYASK, DAYASTK, DAYASTAK,  
DAYASTADK, DAYASTADGK, DIAYASK, DIAYASTK, DIAYASTAK,  
DIAYASTADK, DIAYASTADGK, DLIAYASK, DLIAYASTK, DLIAYASTAK,  
DLIAYASTADK, DLIAYASTADGK, KYASD, KYASTD, KYASTAD,  
KYASTADD, KYASTADGD, KAYASD, KAYASTD, KAYASTAD, KAYASTADD,  
KAYASTADGD, KIAYASD, KIAYASTD, KIAYASTAD, KIAYASTADD,  
KIAYASTADGD, KLIAYASD, KLIAYASTD, KLIAYASTAD, KLIAYASTADD,  
KLIAYASTADGD, YASTA, YASTAD, YASTADG, AYAST, AYASTA,  
AYASTAD, AYASTADG, IAYAS, IAYAST, IAYASTA, IAYASTAD,  
IAYASTADG, LIAYAS, LIAYAST, LIAYASTA, LIAYASTAD and LIAYASTADG.

24. A modulating agent comprising an antibody or antigen-binding fragment thereof that:

(a) specifically binds to a desmocollin CAR sequence selected from the group consisting of LFYIEKDTG (SEQ ID NO:545), LFYVERDTG (SEQ ID NO:559) and LFYIERDTG (SEQ ID NO:571), N-Ac-LYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-LYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-VYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-VYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-IYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-IYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), IIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-LIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:\_\_) and N-Ac-LIAYASTADG-NH<sub>2</sub> (SEQ ID NO:\_\_); and

(b) modulates a desmocollin-mediated function.

25. A modulating agent according to any one of claims 1-5 or 13-14 linked to a detectable marker.

26. A modulating agent according to any one of claims 1-5 or 13-14 linked to a targeting agent.

27. A modulating agent according to any one of claims 1-5 or 13-14 linked to a support material.

28. A modulating agent according to claim 27, wherein the support material is a polymeric matrix.

29. A modulating agent according to claim 27, wherein the support material is selected from the group consisting of plastic dishes, plastic tubes, sutures, membranes, ultra thin films, bioreactors and microparticles.

30. A modulating agent according to any one of claims 1-5 or 13-14, further comprising one or more of:

(a) a CAR sequence that is specifically recognized by an adhesion molecule other than the desmosomal cadherin; and/or

---

(b) an antibody or antigen-binding fragment thereof that specifically binds to a CAR sequence that is specifically recognized by an adhesion molecule other than the desmosomal cadherin.

31. A modulating agent according to claim 30, wherein the adhesion molecule is selected from the group consisting of classical cadherins, integrins, occludin, claudins, nonclassical cadherins, fibronectin, laminin, claudins and other extracellular matrix proteins.

32. A pharmaceutical composition comprising a modulating agent according to any one of claims 1-5 or 13-14, in combination with a physiologically acceptable carrier.

33. A composition according to claim 32, further comprising a drug.

34. A composition according to claim 32, wherein the modulating agent is present within a sustained-release formulation.

35. A pharmaceutical composition according to claim 32, further comprising a modulator of cell adhesion that comprises one or more of:

(a) a CAR sequence that is specifically recognized by an adhesion molecule other than the desmosomal cadherin; and/or

(b) an antibody or antigen-binding fragment thereof that specifically binds to a CAR sequence that is specifically recognized by an adhesion molecule other than the desmosomal cadherin.

36. A pharmaceutical composition according to claim 35, wherein the adhesion molecule is selected from the group consisting of classical cadherins, integrins, occludin, claudins, nonclassical cadherins, fibronectin, laminin and other extracellular matrix proteins.

---



37. A method for modulating a desmosomal cadherin-mediated function, comprising contacting a desmoglein- and/or desmocollin-expressing cell with a modulating agent according to any one of claims 1-5 or 13-14, and thereby modulating a desmosomal cadherin-mediated function.

38. A method according to claim 37, wherein the cell is selected from the group consisting of epithelial cells, endothelial cells and tumor cells.

39. A method according to claim 37, wherein the modulating agent is present within a pharmaceutical composition, in combination with a physiologically acceptable carrier.

40. A method for modulating cell adhesion, comprising contacting a desmosomal cadherin-expressing cell with a modulating agent according to any one of claims 1-5 or 13-14, and thereby modulating cell adhesion.

41. A method according to claim 40, wherein the cell is selected from the group consisting of epithelial cells, endothelial cells and tumor cells.

42. A method according to claim 40, wherein the modulating agent is present within a pharmaceutical composition, in combination with a physiologically acceptable carrier.

43. A method according to claim 40, wherein cell adhesion is inhibited.

44. A method according to claim 40, wherein cell adhesion is enhanced.

45. A method for inhibiting adhesion of desmosomal cadherin-expressing cells in a mammal, comprising administering to a mammal a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent inhibits desmocollin- and/or desmoglein-mediated cell adhesion.

46. A method according to claim 45, wherein the modulating agent is present within a pharmaceutical composition, in combination with a physiologically acceptable carrier.

47. A method according to claim 45, wherein cell adhesion is inhibited.

48. A method according to claim 45, wherein cell adhesion is enhanced.

49. A method for enhancing the delivery of a drug through the skin of a mammal, comprising contacting epithelial cells of a mammal with a drug and a modulating agent according to any one of claims 1-5 or 13-14, wherein the step of contacting is performed under conditions and for a time sufficient to allow passage of the drug across the epithelial cells, and wherein the modulating agent inhibits desmocollin- and/or desmoglein-mediated cell adhesion.

50. A method according to claim 49, wherein the modulating agent passes into the blood stream of the mammal.

51. A method according to claim 49, wherein the modulating agent is linked to the drug.

52. A method according to claim 49, wherein the step of contacting is performed via a skin patch comprising the modulating agent and the drug.

53. A method for facilitating blood sampling in a mammal, comprising contacting epithelial cells of a mammal with a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent inhibits desmocollin and/or desmoglein-mediated cell adhesion, and wherein the step of contacting is performed under conditions and for a time sufficient to allow passage of one or more blood components across the epithelial cells.

54. A method according to claim 53, wherein the step of contacting is performed via a skin patch comprising the modulating agent.

55. A method according to claim 54, wherein the skin patch further comprises a reagent for detecting a blood component of interest.

56. A method according to claim 53, wherein the epithelial cells are skin cells or are gum cells.

57. A method for enhancing adhesion of desmosomal cadherin-expressing cells, comprising contacting desmocollin- and/or desmoglein-expressing cells with a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent enhances desmocollin- and/or desmoglein-mediated cell adhesion, and wherein the step of contacting is performed under conditions and for a time sufficient to detectably enhance adhesion of the cells.

58. A method according to claim 57, wherein the cell is selected from the group consisting of epithelial cells, endothelial cells and tumor cells.

59. A method according to claim 57, wherein the modulating agent is linked to a support molecule or a solid support.

---

60. A method for facilitating wound healing and/or reducing scar tissue in a mammal, comprising contacting a wound in a mammal with a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent enhances desmosomal cadherin-mediated cell adhesion.

61. A method according to claim 60, wherein the modulating agent is linked to a support molecule or a solid support.

62. A method for enhancing adhesion of foreign tissue implanted within a mammal, comprising contacting a site of implantation of foreign tissue in a mammal with a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent enhances desmosomal cadherin-mediated cell adhesion.

63. A method according to claim 62, wherein the foreign tissue is a skin graft or organ implant.

64. A method according to claim 62, wherein the modulating agent is linked to a support molecule or a solid support.

65. A method for treating an autoimmune blistering disorder in a mammal, comprising administering to a mammal a modulating agent according to any one of claims 1-5 or 13-14, wherein the modulating agent enhances desmosomal cadherin-mediated cell adhesion.

66. A method according to claim 65, wherein the modulating agent is administered topically to a blister.

67. A method according to claim 65, wherein the modulating agent is linked to a support molecule or a solid support.

---

68. A method for identifying a compound capable of modulating a desmosomal cadherin-mediated function, comprising: (a) contacting an antibody or antigen-binding fragment thereof that specifically binds to a desmosomal cadherin CAR sequence with a test compound; and (b) detecting the level of antibody or fragment that binds to the test compound, and therefrom identifying a compound capable of modulating desmosomal cadherin-mediated cell adhesion.

69. A method according to claim 68, wherein the antibody or antigen binding fragment specifically binds to a desmoglein CAR sequence selected from the group consisting of N-Ac-IFVINQKTG-NH<sub>2</sub> (SEQ ID NO:105), N-Ac-MFIINRNTG-NH<sub>2</sub> (SEQ ID NO:106), N-Ac-VFYLNKDTG-NH<sub>2</sub> (SEQ ID NO:107), N-Ac-IYRALNSMG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-IYRALNSLG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-LTGYALDARG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-ITCRALNAQG-NH<sub>2</sub> (SEQ ID NO:\_\_) and N-Ac-ITCRALNALG-NH<sub>2</sub> (SEQ ID NO:\_\_).

70. A method according to claim 68, wherein the antibody or antigen binding fragment specifically binds to a desmocollin CAR sequence selected from the group consisting of LFYIEKDTG (SEQ ID NO:545), LFYVERDTG (SEQ ID NO:559) and LFYIERDTG (SEQ ID NO:571), N-Ac-LYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-LYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-VYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-VYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-IYGYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-IYAYATTADG-NH<sub>2</sub> (SEQ ID NO:\_\_), IIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:\_\_), N-Ac-LIAFATTPDG-NH<sub>2</sub> (SEQ ID NO:\_\_) and N-Ac-LIAYASTADG-NH<sub>2</sub> (SEQ ID NO:\_\_).




FIG 1A 11





FIG. 1M-1T



**FIG. 1U-1AA**








Obcad	EC1	RSKRGWVHWOE	FVIEEXTGSDPVLVGRJLISDSDG----	GHIKYI	LSGEGAG----	TIEVIDDKSGHI	IATKTLDBEERAQ----	YTLMAQAVDRDT	NRPLPEPSEFFI	VKVQDINDNDPPEK																					
Obcad	EC2	---JHET	YUANVPEPS	---IHGYSV	LOVTSADADIP	YTHS	AKLVYS	I	LEQCP----	YFSVDAQTG	I	RTAL	PNMDREAKEE	---YHVV	QAKDMG	---HMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Cad5	EC1	RQKRWMTWQH	IMDEKMTS	PLHIIHQ	IKSSVS	KRN----	---AKY	LI	SGSYG----	KVFRVDAETG	VD	---FAL	KLDREHISE	---YHIT	AV	LV	DKDTG	---EMLET	P	SEFT	IK	VDVNDNDPPEK									
Cad6	EC1	RKRSWMMHOF	IEEXTGSDPYQYVK	KLISDQDGH----	GS	KY	LS	SGSYG----	---PLEITH	TEGDI	---QAT	KLDREBPV	---Y	LRAQ	INIRKT	---GRPV	PESEFT	IK	VDVNDNDPPEK												
Cad6	EC2	---TK	YVATVPEMS	---DVGT	EVQVATADADIP	YTHS	AKVVYS	I	LCQP----	YFSESETG	IK	TA	LMHDBREHQ	---YQVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Cad7	EC1	---SK	AY	I	Q	IEDA	---Q	INTT	SGSVTAQPPAAR	---H	SV	YS	DRHTTHIP	---H	IT	ATE	INN	---PKOSS	NRVPL	IK	VDVNDNDPPEK										
Cad7	EC2	RTKRSWMMHOF	VJUEEXTGSDPYVK	KLISDQDGH----	GS	KY	LS	SGSYG----	---HIF	IDSNGIST	---PT	SK	LDRETLW	---H	IT	ATE	INN	---PKOSS	NRVPL	IK	VDVNDNDPPEK										
Cad7	EC2	---JHET	YUANVPEPS	---IHGYSV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Cad7	EC4	---TSRI	YSHNVSEAA	---KVGT	I	GVVA	AIDP	DAASH	---SPV	YS	DRHTTHIP	---H	IT	ATE	INN	---PKOSS	NRVPL	IK	VDVNDNDPPEK												
Cad8	EC1	ISKRGWVHWOE	FVIEEXTGSDPVLVGRJLISDSDG----	GHIKYI	LSGEGAG----	TIEVIDDKSGHI	IATKTLDBEERAQ----	YTLMAQAVDRDT	NRPLPEPSEFFI	VKVQDINDNDPPEK																					
Cad12	EC1	RVKRGWVHWOE	FVIEEXTGSDPVLVGRJLISDSDG----	GHIKYI	LSGEGAG----	TIEVIDDKSGHI	IATKTLDBEERAQ----	YTLMAQAVDRDT	NRPLPEPSEFFI	VKVQDINDNDPPEK																					
Cad12	EC2	---JHET	YUANVPEPS	---IHGYSV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Cad14	EC1	RPKRGWVHWOE	FVIEEXTGSDPVLVGRJLISDSDG----	GHIKYI	LSGEGAG----	TIEVIDDKSGHI	IATKTLDBEERAQ----	YTLMAQAVDRDT	NRPLPEPSEFFI	VKVQDINDNDPPEK																					
Cad14	EC2	---JHET	YUANVPEPS	---IHGYSV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Cad14	EC4	---TNG	Y	LV	PEMS	---IHG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK							
Cad14	EC4	---SMPS	Y	MEV	Y	PEMS	---IHG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK						
Cad15	EC1	---RVIR	AMV	IP	TS	V	EN	SKIR	KE	Y	LV	PEMS	---IHG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK
Cad15	EC2	---I	Q	EA	FC	TRV	IEGA	---V	PG	Y	Y	TR	AEAT	AD	IP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK	
Phcad	EC1	RYKRWVHWOE	FVIEEXTGSDPVLVGRJLISDSDG----	GHIKYI	LSGEGAG----	TIEVIDDKSGHI	IATKTLDBEERAQ----	YTLMAQAVDRDT	NRPLPEPSEFFI	VKVQDINDNDPPEK																					
Phcad	EC2	---JHET	YUANVPEPS	---IHGYSV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK											
Phcad	EC4	---LQ	SG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK										
Phcad	EC4	---HOS	Y	VAR	Y	PEMS	---IHG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK						
Pcad13	EC5	---E	GS	Y	Y	LV	PEMS	---IHG	YS	SV	LOVTSADADIP	YTHS	AKLVYS	I	LCQP----	YFSEVETG	IK	TA	LMHDBREAKEQ	---YHVV	QAKDMG	---QMGSG	SGT	TKVT	TL	VDVNDNDPPEK					
Pcad16	EC3	---E	GS	Y	Y	LV	PEMS																								

**FIG. 2**

Human Dsg1	EWIKFAAACREGEDNSKRNP IAKIHSDCAN--QQVYRISGVGIDOPPYGIFVINOKTGEINITSIVDREVTPEFIIYCRALNSMGQDLERPLELRVRVLDINDNPPVF
Bull Dsg1	EWIKFAAACREGEDNSKRNP IAKIHSDCAN--QQVYRISGVGIDOPPYGIFVINOKTGEINITSIVDREVTPEFIIYCRALNSMGQDLERPLELRVRVLDINDNPPVF
Human Dsg2	AWITAPVALREGEDLSKKNP IAKIHSDLAEEGLKITYKTKGITEPPFGIFVENKDTGELNVTSLDREETPFLLTGYALDARGNNVEKPLELRVRLVLDINDNEPVF
Human Dsg3	EWVKFAKPCREGEDNSKRNP IAKITSYQAT--QKITRYISGVGIDQPPFGIFVVDKNTGDNITAIVDREETPSFLITCRALNAQGLDVEKPLILTVKILDINDNPPVF
Mouse Dsg3	EWVKFAKPCREGEDNSKRNP IAKITSDFQKN--QKITRYISGVGIDQPPFGIFVVDKNTGDNITAIVDREETPSFLITCRALNAQGLDVEKPLILTVKILDINDNPPVF
Human Dsc1	RWAPIPASLMENSLGPPFQHVQIQSDAAQN--YTI FYSISGPGVDKEPFNLFIYIEKDTGDICTRSIDREKYEQFALYGYATTADGYAPEYPLPLIIKIEDDNDNAPYF
Mouse Dsc1	RWAPICSLMENSLGPPFQHIQQIQSDAAQN--YTI FYSISGPGDDKEPYNLFYIEKDTGDICTRSIDREKYDQFVLVGYATTADGYAPDYPLPLFKVEDDNDNAPYF
Bull Dsc1	RWAPICSLMENSLGPPFQHVQVQSDAAQN--YTI FYSISGPGVDKEPFNLFIYIEKDTGDICTRSIDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Human Dsc2	RWAPICSLMENSLGPPFLFLOQVQSDTAQN--YTI FYSISGPGVDKEPFNLFIYIEKDTGDICTRSIDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Dog Dsc2	RWAPIPCSMQENSLGPPFLFLOQVQSDTAQN--YTI FYSIRGPGVDREPKNLFYVERDTGNLCTRPVDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Human Dsc3	RWAPIPCSMQENSLGPPFLFLOQVQSDAAQN--YTV FYSISGRGVGVDKEPPLNLFYIERDTGNLCTRPVDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Mouse Dsc3	RWAPIPCSMQENSLGPPFLFLOQVQSDAAQN--YTV FYSISGRGVGVDKEPPLNLFYIERDTGNLCTRPVDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Bull Dsc3	RWAPIPCSMQENSLGPPFLFLOQVQSDAAQN--YTI FYSISGRGVGVDKEPPLNLFYIERDTGNLCTRPVDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF
Human Dsc4	RWAPIPCSMQENSLGPPFLFLOQVQSDAAQN--YTV FYSISGRGVGVDKEPPLNLFYIERDTGNLCTRPVDREKYDQFVLVGYATTADGYAPEYPLPLFKVEDDNDNAPYF

FIG. 3

N-Ac-CNRNC-NH<sub>2</sub>N-Ac-CEKDC-NH<sub>2</sub>N-Ac-CERDC-NH<sub>2</sub>N-Ac-CNKDC-NH<sub>2</sub>N-Ac-CNOKC-NH<sub>2</sub>**FIG. 4**

**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**